



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### **Usage guidelines**

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



Library No

George James Drury

March 3<sup>rd</sup> 1906.

5432

ENG  
QA451  
H24  
1964  
TIMOSHENKO  
COLL.

ENGINEERING LIBRARY

Get Duty

Good!

Mar 3rd 1956

C. Sunagar

Form I<sub>2</sub> Central School  
Sheffield







By H. S. HALL.

**Algebraical Examples Supplementary to Hall and Knight's Algebra for Beginners and Elementary Algebra.** (Chaps. I.-XXVII.) With or without Answers. Globe 8vo. 2s.

**Short Introduction to Graphical Algebra** Revised Edition. Gl. 8vo. 1s. [Key, in the press.]

By H. S. HALL AND S. R. KNIGHT.

**Elementary Algebra for Schools** 8s. 6d With Answers. Globe 8vo. 4s. 6d. Key. 8s. 6d.

**Answers to Examples in Elementary Algebra** Fcap. 8vo. Sewed. 1s.

**Algebraical Exercises and Examination Papers.** With or without Answers. Globe 8vo. 2s. 6d.

**Higher Algebra** A Sequel to Elementary Algebra for Schools. Cr. 8vo. 7s. 6d. Key. 10s. 6d.

**Algebra for Beginners** Globe 8vo. 2s. With Answers. 2s. 6d.

**Arithmetical Exercises and Examination Papers** With or without Answers. Globe 8vo. 2s. 6d.

**Elementary Trigonometry** Globe 8vo. 4s. 6d. Key, 8s. 6d.

By H. S. HALL AND F. H. STEVENS.

**A Text-Book of Euclid's Elements.** Containing Books I.-VI., XI., and XII. Props. 1 and 3. Globe 8vo. 4s. 6d. Or separately

Book I. 1s.

Books I. and II. 1s. 6d.

Books II. and III. 2s.

Books I.-III., 2s. 6d.; sewed, 2s.

Books I.-IV., 3s.; sewed, 2s. 6d.

Books III. and IV. 2s.

Books III.-VI. 8s.

Books IV.-VI. 2s. 6d.

Books V., VI., XI., and XII., 1

Book XI. 1s. [and 3. 2s. 6d.]

**A Key to the Exercises and Examples contained in a Text-Book of Euclid's Elements** Books I.-VI. and XI. 8s. 6d. Books I.-IV. Cr. 8vo. 6s. 6d. Books VI. and XI. 3s. 6d.

**A School Geometry**, based on the recommendations of the Mathematical Association, and on the recent report of the Cambridge Syndicate on Geometry. Gl. 8vo.

Parts I and II. 1s. 6d.

Part III. 1s.

Parts I., II., and III. 2s. 6d.

Parts III. and IV. 1s. 6d.

Part IV. Sewed, 6d.

Parts I. to IV. 8s.

Part V. 1s. 6d.

Parts I to V. 4s. 6d.

Parts IV. and V. 2s.

Part VI. 2s.

**An Elementary Course of Mathematics** Globe 8vo. 2s. 6d.

By F. H. STEVENS.

**Elementary Mensuration** Globe 8vo. 3s. 6d.

**Mensuration for Beginners** Globe 8vo. 1s. 6d.

By H. S. HALL AND R. J. WOOD.

**Algebra for Elementary Schools** Globe 8vo. Parts I., II., and III., 6d. each. Cloth, 8d. each. Answers to each Part, 4d. each. Sewed.

LONDON: MACMILLAN AND CO., LIMITED.

**A TEXT-BOOK OF EUCLID'S ELEMENTS**



A TEXT-BOOK  
OF  
EUCLID'S ELEMENTS

*FOR THE USE OF SCHOOLS*

BOOKS I.-VI. AND XI.

BY

H. S. HALL, M.A.

FORMERLY SCHOLAR OF CHRIST'S COLLEGE, CAMBRIDGE;

AND

F. H. STEVENS, M.A.

FORMERLY SCHOLAR OF QUEEN'S COLLEGE, OXFORD.

*NEW EDITION*

London

MACMILLAN AND CO., LIMITED

NEW YORK: THE MACMILLAN COMPANY

1904

*All rights reserved.*

First Edition, 1888. Second Edition (Book XI. added), 1889.  
Reprinted 1890, 1891, 1892, 1893, 1894, 1895 (twice), 1896, 1897, 1898, 1899.  
New Edition 1900. Reprinted 1901, 1902, 1903, 1904.

GLASGOW: PRINTED AT THE UNIVERSITY PRESS  
BY ROBERT MACLEHOSE AND CO. LTD.

## EXTRACT FROM THE PREFACE TO THE FIRST EDITION.

THIS volume contains the first Six Books and part of the Eleventh Book of Euclid's Elements, together with additional Theorems and Examples, giving the most important elementary developments of Euclidean Geometry.

The text has been carefully revised, and special attention given to those points which experience has shewn to present difficulties to beginners.

In the course of this revision the enunciations have been altered as little as possible: and very few departures have been made from Euclid's proofs; in each case changes have been adopted only where the old text has been generally found a cause of difficulty; and such changes are for the most part in favour of well-recognised alternatives.

In Book I., for example, the ambiguity has been removed from the Enunciations of Propositions 18 and 19, and the fact that Propositions 8 and 26 establish the complete equality of the two triangles considered has been strongly urged: thus the redundant step has been removed from Proposition 34.

In Book II. Simson's arrangement of Proposition 13 has been abandoned for a well-known equivalent.

In Book III. Propositions 35 and 36 have been treated generally, and it has not been thought necessary to do more than call attention in a note to the special cases.

These are the chief deviations from the ordinary text as regards method and arrangement of proof; they are points familiar as difficulties to most teachers, and to name them indicates sufficiently, without further enumeration, the general principles which have guided our revision.



First Edition, 1888. Second Edition (Book XI. added), 1889.  
Reprinted 1890, 1891, 1892, 1898, 1894, 1895 (twice), 1896, 1897, 1898, 1899.  
New Edition 1900. Reprinted 1901, 1902, 1903, 1904.

GLASGOW: PRINTED AT THE UNIVERSITY PRESS  
BY ROBERT MACLEHOSE AND CO. LTD.

## EXTRACT FROM THE PREFACE TO THE FIRST EDITION.

THIS volume contains the first Six Books and part of the Eleventh Book of Euclid's Elements, together with additional Theorems and Examples, giving the most important elementary developments of Euclidean Geometry.

The text has been carefully revised, and special attention given to those points which experience has shewn to present difficulties to beginners.

In the course of this revision the enunciations have been altered as little as possible: and very few departures have been made from Euclid's proofs; in each case changes have been adopted only where the old text has been generally found a cause of difficulty; and such changes are for the most part in favour of well-recognised alternatives.

In Book I., for example, the ambiguity has been removed from the Enunciations of Propositions 18 and 19, and the fact that Propositions 8 and 26 establish the complete equality of the two triangles considered has been strongly urged: thus the redundant step has been removed from Proposition 34.

In Book II. Simson's arrangement of Proposition 13 has been abandoned for a well-known equivalent.

In Book III. Propositions 35 and 36 have been treated generally, and it has not been thought necessary to do more than call attention in a note to the special cases.

These are the chief deviations from the ordinary text as regards method and arrangement of proof; they are points familiar as difficulties to most teachers, and to name them indicates sufficiently, without further enumeration, the general principles which have guided our revision.

A few alternative proofs of difficult propositions are given for the convenience of those teachers who care to use them.

One purpose of the book is *gradually* to familiarise the student with the use of legitimate symbols and abbreviations; for a geometrical argument may thus be thrown into a form which is not only more readily seized by an advanced reader, but is useful as a guide to the way in which Euclid's propositions may be handled in written work. On the other hand, we think it very desirable to defer the introduction of symbols until the beginner has learnt that they can only be properly used in Pure Geometry as abbreviations for verbal argument: and we hope thus to prevent the slovenly and inaccurate habits which are very apt to arise from their employment before this principle is fully recognised.

Accordingly in Book I. we have used no contractions or symbols of any kind, though we have introduced verbal alterations into the text wherever it appeared that conciseness or clearness would be gained.

In Book II. abbreviated forms of constantly recurring words are used, and the phrases *therefore* and *is equal to* are replaced by the usual symbols.

In the Third and following Books, and in additional matter throughout the whole, we have employed all such signs and abbreviations as we believe to add to the clearness of the reasoning, care being taken that the symbols chosen are compatible with a rigorous geometrical method, and are recognised by the majority of teachers.

If this arrangement should be thought fanciful or wanting in uniformity, we may plead that it is the outcome of long experience in the use of various text-books. For some years, for example, we were accustomed to teach from a symbolical text, but in consequence of the frequent misconceptions and inaccuracies which too great brevity was found to generate among beginners, we were compelled to return to one of the older and unabbreviated editions. The gain to our younger boys was immediate and unmistakeable; but the change has not

been unattended with disadvantage to more advanced students, who on reaching the Third or Fourth Book may not only be safely trusted with a carefully chosen system of abbreviations, but are certainly retarded by the monotonous and lengthy formalities of the old text.

It must be understood that our use of symbols, and the removal of unnecessary verbiage and repetition, by no means implies a desire to secure brevity at all hazards. On the contrary, nothing appears to us more mischievous than an abridgement which is attained by omitting steps, or condensing two or more steps into one. Such uses spring from the pressure of examinations; but an examination is not, or ought not to be, a mere race; and while we wish to indicate generally in the later books how a geometrical argument may be abbreviated for the purposes of written work, we have not attempted to reduce the propositions to the barest skeleton which a lenient Examiner may be supposed to accept. Indeed it does not follow that the form most suitable for the page of a text-book is also best adapted to examination purposes; for the object to be attained in each case is entirely different. The text-book should present the argument in the clearest possible manner to the mind of a reader to whom it is new: the written proposition need only convey to the Examiner the assurance that the proposition has been thoroughly grasped and remembered by the pupil.

From first to last we have kept in mind the undoubted fact that a very small proportion of those who study Elementary Geometry, and study it with profit, are destined to become mathematicians in any special sense; and that, to a large majority of students, Euclid is intended to serve not so much as a first lesson in *mathematical* reasoning, as the first, and sometimes the only, model of formal argument presented in an elementary education.

This consideration has determined not only the full treatment of the earlier Books, but the retention of the formal, if somewhat cumbrous, methods of Euclid in many places where proofs of greater brevity and mathematical elegance are available.

We hope that the additional matter introduced into the book will provide sufficient exercise for pupils whose study of Euclid is preliminary to a mathematical education.

The questions distributed through the text follow very easily from the propositions to which they are attached, and we think that teachers are likely to find in them all that is needed for an average pupil reading the subject for the first time.

The Theorems and Examples at the end of each Book contain questions of a slightly more difficult type : they have been very carefully classified and arranged, and brought into close connection with typical examples worked out either partially or in full ; and it is hoped that this section of the book, on which much thought has been expended, will do something towards removing that extreme want of freedom in solving deductions that is so commonly found even among students who have a good knowledge of the text of Euclid.

To Volumes containing only Books I.-III., or Books I.-IV. an Appendix is added, giving an elementary account of the properties of Pole and Polar, and Radical Axis. In the complete book these subjects, together with a short account of Harmonic Section, Centres of Similitude, and Transversals, appear as *Theorems and Examples on Book VI.*

Throughout the book we have italicised those deductions on which we desired to lay special stress as being in themselves important geometrical results : this arrangement we think will be useful to teachers who have little time to devote to riders, or who wish to sketch out a suitable course for revision.

H. S. HALL.  
F. H. STEVENS.

CLIFTON, *December*, 1886.

## PREFATORY NOTE TO THE NEW EDITION.

In the present edition the text has received further revision, and the notes have been for the most part re-written, with a view to greater clearness and simplicity.

References to the *Definitions* being frequent in the text of Book I., the convenience of a *standard order* has been pointed out to us by many elementary teachers. We have therefore thought it advisable to re-number the Definitions in accordance with Simson's edition. This has involved the insertion of certain definitions hitherto omitted as of slight importance: such insertions have now been printed in subordinate type.

A few typographical improvements have been introduced: notably the italicising of *Particular Enunciations*. Some changes in pagination have also been effected for the purpose of presenting the whole of a proposition at one view, or of bringing notes and exercises into closer connection with the text to which they refer. Further, the symbols " $\therefore$ " for *therefore*, and " $=$ " for *is equal to* are now introduced from the 35th Proposition of Book I.

Groups of Test Questions for Revision have been inserted at various stages. These may be useful to beginners, and suggestive to teachers in framing examination papers, which so often consist of mere monotonous lists of propositions and examples.

One important change has been made. The algebraical treatment of the subject-matter of Book V. has been entirely separated from the stricter general treatment, so as to present in the simplest form such Definitions and Theorems of Proportion as are necessary before entering upon Book VI. This Introduction will be found immediately preceding Book VI. in a chapter called *The Elementary Principles of Proportion*.

H. S. H.

F. H. S.

February, 1900.



# CONTENTS

## BOOK I.

	PAGE
DEFINITIONS, POSTULATES, AXIOMS . . . . .	1
SECTION I. PROPOSITIONS 1-26 . . . . .	12
SECTION II. PARALLELS AND PARALLELOGRAMS.	
PROPOSITIONS 27-34 . . . . .	56
SECTION III. THE AREAS OF PARALLELOGRAMS AND TRIANGLES.	
PROPOSITIONS 35-48 . . . . .	72

### *Theorems and Examples on Book I.*

ANALYSIS, SYNTHESIS . . . . .	95
I. ON THE IDENTICAL EQUALITY OF TRIANGLES . . . . .	98
II. ON INEQUALITIES . . . . .	101
III. ON PARALLELS . . . . .	103
IV. ON PARALLELOGRAMS . . . . .	104
V. MISCELLANEOUS THEOREMS AND EXAMPLES . . . . .	108
VI. ON THE CONCURRENCE OF STRAIGHT LINES IN A TRIANGLE . . . . .	110
VII. ON THE CONSTRUCTION OF TRIANGLES WITH GIVEN PARTS . . . . .	115
VIII. ON AREAS . . . . .	117
IX. ON LOCI . . . . .	122
X. ON THE INTERSECTION OF LOCI . . . . .	125



## BOOK II.

	PAGE
DEFINITIONS, ETC. . . . .	128
PROPOSITIONS 1-14. . . . .	130
THEOREMS AND EXAMPLES ON BOOK II. . . . .	158

## BOOK III.

DEFINITIONS, ETC. . . . .	163
PROPOSITIONS 1-37 . . . . .	167
NOTE ON THE METHOD OF LIMITS AS APPLIED TO TANGENCY . . . . .	231

*Theorems and Examples on Book III.*

I. ON THE CENTRE AND CHORDS OF A CIRCLE . . . . .	233
II. ON THE TANGENT AND THE CONTACT OF CIRCLES. The Common Tangent to Two Circles, Problems on Tangency, Orthogonal Circles . . . . .	235
III. ON ANGLES IN SEGMENTS, AND ANGLES AT THE CENTRES AND CIRCUMFERENCES OF CIRCLES. The Orthocentre of a Triangle, and Properties of the Pedal Triangle, Loci, Simson's Line . . . . .	240
IV. ON THE CIRCLE IN CONNECTION WITH RECTANGLES. Further Problems on Tangency . . . . .	251
V. ON MAXIMA AND MINIMA . . . . .	257
VI. HARDER MISCELLANEOUS EXAMPLES . . . . .	264

## BOOK IV.

DEFINITIONS, . . . . .	268
PROPOSITIONS 1-16. . . . .	269
NOTE ON REGULAR POLYGONS . . . . .	294

*Theorems and Examples on Book IV.*

I. ON THE TRIANGLE AND ITS CIRCLES. Circumscribed, Inscribed, and Escribed Circles, The Nine-Points Circle . . . . .	297
II. MISCELLANEOUS EXAMPLES . . . . .	303

CONTENTS.

xiii

BOOK V.

	PAGE
DEFINITIONS, . . . . .	305
PROPOSITIONS 1-16, . . . . .	309

*Elementary Principles of Proportion.*

INTRODUCTION TO BOOK VI. . . . .	317
----------------------------------	-----

BOOK VI.

DEFINITIONS . . . . .	325
PROPOSITIONS 1-D. . . . .	326

*Theorems and Examples on Book VI.*

I. ON HARMONIC SECTION . . . . .	384
II. ON CENTRES OF SIMILARITY AND SIMILITUDE . . . . .	388
III. ON POLE AND POLAR . . . . .	390
IV. ON THE RADICAL AXIS OF TWO OR MORE CIRCLES . . . . .	396
V. ON TRANSVERSALS . . . . .	399
VI. MISCELLANEOUS EXAMPLES ON BOOK VI. . . . .	403

BOOK XI.

DEFINITIONS . . . . .	409
PROPOSITIONS 1-21 . . . . .	419
EXERCISES ON BOOK XI. . . . .	444
THEOREMS AND EXAMPLES ON BOOK XI. . . . .	446

BOOK XII.

\* \* \* *The First Proposition of Book XII. will be found at p. 364, worked out as an Example on Proposition 20 of Book VI., of which it is a development. Prop. 3 of Book XII. is briefly treated as a Corollary to Prop. 1.*



## EUCLID'S ELEMENTS.

### BOOK I.

#### DEFINITIONS.\*

1. A **point** is that which has position, but no magnitude.
2. A **line** is that which has length without breadth.
3. The extremities of a line are points, and the intersection of two lines is a point.
4. A **straight line** is that which lies evenly between its extreme points.  
Any portion cut off from a straight line is called a **segment** of it.
5. A **surface** (or superficies) is that which has length and breadth, but no thickness.

---

6. The boundaries of a surface are lines.

7. A **plane surface** is one in which any two points being taken, the straight line between them lies wholly in that surface.


A plane surface is frequently referred to simply as a *plane*.

**NOTE.** Euclid regards a point merely as a *mark of position*, and he therefore attaches to it no idea of size and shape.

Similarly he considers that the properties of a line arise only from its *length* and *position*, without reference to that minute breadth which every line must really have if *actually drawn*, even though the most perfect instruments are used.


The definition of a surface is to be understood in a similar way.

8. A **plane angle** is the inclination of two lines to one another, which meet together, but are not in the same direction.





[Definition 8 is not required in Euclid's Geometry, the only angles employed by him being those formed by *straight* lines. See Def. 9.]

9. A **plane rectilinear angle** is the inclination of two *straight* lines to one another, which meet together, but are not in the same straight line.




The point at which the straight lines meet is called the **vertex** of the angle, and the straight lines themselves the **arms** of the angle.

NOTE. When there are several angles at one point, each is expressed by three letters, of which the letter that refers to the vertex is put between the other two. Thus the angle contained by the straight lines OA, OB is named the angle AOB or BOA; and the angle contained by OA, OC is named the angle AOC or COA. But if there is only one angle at a point, it may be expressed by a single letter, as *the angle at O*.



Of the two straight lines OB, OC shewn in the adjoining diagram, we recognize that OC is *more inclined* than OB to the straight line OA: this we express by saying that the angle AOC is greater than the angle AOB. Thus an angle must be regarded as having *magnitude*.




It must be carefully observed that the size of an angle in no way depends on the length of its arms, but only on their *inclination* to one another.

The angle AOC is the *sum* of the angles AOB and BOC; and AOB is the *difference* of the angles AOC and BOC.

[Another view of an angle is recognized in many branches of mathematics; and though not employed by Euclid, it is here given because it furnishes more clearly than any other a conception of what is meant by the *magnitude* of an angle.


Suppose that the straight line OP in the diagram is capable of revolution about the point O, like the hand of a watch, but in the opposite direction; and suppose that in this way it has passed successively from the position OA to the positions occupied by OB and OC. Such a line must have *undergone more turning* in passing from OA to




OC, than in passing from OA to OB; and consequently the angle AOC is said to be greater than the angle AOB.]

Angles which lie on either side of a common arm are called **adjacent angles**.

For example, when one straight line OC is drawn from a point in another straight line AB, the angles COA, COB are *adjacent*.




When two straight lines, such as AB, CD, cross one another at E, the two angles CEA, BED are said to be **vertically opposite**. The two angles CEB, AED are also vertically opposite to one another.



2


10. When a straight line standing on another straight line makes the adjacent angles equal to one another, each of the angles is called a **right angle**; and the straight line which stands on the other is called a **perpendicular** to it.




11. An **obtuse angle** is an angle which is greater than a right angle.



12. An **acute angle** is an angle which is less than a right angle.




[In the adjoining figure the straight line OB may be supposed to have arrived at its present position, from the position occupied by OA, by revolution about the point O in either of the two directions indicated by the arrows: thus two straight lines drawn from a point may be considered as forming *two* angles (marked (i) and (ii) in the figure), of which the greater (ii) is said to be **reflex**.



3

If the arms OA, OB are in the same straight line, the angle formed by them on either side is called a **straight angle**.]



13. A term or boundary is the extremity of anything.


14. Any portion of a plane surface bounded by one or more lines is called a **plane figure**.



The sum of the bounding lines is called the **perimeter** of the figure.  
Two figures are said to be equal in **area** when they enclose equal portions of a plane surface.

3.

15. A **circle** is a plane figure contained by one line, which is called the **circumference**, and is such that all straight lines drawn from a certain point within the figure to the circumference are equal to one another; this point is called the **centre** of the circle.



16. A **radius** of a circle is a straight line drawn from the centre to the circumference.

17. A **diameter** of a circle is a straight line drawn through the centre, and terminated both ways by the circumference.

18. A **semicircle** is the figure bounded by a diameter of a circle and the part of the circumference cut off by the diameter.



4.

19. A **segment of a circle** is the figure bounded by a straight line and the part of the circumference which it cuts off.



20. **Rectilinear figures** are those which are bounded by straight lines.


21. A **triangle** is a plane figure bounded by *three* straight lines.




Any one of the angular points of a triangle may be regarded as its **vertex**; and the opposite side is then called the **base**.

22. A **quadrilateral** is a plane figure bounded by *four* straight lines.

The straight line which joins opposite angular points in a quadrilateral is called a **diagonal**.



23. A **polygon** is a plane figure bounded by more than four straight lines.



5

## TRIANGLES.


24. An **equilateral triangle** is a triangle whose three sides are equal.



25. An **isosceles triangle** is a triangle two of whose sides are equal.



26. A **scalene triangle** is a triangle which has three unequal sides.



27. A **right-angled triangle** is a triangle which has a right angle.



The side opposite to the right angle in a right-angled triangle is called the **hypotenuse**.

28. An **obtuse-angled triangle** is a triangle which has an obtuse angle.



29. An **acute-angled triangle** is a triangle which has *three* acute angles.



[It will be seen hereafter (Book I. Proposition 17) that every triangle must have at least two acute angles.]

6



## QUADRILATERALS.

30. A **square** is a four-sided figure which has all its sides equal and all its angles right angles.

[It may be shewn that if a quadrilateral has all its sides equal and *one* angle a right angle, then *all* its angles will be right angles.]



31. An **oblong** is a four-sided figure which has all its angles right angles, but not all its sides equal.

32. A **rhombus** is a four-sided figure which has all its sides equal, but its angles are not right angles.




33. A **rhomboid** is a four-sided figure which has its opposite sides equal to one another, but all its sides are not equal nor its angles right angles.

34. All other four-sided figures are called **trapeziums**.


It is usual now to restrict the term *trapezium* to a quadrilateral which has two of its sides *parallel*. [See Def. 35.]




35. **Parallel straight lines** are such as, being in the same plane, do not meet, however far they are produced in either direction.



36. A **Parallelogram** is a four-sided figure which has its opposite sides parallel.



37. A **rectangle** is a parallelogram which has one of its angles a right angle.



## THE POSTULATES.

Let it be granted,

1. *That a straight line may be drawn from any one point to any other point.*

2. *That a finite, that is to say a terminated, straight line may be produced to any length in that straight line.*

3. *That a circle may be described from any centre, at any distance from that centre, that is, with a radius equal to any finite straight line drawn from the centre.*

## NOTES ON THE POSTULATES.

1. In order to draw the diagrams required in Euclid's Geometry certain instruments are necessary. These are

(i) A ruler with which to draw straight lines.

(ii) A pair of compasses with which to draw circles.

In the *Postulates*, or requests, Euclid claims the use of these instruments, and assumes that they suffice for the purposes mentioned above.

2. It is important to notice that the Postulates include no means of *direct measurement*: hence the straight ruler is not supposed to be *graduated*; and the compasses are not to be employed for *transferring distances* from one part of a diagram to another.

3. When we draw a straight line from the point A to the point B, we are said to *join* AB.

To *produce* a straight line means to *prolong* or *lengthen* it.

The expression *to describe* is used in Geometry in the sense of *to draw*.

## ON THE AXIOMS.

The science of Geometry is based upon certain simple statements, the truth of which is so evident that they are accepted without proof.

These self-evident truths, called by Euclid *Common Notions*, are known as the **Axioms**.

## GENERAL AXIOMS.


1. *Things which are equal to the same thing are equal to one another.*
2. *If equals be added to equals, the wholes are equal.*
3. *If equals be taken from equals, the remainders are equal.*
4. *If equals be added to unequals, the wholes are unequal, the greater sum being that which includes the greater of the unequals.*
5. *If equals be taken from unequals, the remainders are unequal, the greater remainder being that which is left from the greater of the unequals.*
6. *Things which are double of the same thing, or of equal things, are equal to one another.*
7. *Things which are halves of the same thing, or of equal things, are equal to one another.*
- 9.\* *The whole is greater than its part.*

\* To preserve the classification of general and geometrical axioms, we have placed Euclid's *ninth* axiom before the *eighth*.

## GEOMETRICAL AXIOMS.

8. *Magnitudes which can be made to coincide with one another, are equal.*
10. *Two straight lines cannot enclose a space.*
11. *All right angles are equal.*
12. *If a straight line meet two straight lines so as to make the interior angles on one side of it together less than two right angles, these straight lines will meet if continually produced on the side on which are the angles which are together less than two right angles.*

That is to say, if the two straight lines AB and CD are met by the straight line EH at F and G, in such a way that the angles BFG, DGF are together less than two right angles, it is asserted that AB and CD will meet if continually produced in the direction of B and D.



NOTES ON THE AXIOMS.

1. The necessary characteristics of an Axiom are

- (i) That it should be *self-evident*; that is, that its truth should be immediately accepted without proof.
- (ii) That it should be *fundamental*; that is, that its truth should not be derivable from any other truth more simple than itself.
- (iii) That it should supply a basis for the establishment of further truths.

These characteristics may be summed up in the following definition.

DEFINITION. An **Axiom** is a self-evident truth, which neither requires nor is capable of proof, but which serves as a foundation for future reasoning.

2. Euclid's Axioms may be classified as *general* and *geometrical*.

General Axioms apply to *magnitudes of all kinds*. Geometrical Axioms refer specially to *geometrical magnitudes*, as lines, angles, and figures.

3. Axiom 8 is Euclid's test of the equality of two geometrical magnitudes. It implies that any line, angle, or figure, may be taken up from its position, and without change in size or form, laid down upon a second line, angle, or figure, for the purpose of comparison, and it states that two such magnitudes are equal when one can be exactly placed over the other without overlapping.

This process is called **superposition**, and the first magnitude is said to be **applied** to the other.

4. Axiom 12 has been objected to on the double ground that it cannot be considered self-evident, and that its truth may be deduced from simpler principles. It is employed for the first time in the 29th Proposition of Book I., where a short discussion of the difficulty will be found.

## INTRODUCTORY.

1. Little is known of Euclid beyond the fact that he lived about three centuries before Christ (325-285) at Alexandria, where he became famous as a writer and teacher of Mathematics.

Among the works ascribed to him, the best known and most important is *The Elements*, written in Greek, and consisting of Thirteen Books. Of these it is now usual to read Books I.-IV. and VI. (which deal with Plane Geometry), together with parts of Books XI. and XII. (on the Geometry of Solids). The remaining Books deal with subjects which belong to the theory of Arithmetic.

2. Plane Geometry deals with the properties of all lines and figures that may be drawn upon a plane surface.

Euclid in his first Six Books confines himself to the properties of straight lines, rectilineal figures, and circles.

3. The subject is divided into a number of separate discussions, called **propositions**.

Propositions are of two kinds, **Problems** and **Theorems**.

A **Problem** proposes to perform some geometrical construction, such as to draw some particular line, or to construct some required figure.

A **Theorem** proposes to prove the truth of some geometrical statement.

4. A Proposition consists of the following parts :

The *General Enunciation*, the *Particular Enunciation*, the *Construction*, and the *Proof*.

(i) The **General Enunciation** is a preliminary statement, describing in general terms the purpose of the proposition.

(ii) The **Particular Enunciation** repeats in special terms the statement already made, and refers it to a diagram, which enables the reader to follow the reasoning more easily.

(iii) The **Construction** then directs the drawing of such straight lines and circles as may be required to effect the purpose of a problem, or to prove the truth of a theorem.

(iv) The **Proof** shews that the object proposed in a problem has been accomplished, or that the property stated in a theorem is true.

5. Euclid's reasoning is said to be **Deductive**, because by a connected chain of argument it *deduces* new truths from truths already proved or admitted. Thus each proposition, though in one sense complete in itself, is derived from the Postulates, Axioms, or former propositions, and itself leads up to subsequent propositions.

6. The initial letters Q.E.F., placed at the end of a problem, stand for **Quod erat Faciendum**, *which was to be done*.

The letters Q.E.D. are appended to a theorem, and stand for **Quod erat Demonstrandum**, *which was to be proved*.

7. A **Corollary** is a statement the truth of which follows readily from an established proposition; it is therefore appended to the proposition as an inference or deduction, which usually requires no further proof.

8. The attention of the beginner is drawn to the special use of the *future tense* in the Particular Enunciations of Euclid's propositions.

The future is only used in a statement of which the truth is *about to be proved*. Thus: "*The triangle ABC SHALL BE equilateral*" means that the triangle *has yet to be proved* equilateral. While. "*The triangle ABC IS equilateral*" means that the triangle *has already been proved* (or given) equilateral.

9. The following symbols and abbreviations may be employed in writing out the propositions of Book I., though their use is not recommended to beginners.


$\therefore$	for	therefore,	par <sup>l</sup> (or   )	for	parallel,
=	„	is, or are, equal to,	par <sup>m</sup>	„	parallelogram,
$\sphericalangle$	„	angle,	sq.	„	square,
rt. $\sphericalangle$	„	right angle,	rectil.	„	rectilineal,
$\Delta$	„	triangle,	st. line	„	straight line,
perp.	„	perpendicular,	pt.	„	point;

and all obvious contractions of words, such as opp., adj., diag., etc., for opposite, adjacent, diagonal, etc.

## SECTION I.

## PROPOSITION 1. PROBLEM.

*To describe an equilateral triangle on a given finite straight line.*



Let  $AB$  be the given straight line.

*It is required to describe an equilateral triangle on  $AB$ .*

**Construction.** With centre  $A$ , and radius  $AB$ , describe the circle  $BCD$ . *Post. 3.*

With centre  $B$ , and radius  $BA$ , describe the circle  $ACE$ . *Post. 3.*

From the point  $C$  at which the circles cut one another, draw the straight lines  $CA$  and  $CB$  to the points  $A$  and  $B$ . *Post. 1.*

*Then shall the triangle  $ABC$  be equilateral.*

**Proof.** Because  $A$  is the centre of the circle  $BCD$ , therefore  $AC$  is equal to  $AB$ . *Def. 15.*

And because  $B$  is the centre of the circle  $ACE$ , therefore  $BC$  is equal to  $AB$ . *Def. 15.*

Therefore  $AC$  and  $BC$  are each equal to  $AB$ .

But things which are equal to the same thing are equal to one another. *Ax. 1.*


Therefore  $AC$  is equal to  $BC$ .

Therefore  $AC$ ,  $AB$ ,  $BC$  are equal to one another.

Therefore the triangle  $ABC$  is equilateral;  
and it is described on the given straight line  $AB$ . **Q.E.F.**

## PROPOSITION 2. PROBLEM.

*From a given point to draw a straight line equal to a given straight line.*



Let A be the given point, and BC the given straight line.  
*It is required to draw from A a straight line equal to BC.*

**Construction.** Join AB; *Post. 1.*  
 and on AB describe an equilateral triangle DAB. *I. 1.*  
 With centre B, and radius BC, describe the circle CGH. *Post. 3.*  
 Produce DB to meet the circle CGH at G. *Post. 2.*  
 With centre D, and radius DG, describe the circle GKF.  
 Produce DA to meet the circle GKF at F. *Post. 2.*  
*Then AF shall be equal to BC.*


**Proof.** Because B is the centre of the circle CGH,  
 therefore BC is equal to BG. *Def. 15.*  
 And because D is the centre of the circle GKF,  
 therefore DF is equal to DG. *Def. 15.*  
 And DA, a part of DF, is equal to DB, a part of DG; *Def. 24.*  
 therefore the remainder AF is equal to the remainder BG. *Ax. 3.*  
 But BC has been proved equal to BG;  
 therefore AF and BC are each equal to BG.  
 And things which are equal to the same thing are equal  
 to one another. *Ax. 1.*

Therefore AF is equal to BC;  
 and it has been drawn from the given point A. *Q.E.F.*



## PROPOSITION 3. PROBLEM.

*From the greater of two given straight lines to cut off a part equal to the less.*



Let  $AB$  and  $C$  be the two given straight lines, of which  $AB$  is the greater.

*It is required to cut off from  $AB$  a part equal to  $C$ .*

**Construction.** From the point  $A$  draw the straight line  $AD$  equal to  $C$ ; I. 2.  
and with centre  $A$  and radius  $AD$ , describe the circle  $DEF$ ,  
cutting  $AB$  at  $E$ . Post 3.

*Then  $AE$  shall be equal to  $C$ .*

**Proof.** Because  $A$  is the centre of the circle  $DEF$ ,  
therefore  $AE$  is equal to  $AD$ . Def. 15.

But  $C$  is equal to  $AD$ . Constr.

Therefore  $AE$  and  $C$  are each equal to  $AD$ .

Therefore  $AE$  is equal to  $C$ ; Ax. 1.

and it has been cut off from the given straight line  $AB$ .

Q.E.F.

## EXERCISES ON PROPOSITIONS 1 TO 3.

1. If the two circles in Proposition 1 cut one another again at F, prove that AFB is an equilateral triangle.

2. If the two circles in Proposition 1 cut one another at C and F, prove that the figure ACBF is a rhombus.

3. AB is a straight line of given length: shew how to draw from A a line double the length of AB.

4. Two circles are drawn with the same centre O, and two radii OA, OB are drawn in the smaller circle. If OA, OB are produced to cut the outer circle at D and E, prove that AD = BE.

5. AB is a straight line, and P, Q are two points, one on each side of AB. Shew how to find points in AB, whose distance from P is equal to PQ. How many such points will there be?

6. In the figure of Proposition 2, if AB is equal to BC, shew that D, the vertex of the equilateral triangle, will fall on the circumference of the circle CGH.

7. In Proposition 2 the point A may be joined to either extremity of BC. Draw the figure, and prove the proposition in the case when A is joined to C.

8. On a given straight line AB describe an isosceles triangle having each of its equal sides equal to a given straight line PQ.

9. On a given base describe an isosceles triangle having each of its equal sides double of the base.

10. In a given straight line the points A, M, N, B are taken in order. On AB describe a triangle ABC, such that the side AC may be equal to AN, and the side BC to BM.


## NOTE ON PROPOSITIONS 2 AND 3.

Propositions 2 and 3 are rendered necessary by the restriction tacitly imposed by Euclid, that compasses shall not be used to transfer distances. [See Notes on the Postulates.]

In carrying out the construction of Prop. 2 the point A may be joined to *either* extremity of the line BC; the equilateral triangle may be described on *either* side of the line so drawn; and the sides of the equilateral triangle may be produced in *either* direction. Thus there are in general  $2 \times 2 \times 2$ , or *eight*, possible constructions. The student should exercise himself in drawing the various figures that may arise.

## PROPOSITION 4. THEOREM.

If two triangles have two sides of the one equal to two sides of the other, each to each, and have also the angles contained by those sides equal, then the triangles shall be equal in all respects; that is to say, their bases or third sides shall be equal, and their remaining angles shall be equal, each to each, namely those to which the equal sides are opposite; and the triangles shall be equal in area.



Let  $ABC$ ,  $DEF$  be two triangles, in which  
the side  $AB$  is equal to the side  $DE$ ,  
the side  $AC$  is equal to the side  $DF$ , and  
the contained angle  $BAC$  is equal to the contained angle  $EDF$ .

Then (i) the base  $BC$  shall be equal to the base  $EF$ ;  
(ii) the angle  $ABC$  shall be equal to the angle  $DEF$ ;  
(iii) the angle  $ACB$  shall be equal to the angle  $DFE$ ;  
(iv) the triangle  $ABC$  shall be equal to the triangle  $DEF$  in area.

**Proof.** If the triangle  $ABC$  be applied to the triangle  $DEF$ ,  
so that the point  $A$  may lie on the point  $D$ ,  
and the straight line  $AB$  along the straight line  $DE$ ;  
then because  $AB$  is equal to  $DE$ , *Hyp.*  
therefore the point  $B$  must coincide with the point  $E$ .  
And because  $AC$  falls along  $DF$ ,  
and the angle  $BAC$  is equal to the angle  $EDF$ , *Hyp.*  
therefore  $AC$  must fall along  $DF$ .  
And because  $AC$  is equal to  $DF$ , *Hyp.*  
therefore the point  $C$  must coincide with the point  $F$ .  
Then since  $B$  coincides with  $E$ , and  $C$  with  $F$ ,  
therefore the base  $BC$  must coincide with the base  $EF$ ;

for if not, two straight lines would enclose a space; which is impossible. *Ax. 10.*

Thus the base BC coincides with the base EF, and is therefore equal to it. *Ax. 8.*

And the remaining angles of the triangle ABC coincide with the remaining angles of the triangle DEF, and are therefore equal to them;

namely, the angle ABC is equal to the angle DEF,  
and the angle ACB is equal to the angle DFE.

And the triangle ABC coincides with the triangle DEF, and is therefore equal to it in area. *Ax. 8.*

That is, the triangles are equal in all respects. *Q.E.D.*

**NOTE.** The sides and angles of a triangle are known as its *six parts*. A triangle may also be considered in regard to its *area*.

Two triangles are said to be **equal in all respects**, or **identically equal**, when the sides and angles of one are respectively equal to the sides and angles of the other. We have seen that such triangles may be made to *coincide* with one another by *superposition*, so that they are also equal in *area*. [See Note on Axiom 8.]

[It will be shewn later that triangles can be equal in *area* without being equal in their several parts; that is to say, triangles can have the same *area* without having the same *shape*.]

#### EXERCISES ON PROPOSITION 4.

1. ABCD is a square: prove that the diagonals AC, BD are equal to one another.

2. ABCD is a square, and L, M, and N are the middle points of AB, BC, and CD: prove that

(i)  $LM = MN$ . (ii)  $AM = DM$ .

(iii)  $AN = AM$ . (iv)  $BN = DM$ .


[Draw a separate figure in each case.]

3. ABC is an isosceles triangle: from the equal sides AB, AC two equal parts AX, AY are cut off, and BY and CX are joined. Prove that  $BY = CX$ .

4. ABCD is a quadrilateral having the opposite sides BC, AD equal, and also the angle BCD equal to the angle ADC: prove that BD is equal to AC.

## PROPOSITION 5. THEOREM.

The angles at the base of an isosceles triangle are equal to one another ; and if the equal sides be produced, the angles on the other side of the base shall also be equal to one another.



Let  $ABC$  be an isosceles triangle, in which  
the side  $AB$  is equal to the side  $AC$ ,  
and let the straight lines  $AB, AC$  be produced to  $D$  and  $E$ .

Then (i) the angle  $ABC$  shall be equal to the angle  $ACB$  ;  
(ii) the angle  $CBD$  shall be equal to the angle  $BCE$ .

**Construction.** In  $BD$  take any point  $F$  ;  
and from  $AE$  cut off a part  $AG$  equal to  $AF$ . I. 3.  
Join  $FC, GB$ .

**Proof.** Then in the triangles  $FAC, GAB$ ,  
Because  $\left\{ \begin{array}{l} \text{FA is equal to GA,} \\ \text{and AC is equal to AB,} \\ \text{also the contained angle at A is common to the} \\ \text{two triangles :} \end{array} \right.$  *Constr.*  
*Hyp.*  
therefore the triangle  $FAC$  is equal to the triangle  $GAB$  in  
all respects ; I. 4.

that is, the base  $FC$  is equal to the base  $GB$ ,  
and the angle  $ACF$  is equal to the angle  $ABG$ ,  
also the angle  $AFC$  is equal to the angle  $AGB$ .


Again, because  $AF$  is equal to  $AG$ ,  
and  $AB$ , a part of  $AF$ , is equal to  $AC$ , a part of  $AG$  ; *Hyp.*  
therefore the remainder  $BF$  is equal to the remainder  $CG$ .

Then in the two triangles BFC, CGB,  
 Because  $\left\{ \begin{array}{l} \text{BF is equal to CG,} \\ \text{and FC is equal to GB,} \\ \text{also the contained angle BFC is equal to the} \\ \text{contained angle CGB,} \end{array} \right. \begin{array}{l} \textit{Proved.} \\ \textit{Proved.} \\ \textit{Proved.} \end{array}$   
 therefore the triangle BFC is equal to the triangle CGB in  
 all respects ; I. 4.  
 so that the angle FBC is equal to the angle GCB,  
 and the angle BCF to the angle CBG.

Now it has been shewn that the angle ABG is equal to the  
 angle ACF,  
 and that the angle CBG, a part of ABG, is equal to the angle  
 BCF, a part of ACF ;  
 therefore the remaining angle ABC is equal to the remain-  
 ing angle ACB ; Ax. 3.  
 and these are the angles at the base of the triangle ABC.  
 Also it has been shewn that the angle FBC is equal to the  
 angle GCB ;  
 and these are the angles on the other side of the base. Q.E.D.


**COROLLARY.** Hence if a triangle is equilateral it is also  
 equiangular.

**NOTE.** The difficulty which be-  
 ginners find with this proposition  
 arises from the fact that the triangles  
 to be compared overlap one another  
 in the diagram. This difficulty may  
 be diminished by detaching each  
 pair of triangles from the rest of the  
 figure, as shewn in the margin.



## PROPOSITION 6. THEOREM.

*If two angles of a triangle be equal to one another, then the sides also which subtend, or are opposite to, the equal angles, shall be equal to one another.*



Let  $ABC$  be a triangle, in which  
the angle  $ABC$  is equal to the angle  $ACB$ .  
Then shall the side  $AC$  be equal to the side  $AB$ .

**Construction.** For if  $AC$  be not equal to  $AB$ ,  
one of them must be greater than the other.

If possible, let  $AB$  be the greater ;  
and from it cut off  $BD$  equal to  $AC$ .

I. 3.

Join  $DC$ .

**Proof.** Then in the triangles  $DBC$ ,  $ACB$ ,  
Because  $\left\{ \begin{array}{l} DB \text{ is equal to } AC, \\ \text{and } BC \text{ is common to both,} \\ \text{also the contained angle } DBC \text{ is equal to the} \\ \text{contained angle } ACB ; \end{array} \right. \begin{array}{l} \text{Constr.} \\ \text{Hyp.} \end{array}$   
therefore the triangle  $DBC$  is equal to the triangle  $ACB$   
in area, I. 4.  
the part equal to the whole ; which is absurd. Ax. 9.

Therefore  $AB$  is not unequal to  $AC$  ;  
that is,  $AB$  is equal to  $AC$ .

Q.E.D.

**COROLLARY.** Hence if a triangle is equiangular it is also equilateral.

## NOTE ON PROPOSITIONS 5 AND 6.

The enunciation of a theorem consists of two clauses. The first clause tells us what we are to *assume*, and is called the **hypothesis**; the second tells us what *it is required to prove*, and is called the **conclusion**.

For example, the enunciation of Proposition 5 assumes that in a certain triangle ABC *the side AB = the side AC*: this is the *hypothesis*. From this it is required to prove that *the angle ABC = the angle ACB*: this is the *conclusion*.

If we interchange the hypothesis and conclusion of a theorem, we enunciate a new theorem which is called the **converse** of the first.

For example, in Prop. 5

it is *assumed* that  $AB = AC$  ;  
it is *required to prove* that the angle  $ABC =$  the angle  $ACB$ .

Now in Prop. 6

it is *assumed* that the angle  $ABC =$  the angle  $ACB$  ; }  
it is *required to prove* that  $AB = AC$ . }

Thus we see that Prop. 6 is the converse of Prop. 5 ; for the *hypothesis of each is the conclusion of the other*.

In Proposition 6 Euclid employs for the first time an *indirect method of proof* frequently used in geometry. It consists in shewing that the theorem *cannot be untrue* ; since, if it were, we should be led to some *impossible conclusion*. This form of proof is known as **Reductio ad Absurdum**, and is most commonly used in demonstrating the converse of some foregoing theorem.

The converse of *all* true theorems are not themselves necessarily true. [See Note on Prop 8.]


## EXERCISES ON PROPOSITION 5.

1. ABCD is a rhombus, in which the diagonal BD is drawn : shew that
  - (i) the angle  $ABD =$  the angle  $ADB$  ;
  - (ii) the angle  $CBD =$  the angle  $CDB$  ;
  - (iii) the angle  $ABC =$  the angle  $ADC$ .
2. ABC, DBC are two isosceles triangles drawn on the same base BC, but on *opposite* sides of it: prove (by means of 1. 5) that the angle  $ABD =$  the angle  $ACD$ .
3. ABC, DBC are two isosceles triangles drawn on the same base BC and on *the same* side of it: employ 1. 5 to prove that the angle  $ABD =$  the angle  $ACD$ .



## PROPOSITION 7. THEOREM.

On the same base, and on the same side of it, there cannot be two triangles having their sides which are terminated at one extremity of the base equal to one another, and likewise those which are terminated at the other extremity equal to one another.



If it be possible, on the same base AB, and on the same side of it, let there be two triangles ACB, ADB in which the side AC is equal to the side AD, and also the side BC is equal to the side BD.

CASE I. When the vertex of each triangle is without the other triangle.


Construction. Join CD.

Proof. Then in the triangle ACD, because AC is equal to AD, *Hyp.* therefore the angle ACD is equal to the angle ADC. I. 5.

But the whole angle ACD is greater than its part, the angle BCD ; therefore also the angle ADC is greater than the angle BCD ; still more then is the angle BDC greater than the angle BCD.

Again, in the triangle BCD, because BC is equal to BD, *Hyp.* therefore the angle BDC is equal to the angle BCD : I. 5. but it was shewn to be greater ; which is impossible.

CASE II. When one of the vertices, as D, is within the other triangle ACB.



**Construction.** As before, join CD ;  
and produce AC, AD to E and F.

**Proof.** Then in the triangle ACD,  
because AC is equal to AD, *Hyp.*  
therefore the angle ECD is equal to the angle FDC,  
these being the angles on the other side of the base. 1. 5.  
But the angle ECD is greater than its part, the angle BCD ;  
therefore the angle FDC is also greater than the angle  
BCD :  
still more then is the angle BDC greater than the angle  
BCD.

Again, in the triangle BCD,  
because BC is equal to BD, *Hyp.*  
therefore the angle BDC is equal to the angle BCD : 1. 5.  
but it has been shewn to be greater ; which is impossible.


The case in which the vertex of one triangle is on a side of the other needs no demonstration.

Therefore AC cannot be equal to AD, and *at the same time*, BC equal to BD. Q.E.D.

**NOTE.** The sides AC, AD are called **conterminous** sides ; similarly the sides BC, BD are conterminous.

## PROPOSITION 8. THEOREM.

If two triangles have two sides of the one equal to two sides of the other, each to each, and have likewise their bases equal, then the angle which is contained by the two sides of the one shall be equal to the angle which is contained by the two sides of the other



Let  $ABC$ ,  $DEF$  be two triangles, in which  
the side  $AB$  is equal to the side  $DE$ ,  
the side  $AC$  is equal to the side  $DF$ ,  
and the base  $BC$  is equal to the base  $EF$ .

Then shall the angle  $BAC$  be equal to the angle  $EDF$ .

*Proof.* If the triangle  $ABC$  be applied to the triangle  $DEF$ ,  
so that the point  $B$  falls on the point  $E$ ,  
and the base  $BC$  along the base  $EF$ ;  
then because  $BC$  is equal to  $EF$ , *Hyp.*  
therefore the point  $C$  must coincide with the point  $F$ .

Then since  $BC$  coincides with  $EF$ ,  
it follows that  $BA$  and  $AC$  must coincide with  $ED$  and  $DF$  :  
for if they did not, but took some other position, as  $EG$ ,  $GF$ ,  
then on the same base  $EF$ , and on the same side of it, there  
would be two triangles  $EDF$ ,  $EGF$ , having their *conterminous*  
sides equal : namely  $ED$  equal to  $EG$ , and  $FD$  equal to  $FG$ .

But this is impossible.

I. 7.

Therefore the sides  $BA$ ,  $AC$  coincide with the sides  $ED$ ,  $DF$ .  
That is, the angle  $BAC$  coincides with the angle  $EDF$ , and is  
therefore equal to it. *Ax. 8.*

Q.E.D.

NOTE 1. In this Proposition the three sides of one triangle are given equal respectively to the three sides of the other; and from this it is shewn that the two triangles may be made to coincide with one another.

Hence we are led to the following important Corollary.


COROLLARY. *If in two triangles the three sides of the one are equal to the three sides of the other, each to each, then the triangles are equal in all respects.*

[An alternative proof, which is independent of Prop. 7, will be found on page 26.]

NOTE 2. Proposition 8 furnishes an instance of a true theorem of which the *converse* is not necessarily true.

It is proved above that *if the sides of one triangle are severally equal to the sides of another, then the angles of the first triangle are severally equal to the angles of the second.*

The *converse* of this enunciation would be as follows: *If the angles of one triangle are severally equal to the angles of another, then the sides of the first triangle are equal to the sides of the second.*




But this, as the diagram in the margin shews, is by no means necessarily true.

#### EXERCISES ON PROPOSITION 8.

1. Shew (by drawing a diagonal) that the opposite angles of a rhombus are equal.
2. If ABCD is a quadrilateral, in which  $AB = CD$  and  $AD = CB$ , prove that the angle  $\widehat{ADC} =$  the angle  $\widehat{ABC}$ .
3. If ABC and DBC are two isosceles triangles drawn on the same base BC, prove (by means of I. 8) that the angle  $\widehat{ABD} =$  the angle  $\widehat{ACD}$ , taking (i) the case where the triangles are on the *same* side of BC, (ii) the case where they are on *opposite* sides of BC.
4. If ABC, DBC are two isosceles triangles drawn on opposite sides of the same base BC, and if AD be joined, prove that each of the angles BAC, BDC will be divided into two equal parts.
5. If in the figure of Ex. 4 the line AD meets BC in E, prove that  $BE = EC$ .

PROPOSITION 8. ALTERNATIVE PROOF.



Let  $ABC$  and  $DEF$  be two triangles, which have the sides  $BA, AC$  equal respectively to the sides  $ED, DF$ , and the base  $BC$  equal to the base  $EF$ .

*Then shall the angle  $BAC$  be equal to the angle  $EDF$ .*

For apply the triangle  $ABC$  to the triangle  $DEF$ , so that  $B$  may fall on  $E$ , and  $BC$  along  $EF$ , and so that the point  $A$  may be on the side of  $EF$  remote from  $D$ ;

then  $C$  must fall on  $F$ , since  $BC$  is equal to  $EF$ .

Let  $GEF$  be the new position of the triangle  $ABC$ .

Join  $DG$ .

CASE I. When  $DG$  intersects  $EF$ .

Then because  $ED = EG$ ,

$\therefore$  the angle  $EDG =$  the angle  $EGD$ . I. 5.


Again because  $FD = FG$ ,

$\therefore$  the angle  $FDG =$  the angle  $FGD$ . I. 5.


Hence the whole angle  $EDF =$  the whole angle  $EGF$ ; *Ax. 2.*  
that is, the angle  $EDF =$  the angle  $BAC$ .

Two cases remain which may be dealt with in a similar manner : namely,

CASE II. When  $DG$  meets  $EF$  produced.



CASE III. When one pair of sides, as  $DF, FG$  are in one straight line.




QUESTIONS AND EXERCISES FOR REVISION.

1. Define *adjacent angles*, a *right angle*, *vertically opposite angles*.
2. Explain the words *enunciation*, *hypothesis*, *conclusion*.
3. Distinguish between the meanings of the following statements:
  - (i) then AB is equal to PQ ;
  - (ii) then AB shall be equal to PQ.
4. When are two theorems said to be *converse* to one another. Give an example.
5. Shew by an example that the converse of a true theorem is not itself necessarily true.
6. What is a *corollary* ? Quote the corollary to Proposition 5 ; and shew how its truth follows from that proposition.
7. Name the six *parts* of a triangle. When are triangles said to be *equal in all respects* ?
8. What do you understand by the expression *geometrical magnitudes* ? Give examples ?
9. What is meant by *superposition* ? Explain the test by which Euclid determines if two geometrical magnitudes are equal to one another. Illustrate by an example.
10. Quote and explain the *third postulate*. What restrictions does Euclid impose on the use of compasses, and what problems are thereby made necessary ?
11. Define an *axiom*. Quote the axioms referred to (i) in Proposition 2 ; (ii) in Proposition 7.
12. Prove by the method of *superposition* that two squares are equal in area, if a side of one is equal to a side of the other.
13. Two quadrilaterals ABCD, EFGH have the sides AB, BC, CD, DA equal respectively to the sides EF, FG, GH, HE, and have also the angle BAD equal to the angle FEH. Shew that the figures may be made to coincide with one another.
14. AB, AC are the equal sides of an isosceles triangle ABC ; and L, M, N are the middle points of AB, BC, and CA respectively : prove that
  - (i)  $LM = MN$ .                      (ii)  $BN = CL$ .
  - (iii) the angle ALM = the angle ANM.

## PROPOSITION 9. PROBLEM.

To bisect a given rectilinear angle, that is, to divide it into two equal parts.



Let  $BAC$  be the given angle.  
It is required to bisect the angle  $BAC$ .

**Construction.** In  $AB$  take any point  $D$ ;  
and from  $AC$  cut off  $AE$  equal to  $AD$ . I. 3.

Join  $DE$ ;

and on  $DE$ , on the side remote from  $A$ , describe an equilateral triangle  $DEF$ . I. 1.

Join  $AF$ .

Then shall the straight line  $AF$  bisect the angle  $BAC$ .

**Proof.** For in the two triangles  $DAF$ ,  $EAF$ ,

Because {	DA is equal to EA,	<i>Constr.</i>
	and AF is common to both;	
	and the third side DF is equal to the third side EF;	<i>Def. 24.</i>

therefore the angle  $DAF$  is equal to the angle  $EAF$ . I. 8.


Therefore the given angle  $BAC$  is bisected by the straight line  $AF$ . Q.E.F.

## EXERCISES.

1. If in the above figure the equilateral triangle  $DFE$  were described on the same side of  $DE$  as  $A$ , what different cases would arise? And under what circumstances would the construction fail?
2. In the same figure, shew that  $AF$  also bisects the angle  $DFE$ .
3. Divide an angle into four equal parts.

PROPOSITION 10. PROBLEM.

To bisect a given finite straight line, that is, to divide it into two equal parts.



Let AB be the given straight line.  
It is required to divide AB into two equal parts.

**Constr.** On AB describe an equilateral triangle ABC; I. 1.  
and bisect the angle ACB by the straight line CD, meeting  
AB at D. I. 9.

Then shall AB be bisected at the point D.

**Proof.** For in the triangles ACD, BCD,  
AC is equal to BC, Def. 24.  
and CD is common to both;  
Because { also the contained angle ACD is equal to the con-  
tained angle BCD; Constr.  
therefore the triangle ACD is equal to the triangle BCD in  
all respects: I. 4.

so that the base AD is equal to the base BD.

Therefore the straight line AB is bisected at the point D.  
Q.E.F.

EXERCISES.


1. Shew that the straight line which bisects the vertical angle of an isosceles triangle, also bisects the base.

2. On a given base describe an isosceles triangle such that the sum of its equal sides may be equal to a given straight line.



## PROPOSITION 11. PROBLEM.

To draw a straight line at right angles to a given straight line, from a given point in the same.



Let  $AB$  be the given straight line, and  $C$  the given point in it.

It is required to draw from  $C$  a straight line at right angles to  $AB$ .

**Construction.** In  $AC$  take any point  $D$ ,  
and from  $CB$  cut off  $CE$  equal to  $CD$ . I. 3.

On  $DE$  describe the equilateral triangle  $DFE$ . I. 1.

Join  $CF$ .

Then shall  $CF$  be at right angles to  $AB$ .

**Proof.** For in the triangles  $DCF$ ,  $ECF$ ,  
 DC is equal to EC, *Constr.*  
 and  $CF$  is common to both;  
 and the third side  $DF$  is equal to the third side  
 $EF$ : *Def. 24.*  
 therefore the angle  $DCF$  is equal to the angle  $ECF$ : I. 8.  
 and these are adjacent angles.

But when one straight line, standing on another, makes the adjacent angles equal, each of these angles is called a right angle; *Def. 10.*

therefore each of the angles  $DCF$ ,  $ECF$  is a right angle.


Therefore  $CF$  is at right angles to  $AB$ ,  
and has been drawn from a point  $C$  in it. Q.E.F.

## EXERCISE.

In the figure of the above proposition, shew that any point in  $FC$ , or  $FC$  produced, is equidistant from  $D$  and  $E$ .

PROPOSITION 12. PROBLEM.

To draw a straight line perpendicular to a given straight line of unlimited length, from a given point without it.



Let AB be the given straight line of unlimited length, and let C be the given point without it.

It is required to draw from C a straight line perpendicular to AB.

**Construction.** On the side of AB remote from C take any point D ;  
and with centre C, and radius CD, describe the circle FDG, cutting AB at F and G

Bisect FG at H ; 1. 10.  
and join CH.

Then shall CH be perpendicular to AB.  
Join CF and CG.

**Proof.** Then in the triangles FHC, GHC,  
 FH is equal to GH, Constr.  
 and HC is common to both ;  
 and the third side CF is equal to the third side  
 CG, being radii of the circle FDG ; Def. 15.  
 therefore the angle CHF is equal to the angle CHG ; 1. 8.  
 and these are adjacent angles.

But when one straight line, standing on another, makes the adjacent angles equal, each of these angles is called a right angle, and the straight line which stands on the other is called a perpendicular to it. Def. 10.

Therefore CH is perpendicular to AB,  
and has been drawn from the point C without it. Q.E.F.

**NOTE.** The line AB must be of unlimited length, that is, capable of production to an indefinite length in either direction, to ensure its being intersected in two points by the circle FDG.

## QUESTIONS AND EXERCISES FOR REVISION.

1. Distinguish between a *problem* and a *theorem*.
2. When are two figures said to be *identically equal*? Under what conditions has it so far been proved that two *triangles* are *identically equal*?
3. Explain the method of proof known as *Reductio ad Absurdum*. Quote the enunciations of the propositions in which this method has so far been used.
4. Quote the corollaries of Propositions 5 and 6, and shew that each is the converse of the other.
5. What is meant by saying that Euclid's reasoning is *deductive*? Shew, for instance, that the proof of Proposition 5 is a deductive argument.
6. Two forts defend the mouth of a river, one on each side; the forts are 4000 yards apart, and their guns have a range of 3000 yards. Taking *one inch* to represent a length of 1000 yards, draw a diagram shewing what part of the river is exposed to the fire of both forts.
7. Define *the perimeter* of a rectilineal figure. A square and an equilateral triangle each have a perimeter of 3 feet: compare the lengths of their sides.
8. Shew how to draw a rhombus each of whose sides is equal to a given straight line PQ, which is also to be one diagonal of the figure.
9. A and B are two given points. Shew how to draw a rhombus having A and B as opposite vertices, and having each side equal to a given line PQ. Is this always possible?
10. Two circles are described with the same centre O; and two radii OA, OB are drawn to the inner circle, and produced to cut the outer circle at D and E: prove that
  - (i)  $DB = EA$ ;
  - (ii) the angle BAD = the angle ABE;
  - (iii) the angle ODB = the angle OEA.

## EXERCISES ON PROPOSITIONS 1 TO 12.

1. Shew that the straight line which joins the vertex of an isosceles triangle to the middle point of the base is perpendicular to the base.

2. Shew that the straight lines which join the extremities of the base of an isosceles triangle to the middle points of the opposite sides, are equal to one another.

3. Two given points in the base of an isosceles triangle are equidistant from the extremities of the base : shew that they are also equidistant from the vertex.

4. If the opposite sides of a quadrilateral are equal, shew that the opposite angles are also equal.

5. Any two isosceles triangles  $XAB$ ,  $YAB$  stand on the same base  $AB$  : shew that the angle  $XAY$  is equal to the angle  $XB Y$  ; and if  $XY$  be joined, that the angle  $AXY$  is equal to the angle  $BXY$ .

6. Shew that the opposite angles of a rhombus are bisected by the diagonal which joins them.

7. Shew that the straight lines which bisect the base angles of an isosceles triangle form with the base a triangle which is also isosceles.

8.  $ABC$  is an isosceles triangle having  $AB$  equal to  $AC$  ; and the angles at  $B$  and  $C$  are bisected by straight lines which meet at  $O$  : shew that  $OA$  bisects the angle  $BAC$ .

9. Shew that the triangle formed by joining the middle points of the sides of an equilateral triangle is also equilateral.

10. The equal sides  $BA$ ,  $CA$  of an isosceles triangle  $BAC$  are produced beyond the vertex  $A$  to the points  $E$  and  $F$ , so that  $AE$  is equal to  $AF$  ; and  $FB$ ,  $EC$  are joined : shew that  $FB$  is equal to  $EC$ .

11. Shew that the diagonals of a rhombus bisect one another at right angles.

12. In the equal sides  $AB$ ,  $AC$  of an isosceles triangle  $ABC$  two points  $X$  and  $Y$  are taken, so that  $AX$  is equal to  $AY$  ; and  $CX$  and  $BY$  are drawn intersecting in  $O$  : shew that


- (i) the triangle  $BOC$  is isosceles ;
- (ii)  $AO$  bisects the vertical angle  $BAC$  ;
- (iii)  $AO$ , if produced, bisects  $BC$  at right angles.

13. Describe an isosceles triangle, having given the base and the length of the perpendicular drawn from the vertex to the base.

14. In a given straight line find a point that is equidistant from two given points. In what case is this impossible ?

## PROPOSITION 13. THEOREM.

The adjacent angles which one straight line makes with another straight line, on one side of it, are either two right angles, or are together equal to two right angles.



Let the straight line AB meet the straight line DC.

Then the adjacent angles DBA, ABC shall be either two right angles, or together equal to two right angles.

CASE I. For if the angle DBA is equal to the angle ABC, each of them is a right angle. *Def. 10.*

CASE II. But if the angle DBA is not equal to the angle ABC,

from B draw BE at right angles to CD. *I. 11.*

*Proof.* Now the angle DBA is made up of the two angles DBE, EBA;

to each of these equals add the angle ABC;  
then the two angles DBA, ABC are together equal to the three angles DBE, EBA, ABC. *Ax. 2.*

Again, the angle EBC is made up of the two angles EBA, ABC;

to each of these equals add the angle DBE;  
then the two angles DBE, EBC are together equal to the three angles DBE, EBA, ABC. *Ax. 2.*

But the two angles DBA, ABC have been shewn to be equal to the same three angles;

therefore the angles DBA, ABC are together equal to the angles DBE, EBC. *Ax. 1.*

But the angles DBE, EBC are two right angles; *Constr.*  
therefore the angles DBA, ABC are together equal to two right angles. *Q.E.D.*

## DEFINITIONS.

(1) The **complement** of an acute angle is its *defect from a right angle*, that is, the angle by which it falls short of a right angle.

Thus two angles are **complementary**, when their sum is a right angle.

(ii) The **supplement** of an angle is its *defect from two right angles*, that is, the angle by which it falls short of two right angles.

Thus two angles are **supplementary**, when their sum is two right angles.

**COROLLARY.** *Angles which are complementary or supplementary to the same angle are equal to one another.*

## EXERCISES.


1. If the two exterior angles formed by producing a side of a triangle both ways are equal, shew that the triangle is isosceles.

2. *The bisectors of the adjacent angles which one straight line makes with another contain a right angle.*

**NOTE** In the adjoining diagram  $\text{AOB}$  is a given angle; and one of its arms  $\text{AO}$  is produced to  $\text{C}$ : the adjacent angles  $\text{AOB}$ ,  $\text{BOC}$  are bisected by  $\text{OX}$ ,  $\text{OY}$ .

Then  $\text{OX}$  and  $\text{OY}$  are called respectively the **internal** and **external bisectors** of the angle  $\text{AOB}$ .

Hence Exercise 2 may be thus enunciated:




*The internal and external bisectors of an angle are at right angles to one another.*

3. Shew that the angles  $\text{AOX}$  and  $\text{COY}$  are complementary.

4. Shew that the angles  $\text{BOX}$  and  $\text{COX}$  are supplementary; and also that the angles  $\text{AOY}$  and  $\text{BOY}$  are supplementary.

## PROPOSITION 14. THEOREM.

If, at a point in a straight line, two other straight lines, on opposite sides of it, make the adjacent angles together equal to two right angles, then these two straight lines shall be in one and the same straight line.



At the point B in the straight line AB, let the two straight lines BC, BD, on the opposite sides of AB, make the adjacent angles ABC, ABD together equal to two right angles.

Then BD shall be in the same straight line with BC.

**Proof.** For if BD be not in the same straight line with BC, if possible, let BE be in the same straight line with BC.

Then because AB meets the straight line CBE, therefore the adjacent angles CBA, ABE are together equal to two right angles. I. 13.

But the angles CBA, ABD are also together equal to two right angles. *Hyp.*

Therefore the angles CBA, ABE are together equal to the angles CBA, ABD. *Ax. 11.*

From each of these equals take the common angle CBA; then the remaining angle ABE is equal to the remaining angle ABD; the part equal to the whole; which is impossible.

Therefore BE is not in the same straight line with BC.

And in the same way it may be shewn that no other line but BD can be in the same straight line with BC.


Therefore BD is in the same straight line with BC. Q.E.D.

## EXERCISE.

ABCD is a rhombus; and the diagonal AC is bisected at O. If O is joined to the angular points B and D; shew that OB and OD are in one straight line.

## PROPOSITION 15. THEOREM.

*If two straight lines intersect one another, then the vertically opposite angles shall be equal.*



Let the two straight lines AB, CD cut one another at the point E.

Then (i) the angle AEC shall be equal to the angle DEB ;  
 (ii) the angle CEB shall be equal to the angle AED.

**Proof.** Because AE meets the straight line CD, therefore the adjacent angles CEA, AED are together equal to two right angles. I. 13.

Again, because DE meets the straight line AB, therefore the adjacent angles AED, DEB are together equal to two right angles. I. 13.

Therefore the angles CEA, AED are together equal to the angles AED, DEB.

From each of these equals take the common angle AED ; then the remaining angle CEA is equal to the remaining angle DEB. Ax. 3.

In the same way it may be proved that the angle CEB is equal to the angle AED. Q.E.D.


**COROLLARY 1.** *From this it follows that, if two straight lines cut one another, the four angles so formed are together equal to four right angles.*

**COROLLARY 2.** *Consequently, when any number of straight lines meet at a point, the sum of the angles made by consecutive lines is equal to four right angles.*



## PROPOSITION 16. THEOREM.

*If one side of a triangle be produced, then the exterior angle shall be greater than either of the interior opposite angles.*



Let  $ABC$  be a triangle, and let  $BC$  be produced to  $D$ .

*Then shall the exterior angle  $ACD$  be greater than either of the interior opposite angles  $ABC$ ,  $BAC$ .*

**Construction.** Bisect  $AC$  at  $E$ ; I. 10.  
Join  $BE$ ; and produce it to  $F$ , making  $EF$  equal to  $BE$ . I. 3.  
Join  $FC$ .

**Proof.** Then in the triangles  $AEB$ ,  $CEF$ ,  
 Because  $\left\{ \begin{array}{l} AE \text{ is equal to } CE, \\ \text{and } EB \text{ is equal to } EF; \\ \text{also the angle } AEB \text{ is equal to the vertically} \\ \text{opposite angle } CEF; \end{array} \right. \begin{array}{l} \text{Constr.} \\ \text{Constr.} \\ \text{I. 15.} \end{array}$   
 therefore the triangle  $AEB$  is equal to the triangle  $CEF$  in  
 all respects: I. 4.  
 so that the angle  $BAE$  is equal to the angle  $ECF$ .

But the angle  $ECD$  is greater than its part, the angle  $ECF$ ;  
 therefore the angle  $ECD$  is greater than the angle  $BAE$ ;  
 that is, the angle  $ACD$  is greater than the angle  $BAC$ .


In the same way, if  $BC$  be bisected, and the side  $AC$   
 produced to  $G$ , it may be proved that the angle  $BCG$  is  
 greater than the angle  $ABC$ .

But the angle  $BCG$  is equal to the angle  $ACD$ : I. 15.  
 therefore also the angle  $ACD$  is greater than the angle  $ABC$ .

Q.E.D.

## PROPOSITION 17. THEOREM.

*Any two angles of a triangle are together less than two right angles.*



Let ABC be a triangle.

*Then shall any two of the angles of the triangle ABC be together less than two right angles.*

**Construction.** Produce the side BC to D.

**Proof.** Then because BC, a side of the triangle ABC, is produced to D ; therefore the exterior angle ACD is greater than the interior opposite angle ABC. I. 16.

To each of these add the angle ACB :

then the angles ACD, ACB are together greater than the angles ABC, ACB. Ax. 4.

But the adjacent angles ACD, ACB are together equal to two right angles. I. 13.

Therefore the angles ABC, ACB are together less than two right angles.

Similarly it may be shewn that the angles BAC, ACB, as also the angles CAB, ABC, are together less than two right angles. Q.E.D.


**NOTE.** It follows from this Proposition that *every triangle must have at least two acute angles* : for if one angle is obtuse, or a right angle, each of the other angles must be less than a right angle.

## EXERCISES.

1. Enunciate this Proposition so as to shew that it is the converse of Axiom 12.
2. If any side of a triangle is produced both ways, the exterior angles so formed are together greater than two right angles.
3. Shew how a proof of Proposition 17 may be obtained by joining each vertex in turn to any point in the opposite side.

## PROPOSITION 18. THEOREM.

*If one side of a triangle be greater than another, then the angle opposite to the greater side shall be greater than the angle opposite to the less.*



Let  $ABC$  be a triangle, in which the side  $AC$  is greater than the side  $AB$ .

*Then shall the angle  $ABC$  be greater than the angle  $ACB$ .*

**Construction.** From  $AC$  cut off a part  $AD$  equal to  $AB$ . I. 3.  
Join  $BD$ .

**Proof.** Then in the triangle  $ABD$ ,  
because  $AB$  is equal to  $AD$ ,  
therefore the angle  $ABD$  is equal to the angle  $ADB$ . I. 5.

But the exterior angle  $ADB$  of the triangle  $DCB$  is greater than the interior opposite angle  $DCB$ , that is, greater than the angle  $ACB$ . I. 16.

Therefore also the angle  $ABD$  is greater than the angle  $ACB$ ; still more then is the angle  $ABC$  greater than the angle  $ACB$ . Q.E.D.


Euclid enunciated Proposition 18 as follows :

*The greater side of every triangle has the greater angle opposite to it.*

[This form of enunciation is found to be a common source of difficulty with beginners, who fail to distinguish what is *assumed* in it and what is *to be proved*. If Euclid's enunciations of Props. 18 and 19 are adopted, it is important to remember that in each case the part of the triangle *first named* points out the hypothesis.]

## PROPOSITION 19. THEOREM.

*If one angle of a triangle be greater than another, then the side opposite to the greater angle shall be greater than the side opposite to the less.*



Let ABC be a triangle in which the angle ABC is greater than the angle ACB.

*Then shall the side AC be greater than the side AB.*

**Proof.** For if AC be not greater than AB, it must be either equal to, or less than AB.

But AC is not equal to AB,  
for then the angle ABC would be equal to the angle ACB; I. 5.  
but it is not. *Hyp.*

Neither is AC less than AB;  
for then the angle ABC would be less than the angle ACB; I. 18.  
but it is not. *Hyp.*

That is, AC is neither equal to, nor less than AB.  
Therefore AC is greater than AB. Q.E.D.

**NOTE.** The mode of demonstration used in this Proposition is known as the **Proof by Exhaustion**. It is applicable to cases in which *one* of certain suppositions must necessarily be true; and it consists in shewing that each of these suppositions is false *with one exception*: hence the truth of the remaining supposition is inferred.


Euclid enunciated Proposition 19 as follows:

*The greater angle of every triangle is subtended by the greater side, or, has the greater side opposite to it.*

[For Exercises on Props. 18 and 19 see page 44.]

## PROPOSITION 20. THEOREM.

*Any two sides of a triangle are together greater than the third side.*



Let ABC be a triangle.

*Then shall any two of its sides be together greater than the third side :*

*namely, BA, AC, shall be greater than CB ;*

*AC, CB shall be greater than BA ;*

*and CB, BA shall be greater than AC.*

**Construction.** Produce BA to D, making AD equal to AC. I. 3.  
Join DC.

**Proof.**

Then in the triangle ADC,

because AD is equal to AC,

*Constr.*

therefore the angle ACD is equal to the angle ADC. I. 5.

But the angle BCD is greater than its part the angle ACD ;

therefore also the angle BCD is greater than the angle ADC,

that is, than the angle BDC.

And in the triangle BCD,

because the angle BCD is greater than the angle BDC,

therefore the side BD is greater than the side CB. I. 19.

But BA and AC are together equal to BD ;

therefore BA and AC are together greater than CB.

Similarly it may be shewn


that AC, CB are together greater than BA ;

and CB, BA are together greater than AC. Q.E.D.

[For Exercises see page 44.]

PROPOSITION 21. THEOREM.

*If from the ends of a side of a triangle, there be drawn two straight lines to a point within the triangle, then these straight lines shall be less than the other two sides of the triangle, but shall contain a greater angle.*



Let ABC be a triangle, and from B, C, the ends of the side BC, let the straight lines BD, CD be drawn to a point D within the triangle

Then (i) BD and DC shall be together less than BA and AC ;  
 (ii) the angle BDC shall be greater than the angle BAC.

**Construction.** Produce BD to meet AC in E.

**Proof** (i) In the triangle BAE, the two sides BA, AE are together greater than the third side BE ; I. 20.  
 to each of these add EC ;

then BA, AC are together greater than BE, EC. Ax. 4.

Again, in the triangle DEC, the two sides DE, EC are together greater than DC ; I. 20.

to each of these add BD ;

then BE, EC are together greater than BD, DC.

But it has been shewn that BA, AC are together greater than BE, EC :

still more then are BA, AC greater than BD, DC.

(ii) Again, the exterior angle BDC of the triangle DEC is greater than the interior opposite angle DEC ; I. 16.

and the exterior angle DEC of the triangle BAE is greater than the interior opposite angle BAE, that is, than the angle BAC ; I. 16.

still more then is the angle BDC greater than the angle BAC.

Q.E.D.

## EXERCISES.

## ON PROPOSITIONS 18 AND 19.

1. The hypotenuse is the greatest side of a right-angled triangle.
2. If two angles of a triangle are equal to one another, the sides also, which subtend the equal angles, are equal to one another. Prove this [*i.e.* Prop. 6] indirectly by using the result of Prop. 18.
3. BC, the base of an isosceles triangle ABC, is produced to any point D; shew that AD is greater than either of the equal sides.
4. If in a quadrilateral the greatest and least sides are opposite to one another, then each of the angles adjacent to the least side is greater than its opposite angle.
5. In a triangle ABC, if AC is not greater than AB, shew that any straight line drawn through the vertex A and terminated by the base BC, is less than AB.
6. ABC is a triangle, in which OB, OC bisect the angles ABC, ACB respectively: shew that, if AB is greater than AC, then OB is greater than OC.

## ON PROPOSITION 20.

7. The difference of any two sides of a triangle is less than the third side.
8. In a quadrilateral, if two opposite sides which are not parallel are produced to meet one another; shew that the perimeter of the greater of the two triangles so formed is greater than the perimeter of the quadrilateral.
9. The sum of the distances of any point from the three angular points of a triangle is greater than half its perimeter.
10. The perimeter of a quadrilateral is greater than the sum of its diagonals.
11. Obtain a proof of Proposition 20 by bisecting an angle by a straight line which meets the opposite side.

## ON PROPOSITION 21.

12. In Proposition 21 shew that the angle BDC is greater than the angle BAC by joining AD, and producing it towards the base.
13. The sum of the distances of any point within a triangle from its angular points is less than the perimeter of the triangle.

QUESTIONS FOR REVISION.

1. Define the *complement* of an angle. When are two angles said to be *supplementary*? Shew that two angles which are supplementary to the same angle are equal to one another.

2. What is meant by an angle being *bisected internally and externally*?

Prove that the internal and external bisectors of an angle are at right angles to one another.

3. Prove that the sum of the angles formed by any number of straight lines drawn from a point is equal to four right angles.

4. Why must every triangle have *at least two acute angles*? Quote the enunciation of the proposition from which this inference is drawn.

5. In the enunciation *The greater side of a triangle has the greater angle opposite to it*, point out what is assumed and what is to be proved.

6. What is meant by the *Proof by Exhaustion*? Illustrate the use of this method by naming the steps in the proof of Proposition 19.

7. What inference may be drawn respecting the triangles whose sides measure

(i) 4 inches, 5 inches, 4 inches;

(ii) 8 inches, 9 inches, 10 inches;

(iii) 6 inches, 10 inches, 4 inches?

8. Quote the enunciations of propositions which, from a hypothesis relating to the *sides* of triangle, establish a conclusion relating to the *angles*.

9. Quote the enunciations of propositions which, from a hypothesis relating to the *angles* of a triangle, establish a conclusion relating to the *sides*.

10. Explain why parallel straight lines must be *in the same plane*.


11. Prove by means of Prop. 7 that on a given base and on the same side of it only one equilateral triangle can be drawn.

12. In an isosceles triangle, if the equal sides are produced, shew that the angles on the other side of the base must be obtuse.



## PROPOSITION 22. PROBLEM.

To describe a triangle having its sides equal to three given straight lines, any two of which are together greater than the third.



Let  $A, B, C$  be the three given straight lines, of which any two are together greater than the third.

It is required to describe a triangle of which the sides shall be equal to  $A, B, C$ .

**Construction.** Take a straight line  $DE$  terminated at the point  $D$ , but unlimited towards  $E$ .

Make  $DF$  equal to  $A$ ,  $FG$  equal to  $B$ , and  $GH$  equal to  $C$ . I. 3.

With centre  $F$  and radius  $FD$ , describe the circle  $DLK$ .

With centre  $G$  and radius  $GH$ , describe the circle  $MHK$  cutting the former circle at  $K$ .

Join  $FK, GK$ .

Then shall the triangle  $KFG$  have its sides equal to the three straight lines  $A, B, C$ .


**Proof.** Because  $F$  is the centre of the circle  $DLK$ ,  
 therefore  $FK$  is equal to  $FD$ : Def. 15.  
 but  $FD$  is equal to  $A$ ; Constr.  
 therefore also  $FK$  is equal to  $A$ . Ax. 1.

Again, because  $G$  is the centre of the circle  $MHK$ ,  
 therefore  $GK$  is equal to  $GH$ : Def. 15.  
 but  $GH$  is equal to  $C$ ; Constr.  
 therefore also  $GK$  is equal to  $C$ . Ax. 1.  
 And  $FG$  is equal to  $B$ . Constr.

Therefore the triangle  $KFG$  has its sides  $KF, FG, GK$  equal respectively to the three given lines  $A, B, C$ . Q.E.F.

PROPOSITION 23. PROBLEM.

*At a given point in a given straight line, to make an angle equal to a given rectilineal angle.*



Let AB be the given straight line, and A the given point in it, and let LCM be the given angle.

*It is required to draw from A a straight line making with AB an angle equal to the given angle DCE.*

**Construction.** In CL, CM take any points D and E ;  
and join DE.

From AB cut off AF equal to CD. I. 3.

On AF describe the triangle FAG, having the remaining sides AG, GF equal respectively to CE, ED. I. 22.

*Then shall the angle FAG be equal to the angle DCE.*

**Proof.** For in the triangles FAG, DCE,  
Because { FA is equal to DC, Constr.  
          { and AG is equal to CE ; Constr.  
          { and the base FG is equal to the base DE: Constr.  
therefore the angle FAG is equal to the angle DCE. I. 8.


That is, AG makes with AB, at the given point A, an angle equal to the given angle DCE. Q.E.F.

EXERCISE.

On a given base describe a triangle, whose remaining sides shall be equal to two given straight lines. Point out how the construction fails, if any one of the three given lines is greater than the sum of the other two.

## PROPOSITION 24. THEOREM.

If two triangles have two sides of the one equal to two sides of the other, each to each, but the angle contained by the two sides of one greater than the angle contained by the corresponding sides of the other; then the base of that which has the greater angle shall be greater than the base of the other.



Let  $ABC$ ,  $DEF$  be two triangles, in which  
the side  $BA$  is equal to the side  $ED$ ,  
and the side  $AC$  is equal to the side  $DF$ ,  
but the angle  $BAC$  is greater than the angle  $EDF$ .  
Then shall the base  $BC$  be greater than the base  $EF$ .

Of the two sides  $DE$ ,  $DF$ , let  $DE$  be that which is not greater than the other.\*

**Construction.** At  $D$  in the straight line  $ED$ , and on the same side of it as  $DF$ , make the angle  $EDG$  equal to the angle  $BAC$ . I. 23.

Make  $DG$  equal to  $DF$  or  $AC$ ; I. 3.  
and join  $EG$ ,  $GF$ .

**Proof.** Then in the triangles  $BAC$ ,  $EDG$ ,  
Because  $\left\{ \begin{array}{l} BA \text{ is equal to } ED, \\ \text{and } AC \text{ is equal to } DG, \\ \text{also the contained angle } BAC \text{ is equal to the} \\ \text{contained angle } EDG; \end{array} \right.$  Hyp.  
Constr.  
Constr.  
therefore the triangle  $BAC$  is equal to the triangle  $EDG$  in  
all respects: I. 4.  
so that the base  $BC$  is equal to the base  $EG$ .

Again, in the triangle FDG,  
 because DG is equal to DF,  
 therefore the angle DFG is equal to the angle DGF. I. 5.

But the angle DGF is greater than its part the angle EGF;  
 therefore also the angle DFG is greater than the angle EGF;  
 still more then is the angle EFG greater than the angle EGF.

And in the triangle EFG,  
 because the angle EFG is greater than the angle EGF,  
 therefore the side EG is greater than the side EF; I. 19.  
 but EG was shewn to be equal to BC;  
 therefore BC is greater than EF. Q.E.D.

\*The object of this step is to make the point F fall *below* EG. Otherwise F might fall *above*, *upon*, or *below* EG; and each case would require separate treatment. But as it is not *proved* that this condition fulfils its object, this demonstration of Prop. 24 must be considered defective.

An alternative construction and proof are given below.

**Construction.** At D in ED make the angle EDG equal to the angle BAC; and make DG equal to DF. Join EG.

Then, as before, it may be shewn that the triangle EDG = the triangle BAC in all respects.


Now if EG passes through F, then EG is greater than EF; that is, BC is greater than EF.

But if not, bisect the angle FDG by DK, meeting EG at K. Join FK.

**Proof.** Then in the triangles FDK, GDK,


Because  $\begin{cases} FD = GD, \\ \text{and } DK \text{ is common to both,} \\ \text{and the angle } FDK = \text{the angle } GDK; \end{cases}$  *Constr.*  
 $\therefore FK = GK.$  I. 4.

But in the triangle EKF, the two sides EK, KF are greater than EF;  
 that is, EK, KG are greater than EF.  
 Hence EG (or BC) is greater than EF.



## PROPOSITION 25. THEOREM.

*If two triangles have two sides of the one equal to two sides of the other, each to each, but the base of one greater than the base of the other ; then the angle contained by the sides of that which has the greater base, shall be greater than the angle contained by the corresponding sides of the other.*



Let  $ABC$ ,  $DEF$  be two triangles in which  
the side  $BA$  is equal to the side  $ED$ ,  
and the side  $AC$  is equal to the side  $DF$ ,  
but the base  $BC$  is greater than the base  $EF$ .  
Then shall the angle  $BAC$  be greater than the angle  $EDF$ .

*Proof.* For if the angle  $BAC$  be not greater than the angle  $EDF$ , it must be either equal to, or less than the angle  $EDF$ .

But the angle  $BAC$  is not equal to the angle  $EDF$ ,  
for then the base  $BC$  would be equal to the base  $EF$ ; I. 4.  
but it is not. *Hyp.*

Neither is the angle  $BAC$  less than the angle  $EDF$ ,  
for then the base  $BC$  would be less than the base  $EF$ ; I. 24.  
but it is not. *Hyp.*

Therefore the angle  $BAC$  is neither equal to, nor less than  
the angle  $EDF$  ;  
that is, the angle  $BAC$  is greater than the angle  $EDF$ . Q.E.D.


## EXERCISE.

In a triangle  $ABC$ , the vertex  $A$  is joined to  $X$ , the middle point of the base  $BC$  ; shew that the angle  $AXB$  is obtuse or acute, according as  $AB$  is greater or less than  $AC$ .

PROPOSITION 26. THEOREM.

If two triangles have two angles of the one equal to two angles of the other, each to each, and a side of one equal to a side of the other, these sides being either adjacent to the equal angles, or opposite to equal angles in each; then shall the triangles be equal in all respects.

CASE I. When the equal sides are adjacent to the equal angles in the two triangles.



Let  $ABC$ ,  $DEF$  be two triangles, in which  
the angle  $ABC$  is equal to the angle  $DEF$ ,  
and the angle  $ACB$  is equal to the angle  $DFE$ ,  
and the side  $BC$  is equal to the side  $EF$ .

Then shall the triangle  $ABC$  be equal to the triangle  $DEF$  in all respects; that is,  $AB$  shall be equal to  $DE$ , and  $AC$  to  $DF$ , and the angle  $BAC$  shall be equal to the angle  $EDF$ .


For if  $AB$  be not equal to  $DE$ , one must be greater than the other. If possible, let  $AB$  be greater than  $DE$ .

**Construction.** From  $BA$  cut off  $BG$  equal to  $ED$ ,  
and join  $GC$ . I. 3.

**Proof.** Then in the two triangles  $GBC$ ,  $DEF$ ,  
Because  $\left\{ \begin{array}{l} \text{GB is equal to DE,} \\ \text{and BC is equal to EF,} \\ \text{also the contained angle GBC is equal to the} \\ \text{contained angle DEF;} \end{array} \right. \begin{array}{l} \text{Constr.} \\ \text{Hyp.} \\ \text{Hyp.} \end{array}$   
therefore the triangle  $GBC$  is equal to the triangle  $DEF$  in all respects; I. 4.

so that the angle  $GCB$  is equal to the angle  $DFE$ .

But the angle  $ACB$  is equal to the angle  $DFE$ ; *Hyp.*  
therefore also the angle  $GCB$  is equal to the angle  $ACB$ ; *Ax. 1.*  
the part equal to the whole, which is impossible.




Therefore AB is not unequal to DE ;  
that is, AB is equal to DE.

Hence in the triangles ABC, DEF,  
AB is equal to DE, *Proved.*  
and BC is equal to EF; *Hyp.*  
Because { also the contained angle ABC is equal to the  
contained angle DEF; *Hyp.*  
therefore the triangle ABC is equal to the triangle DEF in  
all respects : I. 4.  
so that the side AC is equal to the side DF ;  
and the angle BAC is equal to the angle EDF.

Q.E.D.

CASE II. When the equal sides are *opposite* to equal angles in the two triangles.



Let ABC, DEF be two triangles, in which  
the angle ABC is equal to the angle DEF,  
and the angle ACB is equal to the angle DFE,  
and the side AB is equal to the side DE.

Then the triangle ABC shall be equal to the triangle DEF in all respects ;

*namely, BC shall be equal to EF,  
and AC shall be equal to DF,  
and the angle BAC shall be equal to the angle EDF.*

For if BC be not equal to EF, one must be greater than the other. If possible, let BC be greater than EF.

**Construction.** From BC cut off BH equal to EF, and join AH. I. 3.

**Proof.** Then in the triangles ABH, DEF,  
 Because { AB is equal to DE, *Hyp.*  
           and BH is equal to EF, *Constr.*  
           also the contained angle ABH is equal to the  
           contained angle DEF; *Hyp.*  
 therefore the triangle ABH is equal to the triangle DEF in  
 all respects; I. 4.  
 so that the angle AHB is equal to the angle DFE.

But the angle DFE is equal to the angle ACB; *Hyp.*  
 therefore the angle AHB is equal to the angle ACB; *Ax. 1.*  
 that is, an exterior angle of the triangle ACH is equal to an  
 interior opposite angle; which is impossible. I. 16.

Therefore BC is not unequal to EF,  
 that is, BC is equal to EF.

Hence in the triangles ABC, DEF,  
 Because { AB is equal to DE, *Hyp.*  
           and BC is equal to EF; *Proved.*  
           also the contained angle ABC is equal to the  
           contained angle DEF; *Hyp.*  
 therefore the triangle ABC is equal to the triangle DEF in  
 all respects; I. 4.  
 so that the side AC is equal to the side DF,  
 and the angle BAC is equal to the angle EDF.

Q.E.D.

**COROLLARY.** In both cases of this Proposition it is seen that the triangles may be made to coincide with one another; and they are therefore equal in area.



## ON THE IDENTICAL EQUALITY OF TRIANGLES.


Three cases have been already dealt with in Propositions 4, 8, and 26, the results of which may be summarized as follows :

Two triangles are equal in all respects when the following three parts in each are severally equal :


1. Two sides, and the included angle. *Prop. 4.*
2. The three sides. *Prop. 8, Cor.*
3. (a) Two angles, and the adjacent side ;  
(b) Two angles, and a side opposite one of them. } *Prop. 26.*

Two triangles are not, however, necessarily equal in all respects when *any three parts* of one are equal to the corresponding parts of the other. For example

(i) When the *three angles* of one are equal to the *three angles* of the other, each to each, the adjoining diagram shews that the triangles need not be equal in all respects.



(ii) When *two sides and one angle* in one are equal to *two sides and one angle* in the other, the given angles being *opposite* to equal sides, the diagram shews that the triangles need not be equal in all respects.



For it will be seen that if  $AB=DE$ , and  $AC=DF$ , and the angle  $ABC =$  the angle  $DEF$ , then the shorter of the given sides in the triangle  $DEF$  may lie in either of the positions  $DF$  or  $DF'$ .

In cases (i) and (ii) a further condition must be given before we can prove that the two triangles are identically equal.

We observe that in each of the three cases in which two triangles have been proved equal in all respects, namely in Propositions 4, 8, 26, it is shewn that the triangles may be made to *coincide with one another* ; so that they are equal in *area*. Euclid however restricted himself to the use of Prop. 4, when he required to deduce the equality in *area* of two triangles from the equality of certain of their parts. This restriction is now generally abandoned.

## EXERCISES ON PROPOSITIONS 12-26.

1. If  $BX$  and  $CY$ , the bisectors of the angles at the base  $BC$  of an isosceles triangle  $ABC$ , meet the opposite sides in  $X$  and  $Y$ , shew that the triangles  $YBC$ ,  $XCB$  are equal in all respects.

2. Shew that the perpendiculars drawn from the extremities of the base of an isosceles triangle to the opposite sides are equal.

3. *Any point on the bisector of an angle is equidistant from the arms of the angle.*

4. Through  $O$ , the middle point of a straight line  $AB$ , any straight line is drawn, and perpendiculars  $AX$  and  $BY$  are dropped upon it from  $A$  and  $B$ : shew that  $AX$  is equal to  $BY$ .

5. If the bisector of the vertical angle of a triangle is at right angles to the base, the triangle is isosceles.


6. *The perpendicular is the shortest straight line that can be drawn from a given point to a given straight line; and of others, that which is nearer to the perpendicular is less than the more remote; and two, and only two equal straight lines can be drawn from the given point to the given straight line, one on each side of the perpendicular.*

7. *From two given points on the same side of a given straight line, draw two straight lines, which shall meet in the given straight line, and make equal angles with it.*

Let  $AB$  be the given straight line, and  $P$ ,  $Q$  the given points.

It is required to draw from  $P$  and  $Q$  to a point in  $AB$ , two straight lines that shall be equally inclined to  $AB$ .

*Construction.* From  $P$  draw  $PH$  perpendicular to  $AB$ : produce  $PH$  to  $P'$ , making  $HP'$  equal to  $PH$ . Draw  $QP'$ , meeting  $AB$  in  $K$ . Join  $PK$ .



Then  $PK$ ,  $QK$  shall be the required lines. [Supply the proof.]

8. In a given straight line find a point which is equidistant from two given intersecting straight lines. In what case is this impossible?

9. Through a given point draw a straight line such that the perpendiculars drawn to it from two given points may be equal.

In what case is this impossible?


## SECTION II.

## PARALLEL STRAIGHT LINES AND PARALLELOGRAMS.

**DEFINITION.** Parallel straight lines are such as, being in the same plane, do not meet however far they are produced in both directions.

When two straight lines  $AB$ ,  $CD$  are met by a third straight line  $EF$ , *eight* angles are formed, to which for the sake of distinction particular names are given.

Thus in the adjoining figure,  
 1, 2, 7, 8 are called **exterior** angles,  
 3, 4, 5, 6 are called **interior** angles;  
 4 and 6 are said to be **alternate** angles;  
 so also the angles 3 and 5 are alternate to one another.



Of the angles 2 and 6, 2 is referred to as the exterior angle, and 6 as the **interior opposite** angle on the same side of  $EF$ .

2 and 6 are sometimes called **corresponding** angles.

So also, 1 and 5, 7 and 3, 8 and 4 are corresponding angles.

Euclid's treatment of parallel straight lines is based upon his twelfth Axiom, which we here repeat.


**AXIOM 12.** If a straight line cut two straight lines so as to make the two interior angles on the same side of it together less than two right angles, these straight lines, being continually produced, will at length meet on that side on which are the angles which are together less than two right angles.

Thus in the figure given above, if the two angles 3 and 6 are together less than two right angles, it is asserted that  $AB$  and  $CD$  will meet towards  $B$  and  $D$ .

This Axiom is used to establish *i. 29*: some remarks upon it will be found in a note on that Proposition.

PROPOSITION 27. THEOREM.

*If a straight line, falling on two other straight lines, make the alternate angles equal to one another, then these two straight lines shall be parallel.*



Let the straight line  $EF$  cut the two straight lines  $AB$ ,  $CD$  at  $G$  and  $H$ , so as to make the alternate angles  $AGH$ ,  $GHD$  equal to one another.

*Then shall  $AB$  and  $CD$  be parallel.*

**Proof.** For if  $AB$  and  $CD$  be not parallel, they will meet, if produced, either towards  $B$  and  $D$ , or towards  $A$  and  $C$ .

If possible, let  $AB$  and  $CD$ , when produced, meet towards  $B$  and  $D$ , at the point  $K$ .

Then  $KGH$  is a triangle, of which one side  $KG$  is produced to  $A$ ;

therefore the exterior angle  $AGH$  is greater than the interior opposite angle  $GHK$ . I. 16.

But the angle  $AGH$  was given equal to the angle  $GHK$ : *Hyp.* hence the angles  $AGH$  and  $GHK$  are both equal and unequal; which is impossible.


Therefore  $AB$  and  $CD$  cannot meet when produced towards  $B$  and  $D$ .

Similarly it may be shewn that they cannot meet towards  $A$  and  $C$ :

therefore  $AB$  and  $CD$  are parallel.

## PROPOSITION 28. THEOREM.

*If a straight line, falling on two other straight lines, make an exterior angle equal to the interior opposite angle on the same side of the line; or if it make the interior angles on the same side together equal to two right angles, then the two straight lines shall be parallel.*



Let the straight line EF cut the two straight lines AB, CD in G and H: and

*First*, let the exterior angle EGB be equal to the interior opposite angle GHD.

*Then shall AB and CD be parallel.*

**Proof.** Because the angle EGB is equal to the angle GHD; and because the angle EGB is also equal to the vertically opposite angle AGH; I. 15.

therefore the angle AGH is equal to the angle GHD;

but these are alternate angles;

therefore AB and CD are parallel. I. 27.

Q.E.D.

*Secondly*, let the two interior angles BGH, GHD be together equal to two right angles.

*Then shall AB and CD be parallel.*

**Proof.** Because the angles BGH, GHD are together equal to two right angles; *Hyp.*

and because the adjacent angles BGH, AGH are also together equal to two right angles; I. 13.

therefore the angles BGH, AGH are together equal to the two angles BGH, GHD.

From these equals take the common angle BGH:


then the remaining angle AGH is equal to the remaining angle GHD: and these are alternate angles;

therefore AB and CD are parallel. I. 27.

Q.E.D.

## PROPOSITION 29. THEOREM.

If a straight line fall on two parallel straight lines, then it shall make the alternate angles equal to one another, and the exterior angle equal to the interior opposite angle on the same side; and also the two interior angles on the same side equal to two right angles.



Let the straight line EF fall on the parallel straight lines AB, CD.

Then (i) the angle AGH shall be equal to the alternate angle GHD;

(ii) the exterior angle EGB shall be equal to the interior opposite angle GHD;

(iii) the two interior angles BGH, GHD shall be together equal to two right angles.

**Proof.** (i) For if the angle AGH be not equal to the angle GHD, one of them must be greater than the other.

If possible, let the angle AGH be greater than the angle GHD;

add to each the angle BGH:

then the angles AGH, BGH are together greater than the angles BGH, GHD.

But the adjacent angles AGH, BGH are together equal to two right angles;


therefore the angles BGH, GHD are together less than two right angles;

therefore, by *Axiom 12*, AB and CD meet towards B and D.

But they never meet, since they are parallel. *Hyp.*

Therefore the angle AGH is not unequal to the angle GHD: that is, the angle AGH is equal to the alternate angle GHD.

(Over)



(ii) Again, because the angle AGH is equal to the vertically opposite angle EGB ; I. 15.  
and because the angle AGH is equal to the angle GHD ;

*Proved.*

therefore the exterior angle EGB is equal to the interior opposite angle GHD.

(iii) Lastly, the angle EGB is equal to the angle GHD ;

*Proved.*

add to each the angle BGH ;

then the angles EGB, BGH are together equal to the angles BGH, GHD.

But the adjacent angles EGB, BGH are together equal to two right angles : I. 13.

therefore also the two interior angles BGH, GHD are together equal to two right angles. Q.E.D.

#### EXERCISES ON PROPOSITIONS 27, 28, 29.

1. Two straight lines AB, CD bisect one another at O : shew that the straight lines joining AC and BD are parallel. [I. 27.]

2. *Straight lines which are perpendicular to the same straight line are parallel to one another.* [I. 27 or I. 28.]

3. *If a straight line meet two or more parallel straight lines, and is perpendicular to one of them, it is also perpendicular to all the others.* [I. 29.]

4. *If two straight lines are parallel to two other straight lines, each to each, then the angles contained by the first pair are equal respectively to the angles contained by the second pair.* [I. 29.]

NOTE ON THE TWELFTH AXIOM.

Euclid's twelfth Axiom is unsatisfactory as the basis of a theory of parallel straight lines. It cannot be regarded as either simple or self-evident, and it therefore falls short of the essential characteristics of an axiom: nor is the difficulty entirely removed by considering it as a corollary to Proposition 17, of which it is the converse.

Of the many substitutes which have been proposed, we need only notice the following :

**AXIOM.** *Two intersecting straight lines cannot be both parallel to a third straight line.*

This statement is known as **Playfair's Axiom** ; and though it is not altogether free from objection, it is no doubt simpler and more fundamental than that employed by Euclid, and more readily admitted without proof.

Propositions 27 and 28 having been proved in the usual way, the first part of Proposition 29 is then given thus.

PROPOSITION 29. [ALTERNATIVE PROOF.]

*If a straight line fall on two parallel straight lines, then it shall make the alternate angles equal.*

Let the straight line EF meet the two parallel straight lines AB, CD at G and H.

*Then shall the alternate angles AGH, GHD be equal.*

For if the angle AGH is not equal to the angle GHD :

at G in the straight line HG make the angle HGP equal to the angle GHD, and alternate to it. I. 23.

Then PG and CD are parallel. I. 27.


But AB and CD are parallel : *Hyp.*

therefore the two intersecting straight lines AG, PG are both parallel to CD :

which is impossible. *Playfair's Axiom.*

Therefore the angle AGH is not unequal to the angle GHD ; that is, the alternate angles AGH, GHD are equal. Q.E.D.


The second and third parts of the Proposition may then be deduced as in the text ; and Euclid's Axiom 12 follows as a Corollary.





## PROPOSITION 30. THEOREM.

*Straight lines which are parallel to the same straight line are parallel to one another.*



Let the straight lines  $AB$ ,  $CD$  be each parallel to the straight line  $PQ$ .

*Then shall  $AB$  and  $CD$  be parallel to one another.*

**Construction.** Draw any straight line  $EF$  cutting  $AB$ ,  $CD$ , and  $PQ$  in the points  $G$ ,  $H$ , and  $K$ .

**Proof.** Then because  $AB$  and  $PQ$  are parallel, and  $EF$  meets them,

therefore the angle  $AGK$  is equal to the alternate angle  $GKQ$ .  
I. 29.

And because  $CD$  and  $PQ$  are parallel, and  $EF$  meets them, therefore the exterior angle  $GHD$  is equal to the interior opposite angle  $GKQ$ .  
I. 29.

Therefore the angle  $AGH$  is equal to the angle  $GHD$ ;  
and these are alternate angles;

therefore  $AB$  and  $CD$  are parallel.  
I. 27.

Q.E.D.

**NOTE.** If  $PQ$  lies between  $AB$  and  $CD$ , the Proposition may be established in a similar manner, though in this case it scarcely needs proof; for it is inconceivable that two straight lines, which do not meet an intermediate straight line, should meet one another.


The truth of this Proposition may be readily deduced from Playfair's Axiom, of which it is the converse.

For if  $AB$  and  $CD$  were not parallel, they would meet when produced. Then there would be two intersecting straight lines both parallel to a third straight line: which is impossible.

Therefore  $AB$  and  $CD$  never meet; that is, they are parallel.

## PROPOSITION 31. PROBLEM.

To draw a straight line through a given point parallel to a given straight line.



Let A be the given point, and BC the given straight line.  
It is required to draw through A a straight line parallel to BC.

**Construction.** In BC take any point D; and join AD.  
At the point A in DA, make the angle DAE equal to the angle ADC, and alternate to it, I. 23.  
and produce EA to F.

Then shall EF be parallel to BC.

**Proof.** Because the straight line AD, meeting the two straight lines EF, BC, makes the alternate angles EAD, ADC equal;

*Constr.*

therefore EF is parallel to BC; I. 27.

and it has been drawn through the given point A.


Q.E.F.

## EXERCISES.

1. Any straight line drawn parallel to the base of an isosceles triangle makes equal angles with the sides.
2. If from any point in the bisector of an angle a straight line is drawn parallel to either arm of the angle, the triangle thus formed is isosceles.
3. From a given point draw a straight line that shall make with a given straight line an angle equal to a given angle.
4. From X, a point in the base BC of an isosceles triangle ABC, a straight line is drawn at right angles to the base, cutting AB in Y, and CA produced in Z: shew the triangle AYZ is isosceles.
5. If the straight line which bisects an exterior angle of a triangle is parallel to the opposite side, shew that the triangle is isosceles.

## PROPOSITION 32. THEOREM.

*If a side of a triangle be produced, then the exterior angle shall be equal to the sum of the two interior opposite angles ; also the three interior angles of a triangle are together equal to two right angles.*



Let ABC be a triangle, and let one of its sides BC be produced to D.

Then (i) *the exterior angle ACD shall be equal to the sum of the two interior opposite angles CAB, ABC ;*

(ii) *the three interior angles ABC, BCA, CAB shall be together equal to two right angles.*

**Construction.** Through C draw CE parallel to BA. I. 31.

**Proof.** (i) Then because BA and CE are parallel, and AC meets them, therefore the angle ACE is equal to the alternate angle CAB. I. 29.

Again, because BA and CE are parallel, and BD meets them, therefore the exterior angle ECD is equal to the interior opposite angle ABC. I. 29.

Therefore the whole exterior angle ACD is equal to the sum of the two interior opposite angles CAB, ABC.

(ii) Again, since the angle ACD is equal to the sum of the angles CAB, ABC ; *Proved.*

to each of these equals add the angle BCA : then the angles BCA, ACD are together equal to the three angles BCA, CAB, ABC.

But the adjacent angles BCA, ACD are together equal to two right angles. I. 13.

Therefore also the angles BCA, CAB, ABC are together equal to two right angles. Q.E.D.

From this Proposition we draw the following important inferences.

1. *If two triangles have two angles of the one equal to two angles of the other, each to each, then the third angle of the one is equal to the third angle of the other.*
2. *In any right-angled triangle the two acute angles are complementary.*
3. *In a right-angled isosceles triangle each of the equal angles is half a right angle.*
4. *If one angle of a triangle is equal to the sum of the other two, the triangle is right-angled.*
5. *The sum of the angles of any quadrilateral figure is equal to four right angles.*
6. *Each angle of an equilateral triangle is two-thirds of a right angle.*

---

#### EXERCISES ON PROPOSITION 32.

1. Prove that the three angles of a triangle are together equal to two right angles,
  - (i) by drawing through the vertex a straight line parallel to the base ;
  - (ii) by joining the vertex to any point in the base.
2. If the base of any triangle is produced both ways, shew that the sum of the two exterior angles diminished by the vertical angle is equal to two right angles.
3. *If two straight lines are perpendicular to two other straight lines, each to each, the acute angle between the first pair is equal to the acute angle between the second pair.*
4. *Every right-angled triangle is divided into two isosceles triangles by a straight line drawn from the right angle to the middle point of the hypotenuse.*  
*Hence the joining line is equal to half the hypotenuse.*
5. *Draw a straight line at right angles to a given finite straight line from one of its extremities, without producing the given straight line.*

[Let AB be the given straight line. On AB describe any isosceles triangle ACB. Produce BC to D, making CD equal to BC. Join AD. Then shall AD be perpendicular to AB.]


6. *Trisect a right angle.*

7. The angle contained by the bisectors of the angles at the base of an isosceles triangle is equal to an exterior angle formed by producing the base.

8. The angle contained by the bisectors of two adjacent angles of a quadrilateral is equal to half the sum of the remaining angles.

The following theorems were added as corollaries to Proposition 32 by Robert Simson, who edited Euclid's text in 1756.

**COROLLARY 1.** *All the interior angles of any rectilineal figure, together with four right angles, are equal to twice as many right angles as the figure has sides.*



Let ABCDE be any rectilineal figure.

Take F, any point within it,

and join F to each of the angular points of the figure.

Then the figure is divided into as many triangles as it has sides.

And the three angles of each triangle are together equal to two right angles. I. 32.

Hence *all* the angles of *all* the triangles are together equal to twice as many right angles as the figure has sides.

But all the angles of all the triangles make up all the interior angles of the figure, together with the angles at F, which are equal to four right angles. I. 15, *Cor.*

Therefore all the interior angles of the figure, together with four right angles, are equal to twice as many right angles as the figure has sides. Q.E.D.

**COROLLARY 2.** *If the sides of a rectilineal figure, which has no re-entrant angle, are produced in order, then all the exterior angles so formed are together equal to four right angles.*



For at each angular point of the figure, the interior angle and the exterior angle are together equal to two right angles. I. 13.

Therefore all the interior angles, with all the exterior angles, are together equal to twice as many right angles as the figure has sides.

But all the interior angles, with four right angles, are together equal to twice as many right angles as the figure has sides. I. 32, Cor. 1.

Therefore all the interior angles, with all the exterior angles, are together equal to all the interior angles, with four right angles.

Therefore the exterior angles are together equal to four right angles. Q.E.D.


#### EXERCISES ON SIMSON'S COROLLARIES.

[A polygon is said to be **regular** when it has all its sides and all its angles equal.]

1. Express in terms of a right angle the magnitude of each angle of (i) a regular hexagon, (ii) a regular octagon.
2. If one side of a regular hexagon is produced, shew that the exterior angle is equal to the angle of an equilateral triangle.
3. Prove Simson's first Corollary by joining one vertex of the rectilineal figure to each of the other vertices.
4. Find the magnitude of each angle of a regular polygon of  $n$  sides.
5. If the alternate sides of any polygon be produced to meet, the sum of the included angles, together with eight right angles, will be equal to twice as many right angles as the figure has sides.

## PROPOSITION 33. THEOREM.

*The straight lines which join the extremities of two equal and parallel straight lines towards the same parts are themselves equal and parallel.*



Let AB and CD be equal and parallel straight lines ; and let them be joined towards the same parts by the straight lines AC and BD.

*Then shall AC and BD be equal and parallel.*

Construction.

Join BC.

**Proof.** Then because AB and CD are parallel, and BC meets them, therefore the angle ABC is equal to the alternate angle DCB. I. 29.

Now in the triangles ABC, DCB,

Because { AB is equal to DC, *Hyp.*  
and BC is common to both ;  
also the angle ABC is equal to the angle DCB ; *Proved.*  
therefore the triangle ABC is equal to the triangle DCB in all respects ; I. 4.

so that the base AC is equal to the base DB,  
and the angle ACB equal to the angle DBC.

But these are alternate angles.

Therefore AC and BD are parallel : I. 27.


and it has been shewn that they are also equal.

Q.E.D.

**DEFINITION.** A **Parallelogram** is a four-sided figure whose opposite sides are parallel.

PROPOSITION 34. THEOREM.

*The opposite sides and angles of a parallelogram are equal to one another, and each diagonal bisects the parallelogram.*



Let ACDB be a parallelogram, of which BC is a diagonal.

*Then shall the opposite sides and angles of the figure be equal to one another ; and the diagonal BC shall bisect it*

**Proof.** Because AB and CD are parallel, and BC meets them,

therefore the angle ABC is equal to the alternate angle DCB ; I. 29.

Again, because AC and BD are parallel, and BC meets them,

therefore the angle ACB is equal to the alternate angle DBC. I. 29.

Hence in the triangles ABC, DCB,

Because  $\left\{ \begin{array}{l} \text{the angle ABC is equal to the angle DCB,} \\ \text{and the angle ACB is equal to the angle DBC ;} \\ \text{also the side BC is common to both ;} \end{array} \right.$

therefore the triangle ABC is equal to the triangle DCB in all respects ; I. 26.

so that AB is equal to DC, and AC to DB ;  
and the angle BAC is equal to the angle CDB.

Also, because the angle ABC is equal to the angle DCB,  
and the angle CBD equal to the angle BCA,  
therefore the whole angle ABD is equal to the whole angle DCA.

And the triangles ABC, DCB having been proved equal in all respects are equal in area.

Therefore the diagonal BC bisects the parallelogram ACDB. Q.E.D.



## EXERCISES ON PARALLELOGRAMS.

1. *If one angle of a parallelogram is a right angle, all its angles are right angles.*
2. *If the opposite sides of a quadrilateral are equal, the figure is a parallelogram.*
3. *If the opposite angles of a quadrilateral are equal, the figure is a parallelogram.*
4. *If a quadrilateral has all its sides equal and one angle a right angle, all its angles are right angles.*
5. *The diagonals of a parallelogram bisect each other.*
6. *If the diagonals of a quadrilateral bisect each other, the figure is a parallelogram.*
7. *If two opposite angles of a parallelogram are bisected by the diagonal which joins them, the figure is equilateral.*
8. *If the diagonals of a parallelogram are equal, all its angles are right angles.*
9. *In a parallelogram which is not rectangular the diagonals are unequal.*
10. *Any straight line drawn through the middle point of a diagonal of a parallelogram and terminated by a pair of opposite sides, is bisected at that point.*
11. *If two parallelograms have two adjacent sides of one equal to two adjacent sides of the other, each to each, and one angle of one equal to one angle of the other, the parallelograms are equal in all respects.*
12. *Two rectangles are equal if two adjacent sides of one are equal to two adjacent sides of the other, each to each.*
13. *In a parallelogram the perpendiculars drawn from one pair of opposite angles to the diagonal which joins the other pair are equal.*
14. *If ABCD is a parallelogram, and X, Y respectively the middle points of the sides AD, BC; shew that the figure AYCX is a parallelogram.*

MISCELLANEOUS EXERCISES ON SECTIONS I. AND II.

1. Shew that the construction in Proposition 2 may generally be performed in eight different ways. Point out the exceptional case.
2. The bisectors of two vertically opposite angles are in the same straight line.
3. In the figure of Proposition 16, if AF is joined, shew
  - (i) that AF is equal to BC ;
  - (ii) that the triangle ABC is equal to the triangle CFA in all respects.
4. ABC is a triangle right-angled at B, and BC is produced to D : shew that the angle ACD is obtuse.
5. Shew that in any regular polygon of  $n$  sides each angle contains  $\frac{2(n-2)}{n}$  right angles.
6. The angle contained by the bisectors of the angles at the base of any triangle is equal to the vertical angle together with half the sum of the base angles.
7. The angle contained by the bisectors of two exterior angles of any triangle is equal to half the sum of the two corresponding interior angles.
8. If perpendiculars are drawn to two intersecting straight lines from any point between them, shew that the bisector of the angle between the perpendiculars is parallel to (or coincident with) the bisector of the angle between the given straight lines.
9. If two points P, Q be taken in the equal sides of an isosceles triangle ABC, so that BP is equal to CQ, shew that PQ is parallel to BC.
10. ABC and DEF are two triangles, such that AB, BC are equal and parallel to DE, EF, each to each ; shew that AC is equal and parallel to DF.
11. Prove the second Corollary to Prop. 32 by drawing through any angular point lines parallel to all the sides.
12. If two sides of a quadrilateral are parallel, and the remaining two sides equal but not parallel, shew that the opposite angles are supplementary ; also that the diagonals are equal.

## SECTION III.

## THE AREAS OF PARALLELOGRAMS AND TRIANGLES.

Hitherto when two figures have been said to be *equal*, it has been implied that they are *identically equal*, that is, equal in all respects.

But figures may be equal *in area* without being equal in all respects, that is, without having the same shape.

The present section deals with parallelograms and triangles which are equal in area but not necessarily identically equal.

[The ultimate test of equality, as we have already seen, is afforded by Axiom 8, which asserts that magnitudes which *may be made to coincide with one another* are equal. Now figures which are not equal in all respects, cannot be made to coincide without first undergoing some change of form: hence the method of direct *superposition* is unsuited to the purposes of the present section.

We shall see however from Euclid's proof of Proposition 35, that two figures which are not identically equal, may nevertheless be so related to a third figure, that it is possible to infer the equality of their areas.]

## DEFINITIONS.

1. The **Altitude** of a parallelogram with reference to a given side as base, is the perpendicular distance between the base and the opposite side.

2. The **Altitude** of a triangle with reference to a given side as base, is the perpendicular distance of the opposite vertex from the base.


[From this point the following symbols will be introduced into the text:

= for *is equal to*; ∴ for *therefore*.

If it is thought desirable to shorten *written work* by the use of symbols and abbreviations, it is strongly recommended that only some well recognized system should be allowed, such, for example, as that given on page 11.]

PROPOSITION 35. THEOREM.

*Parallelograms on the same base, and between the same parallels, are equal in area.*



Let the parallelograms ABCD, EBCF be on the same base BC, and between the same parallels BC, AF.

*Then shall the parallelogram ABCD be equal in area to the parallelogram EBCF.*

CASE I. If the sides AD, EF, opposite to the base BC, are terminated at the same point D :  
then each of the parallelograms ABCD, EBCF is double of the triangle BDC ; I. 34.

$\therefore$  they are equal to one another. Ax. 6.

CASE II. But if the sides AD, EF are not terminated at the same point :

then because ABCD is a parallelogram,  
 $\therefore$  the side AD = the opposite side BC ; I. 34.

similarly EF = BC ;

$\therefore$  AD = EF. Ax. 1.

$\therefore$  the whole, or remainder, EA = the whole, or remainder, FD.

Then in the triangles FDC, EAB,

FD = EA, *Proved.*

Because  $\left\{ \begin{array}{l} \text{and the side DC = the opposite side AB, I. 34.} \\ \text{also the exterior angle FDC = the interior opposite} \\ \text{angle EAB, I. 29.} \end{array} \right.$

$\therefore$  the triangle FDC = the triangle EAB. I. 4.


From the whole figure ABCF take the triangle FDC ;  
and from the same figure take the equal triangle EAB ;

then the remainders are equal. Ax. 3.

Therefore the parallelogram ABCD is equal to the parallelogram EBCF.

## PROPOSITION 36. THEOREM.

*Parallelograms on equal bases, and between the same parallels, are equal in area.*



Let ABCD, EFGH be parallelograms on equal bases BC, FG, and between the same parallels AH, BG.

*Then shall the parallelogram ABCD be equal to the parallelogram EFGH.*

**Construction.** Join BE, CH.

**Proof.** Then because  $BC = FG$  ; *Hyp.*  
 and the side  $FG =$  the opposite side  $EH$  ; I. 34.  
 $\therefore BC = EH$  ; *Ax. 1.*  
 and  $BC$  is parallel to  $EH$  ; *Hyp.*  
 $\therefore BE$  and  $CH$  are also equal and parallel. I. 33.  
 Therefore  $EBCH$  is a parallelogram. *Def. 36.*

Now the parallelograms ABCD, EBCH are on the same base BC, and between the same parallels BC, AH ;  
 $\therefore$  the parallelogram ABCD = the parallelogram EBCH. I. 35.

Also the parallelograms EFGH, EBCH are on the same base EH, and between the same parallels EH, BG ;  
 $\therefore$  the parallelogram EFGH = the parallelogram EBCH. I. 35.

Therefore the parallelogram ABCD is equal to the parallelogram EFGH. *Ax. 1.*

Q.E.D.


From the last two Propositions we infer that :

(i) *A parallelogram is equal in area to a rectangle of equal base and equal altitude.*

(ii) *Parallelograms on equal bases and of equal altitudes are equal in area.*

PROPOSITION 37. THEOREM.

*Triangles on the same base, and between the same parallels, are equal in area.*



Let the triangles  $ABC$ ,  $DBC$  be upon the same base  $BC$ , and between the same parallels  $BC$ ,  $AD$ .

*Then shall the triangle  $ABC$  be equal to the triangle  $DBC$ .*

**Construction.** Through  $B$  draw  $BE$  parallel to  $CA$ , to meet  $DA$  produced in  $E$ ; I. 31.  
through  $C$  draw  $CF$  parallel to  $BD$ , to meet  $AD$  produced in  $F$ .

**Proof.** Then, by construction, each of the figures  $EBCA$ ,  $DBCF$  is a parallelogram. *Def.* 36.

And since they are on the same base  $BC$ , and between the same parallels  $BC$ ,  $EF$ ;

$\therefore$  the parallelogram  $EBCA$  = the parallelogram  $DBCF$ . I. 35.

Now the diagonal  $AB$  bisects  $EBCA$ ; I. 34.

$\therefore$  the triangle  $ABC$  is half the parallelogram  $EBCA$ .

And the diagonal  $DC$  bisects  $DBCF$ ; I. 34.

$\therefore$  the triangle  $DBC$  is half the parallelogram  $DBCF$ .

And the halves of equal things are equal. *Ax.* 7.


Therefore the triangle  $ABC$  is equal to the triangle  $DBC$ .

Q.E.D.

[For Exercises see page 79.]

## PROPOSITION 38. THEOREM.

*Triangles on equal bases, and between the same parallels, are equal in area.*



Let the triangles  $ABC$ ,  $DEF$  be on equal bases  $BC$ ,  $EF$ , and between the same parallels  $BF$ ,  $AD$ .

*Then shall the triangle  $ABC$  be equal to the triangle  $DEF$ .*

**Construction.** Through  $B$  draw  $BG$  parallel to  $CA$ , to meet  $DA$  produced in  $G$ ; I. 31.  
through  $F$  draw  $FH$  parallel to  $ED$ , to meet  $AD$  produced in  $H$ .

**Proof.** Then, by construction, each of the figures  $GBCA$ ,  $DEFH$  is a parallelogram. Def. 36.

And since they are on equal bases  $BC$ ,  $EF$ , and between the same parallels  $BF$ ,  $GH$ ;

$\therefore$  the parallelogram  $GBCA =$  the parallelogram  $DEFH$ . I. 36.

Now the diagonal  $AB$  bisects  $GBCA$ ; I. 34.

$\therefore$  the triangle  $ABC$  is half the parallelogram  $GBCA$ .

And the diagonal  $DF$  bisects  $DEFH$ ; I. 34.

$\therefore$  the triangle  $DEF$  is half the parallelogram  $DEFH$ .

And the halves of equal things are equal. Ax. 7.

Therefore the triangle  $ABC$  is equal to the triangle  $DEF$ .

Q.E.D.


From this Proposition we infer that :

(i) *Triangles on equal bases and of equal altitude are equal in area.*

(ii) *Of two triangles of the same altitude, that is the greater which has the greater base; and of two triangles on the same base, or on equal bases, that is the greater which has the greater altitude.*

PROPOSITION 39. THEOREM.

*Equal triangles on the same base, and on the same side of it, are between the same parallels.*



Let the triangles  $ABC$ ,  $DBC$  which stand on the same base  $BC$ , and on the same side of it be equal in area.

*Then shall the triangles  $ABC$ ,  $DBC$  be between the same parallels; that is, if  $AD$  be joined,  $AD$  shall be parallel to  $BC$ .*

**Construction.** For if  $AD$  be not parallel to  $BC$ , if possible, through  $A$  draw  $AE$  parallel to  $BC$ , I. 31. meeting  $BD$ , or  $BD$  produced, in  $E$ .  
Join  $EC$ .

**Proof.** Now the triangles  $ABC$ ,  $EBC$  are on the same base  $BC$ , and between the same parallels  $BC$ ,  $AE$ ;

$\therefore$  the triangle  $ABC =$  the triangle  $EBC$ . I. 37.

But the triangle  $ABC =$  the triangle  $DBC$ ; Hyp.

$\therefore$  the triangle  $DBC =$  the triangle  $EBC$ ;

that is, the whole is equal to a part; which is impossible.

$\therefore$   $AE$  is not parallel to  $BC$ .

Similarly it can be shewn that no other straight line through  $A$ , except  $AD$ , is parallel to  $BC$ .

Therefore  $AD$  is parallel to  $BC$ .

Q.E.D.

From this Proposition it follows that :


*Equal triangles on the same base have equal altitudes.*

[For Exercises see page 79.]



## PROPOSITION 40. THEOREM.

*Equal triangles, on equal bases in the same straight line, and on the same side of it, are between the same parallels.*



Let the triangles  $ABC$ ,  $DEF$  which stand on equal bases  $BC$ ,  $EF$ , in the same straight line  $BF$ , and on the same side of it, be equal in area.

*Then shall the triangles  $ABC$ ,  $DEF$  be between the same parallels; that is, if  $AD$  be joined,  $AD$  shall be parallel to  $BF$ .*

**Construction.** For if  $AD$  be not parallel to  $BF$ , if possible, through  $A$  draw  $AG$  parallel to  $BF$ , I. 31. meeting  $ED$ , or  $ED$  produced, in  $G$ .  
Join  $GF$ .

**Proof.** Now the triangles  $ABC$ ,  $GEF$  are on equal bases  $BC$ ,  $EF$ , and between the same parallels  $BF$ ,  $AG$ ;

$\therefore$  the triangle  $ABC =$  the triangle  $GEF$ . I. 38.

But the triangle  $ABC =$  the triangle  $DEF$ : *Hyp.*

$\therefore$  the triangle  $DEF =$  the triangle  $GEF$ :

that is, the whole is equal to a part; which is impossible.

$\therefore$   $AG$  is not parallel to  $BF$ .

Similarly it can be shewn that no other straight line through  $A$ , except  $AD$ , is parallel to  $BF$ .

Therefore  $AD$  is parallel to  $BF$ .

Q.E.D.

From this Proposition it follows that :

- (i) *Equal triangles on equal bases have equal altitudes.*
- (ii) *Equal triangles of equal altitudes have equal bases.*

## EXERCISES ON PROPOSITIONS 37-40.

**DEFINITION.** Each of the three straight lines which join the angular points of a triangle to the middle points of the opposite sides is called a **Median** of the triangle.

## ON PROP. 37.

1. If, in the figure of Prop. 37, AC and BD intersect in K, shew that
  - (i) the triangles AKB, DKC are equal in area.
  - (ii) the quadrilaterals EBKA, FCKD are equal.
2. In the figure of I. 16, shew that the triangles ABC, FBC are equal in area.
3. On the base of a given triangle construct a second triangle, equal in area to the first, and having its vertex in a given straight line.
4. Describe an isosceles triangle equal in area to a given triangle and standing on the same base.

## ON PROP. 38.

5. *A triangle is divided by each of its medians into two parts of equal area.*
6. A parallelogram is divided by its diagonals into four triangles of equal area.
7. ABC is a triangle, and its base BC is bisected at X; if Y be any point in the median AX, shew that the triangles ABY, ACY are equal in area.
8. In AC, a diagonal of the parallelogram ABCD, any point X is taken, and XB, XD are drawn: shew that the triangle BAX is equal to the triangle DAX.
9. If two triangles have two sides of one respectively equal to two sides of the other, and the angles contained by those sides *supplementary*, the triangles are equal in area.

## ON PROP. 39.


10. *The straight line which joins the middle points of two sides of a triangle is parallel to the third side.*
11. *If two straight lines AB, CD intersect in O, so that the triangle AOC is equal to the triangle DOB, shew that AD and CB are parallel.*

## ON PROP. 40.

12. Deduce Prop. 40 from Prop. 39 by joining AE, AF in the figure of page 78.

## PROPOSITION 41. THEOREM.

*If a parallelogram and a triangle be on the same base and between the same parallels, the parallelogram shall be double of the triangle.*



Let the parallelogram  $ABCD$ , and the triangle  $EBC$  be upon the same base  $BC$ , and between the same parallels  $BC$ ,  $AE$ .

*Then shall the parallelogram  $ABCD$  be double of the triangle  $EBC$ .*

**Construction.** Join  $AC$ .

**Proof.** Now the triangles  $ABC$ ,  $EBC$  are on the same base  $BC$ , and between the same parallels  $BC$ ,  $AE$ ;

$\therefore$  the triangle  $ABC =$  the triangle  $EBC$ . I. 37.

And since the diagonal  $AC$  bisects  $ABCD$ ; I. 34.

$\therefore$  the parallelogram  $ABCD$  is double of the triangle  $ABC$ .

Therefore the parallelogram  $ABCD$  is also double of the triangle  $EBC$ . Q.E.D.

## EXERCISES. •

1.  $ABCD$  is a parallelogram, and  $X$ ,  $Y$  are the middle points of the sides  $AD$ ,  $BC$ ; if  $Z$  is any point in  $XY$ , or  $XY$  produced, shew that the triangle  $AZB$  is one quarter of the parallelogram  $ABCD$ .


2. Describe a right-angled isosceles triangle equal to a given square.

3. If  $ABCD$  is a parallelogram, and  $X$ ,  $Y$  any points in  $DC$  and  $AD$  respectively: shew that the triangles  $AXB$ ,  $BYC$  are equal in area.

4.  $ABCD$  is a parallelogram, and  $P$  is any point within it; shew that the sum of the triangles  $PAB$ ,  $PCD$  is equal to half the parallelogram.

PROPOSITION 42. PROBLEM.

To describe a parallelogram that shall be equal to a given triangle, and have one of its angles equal to a given angle.



Let ABC be the given triangle, and D the given angle.

It is required to describe a parallelogram equal to ABC, and having one of its angles equal to D.

Construction. Bisect BC at E. I. 10.

At E in CE, make the angle CEF equal to D; I. 23.

through A draw AFG parallel to EC; I. 31.

and through C draw CG parallel to EF.

Then FECG shall be the parallelogram required.

Join AE.

Proof. Now the triangles ABE, AEC are on equal bases BE, EC, and between the same parallels;

$\therefore$  the triangle ABE = the triangle AEC; I. 38.

$\therefore$  the triangle ABC is double of the triangle AEC.

But FECG is a parallelogram by construction; Def. 36

and it is double of the triangle AEC,

being on the same base EC, and between the same parallels EC and AG. I. 41.

Therefore the parallelogram FECG is equal to the triangle ABC;

and it has one of its angles CEF equal to the given angle D.


Q.E.F.

EXERCISES.

1. Describe a parallelogram equal to a given square standing on the same base, and having an angle equal to half a right angle.

2. Describe a rhombus equal to a given parallelogram and standing on the same base. When does the construction fail?

**DEFINITION.** If in the diagonal of a parallelogram any point is taken, and straight lines are drawn through it parallel to the sides of the parallelogram; then of the four parallelograms into which the whole figure is divided, the two through which the diagonal passes are called **Parallelograms about that diagonal**, and the other two, which with these make up the whole figure, are called the **complements** of the parallelograms about the diagonal.




Thus in the figure given above,  $AEKH$ ,  $KCGF$  are parallelograms about the diagonal  $AC$ ; and the shaded figures  $HKCF$ ,  $EBKG$  are the complements of those parallelograms.

**NOTE.** A parallelogram is often named by *two* letters only, these being placed at opposite angular points.

PROPOSITION 43. THEOREM.

*The complements of the parallelograms about the diagonal of any parallelogram, are equal to one another.*



Let ABCD be a parallelogram, and KD, KB the complements of the parallelograms EH, GF about the diagonal AC. Then shall the complement BK be equal to the complement KD.

**Proof.** Because EH is a parallelogram, and AK its diagonal,  $\therefore$  the triangle AEK = the triangle AHK. I. 34.

Similarly the triangle KGC = the triangle KFC.

Hence the triangles AEK, KGC are together equal to the triangles AHK, KFC.

But since the diagonal AC bisects the parallelogram ABCD;  $\therefore$  the whole triangle ABC = the whole triangle ADC. I. 34.

Therefore the remainder, the complement BK, is equal to the remainder, the complement KD. Q.E.D.


EXERCISES.

In the figure of Prop. 43, prove that

- (i) The parallelogram ED is equal to the parallelogram BH.
- (ii) If KB, KD are joined, the triangle AKB is equal to the triangle AKD,

## PROPOSITION 44. PROBLEM.

To a given straight line to apply a parallelogram which shall be equal to a given triangle, and have one of its angles equal to a given angle.



Let AB be the given straight line, C the given triangle, and D the given angle.

It is required to apply to the straight line AB a parallelogram equal to the triangle C, and having an angle equal to the angle D.

**Construction.** On AB produced describe a parallelogram BEFG equal to the triangle C, and having the angle EBG equal to the angle D. I. 22 and I. 42\*.

Through A draw AH parallel to BG or EF, to meet FG produced in H. I. 31.

Join HB.

Then because AH and EF are parallel, and HF meets them,  $\therefore$  the angles AHF, HFE together = two right angles. I. 29.

Hence the angles BHF, HFE are together less than two right angles;

$\therefore$  HB and FE will meet if produced towards B and E. Ax. 12.

Produce HB and FE to meet at K.

Through K draw KL parallel to EA or FH; I. 31. and produce HA, GB to meet KL in the points L and M.

Then shall BL be the parallelogram required.

**Proof.** Now FHLK is a parallelogram, *Constr.*  
and LB, BF are the complements of the parallelograms  
about the diagonal HK :

$\therefore$  the complement LB = the complement BF. I. 43.

But the triangle C = the figure BF ; *Constr.*  
 $\therefore$  the figure LB = the triangle C.

Again the angle ABM = the vertically opposite angle GBE ;  
also the angle D = the angle GBE ; *Constr.*  
 $\therefore$  the angle ABM = the angle D.

Therefore the parallelogram LB, which is applied to the  
straight line AB, is equal to the triangle C, and has the  
angle ABM equal to the angle D. Q.E.F.

\*This step of the construction is effected by first describing on  
AB produced a triangle whose sides are respectively equal to those of  
the triangle C (I. 22) ; and by then making a parallelogram equal to  
the triangle so drawn, and having an angle equal to D (I. 42).


#### QUESTIONS FOR REVISION.

1. Quote Euclid's Twelfth Axiom. What objections have been raised to it, and what substitute for it has been suggested ?
2. Which of Euclid's Propositions, dealing with parallel straight lines, depends on Axiom 12 ? Furnish an alternative proof.
3. *Straight lines which are parallel to the same straight line are parallel to one another* [Prop. 30]. Deduce this from Playfair's Axiom.
4. Define a *parallelogram*, an *altitude* of a triangle, a *median* of a triangle, *parallelograms about the diagonal* of a parallelogram.
5. What is meant by *superposition* ? On what Axiom does this method depend ? Give instances of figures which are equal *in area*, but which cannot be superposed.
6. In fig. 2 of Prop. 35 shew how one parallelogram may be cut into pieces, which, when fitted together in other positions, make up the other parallelogram.



## PROPOSITION 45. PROBLEM.

To describe a parallelogram equal to a given rectilineal figure, and having an angle equal to a given angle.



Let ABCD be the given rectilineal figure, and E the given angle.

It is required to describe a parallelogram equal to ABCD, and having an angle equal to E.

Suppose the given rectilineal figure to be a quadrilateral.

Construction. Join BD.

Describe the parallelogram FKH equal to the triangle ABD, and having the angle FKH equal to the angle E. I. 42.

To GH apply the parallelogram GM, equal to the triangle DBC, and having the angle GHM equal to E. I. 44.

Then shall FKML be the parallelogram required.

Proof. Because each of the angles GHM, FKH = the angle E;  
 $\therefore$  the angle FKH = the angle GHM.

To each of these equals add the angle GHK;  
 then the angles FKH, GHK together = the angles GHM, GHK.

But since FK, GH are parallel, and KH meets them;  
 $\therefore$  the angles FKH, GHK together = two right angles; I. 29.

$\therefore$  also the angles GHM, GHK together = two right angles;  
 $\therefore$  KH, HM are in the same straight line. I. 14.

Again, because KM, FG are parallel, and HG meets them,  
 $\therefore$  the angle MHG = the alternate angle HGF. I. 29.

To each of these equals add the angle HGL;  
 then the angles MHG, HGL together = the angles HGF, HGL.

But because HM, GL are parallel, and HG meets them,  
 $\therefore$  the angles MHG, HGL together = two right angles : I. 29.

$\therefore$  also the angles HGF, HGL together = two right angles :  
 $\therefore$  FG, GL are in the same straight line. I. 14.

And because KF and ML are each parallel to HG, *Constr.*  
 therefore KF is parallel to ML; I. 30.  
 and KM, FL are parallel; *Constr.*  
 $\therefore$  FKML is a parallelogram. *Def.* 36.


Again, because the parallelogram FH = the triangle ABD,  
 and the parallelogram GM = the triangle DBC; *Constr.*  
 $\therefore$  the whole parallelogram FKML = the whole figure ABCD;  
 and it has the angle FKM equal to the angle E.

By a series of similar steps, a parallelogram may be  
 constructed equal to a rectilineal figure of more than four  
 sides. Q.E.F.

The following Problem is important, and furnishes a useful application of the principles of the foregoing propositions.

ADDITIONAL PROBLEM.

*To describe a triangle equal in area to a given quadrilateral.*



Let ABCD be the given quadrilateral.

*It is required to describe a triangle equal to ABCD in area.*

**Construction.** Join BD.  
Through C draw CX parallel to BD, meeting AD produced in X.  
Join BX.

Then XAB shall be the required triangle.

**Proof.** Now the triangles XDB, CDB are on the same base DB and between the same parallels DB, XC;

$\therefore$  the triangle XDB = the triangle CDB in area. I. 37.

To each of these equals add the triangle ADB;  
then the triangle XAB = the figure ABCD.


EXERCISE.

Construct a rectilinear figure equal to a given rectilinear figure, and having fewer sides by one than the given figure.

Hence shew how to construct a triangle equal to a given rectilinear figure.

PROPOSITION 46. PROBLEM.

To describe a square on a given straight line.



Let AB be the given straight line.

It is required to describe a square on AB.

**Constr.** From A draw AC at right angles to AB ; I. 11.  
and make AD equal to AB. I. 3.

Through D draw DE parallel to AB ; I. 31.  
and through B draw BE parallel to AD, meeting DE in E.  
Then shall ADEB be a square.

**Proof.** For, by construction, ADEB is a parallelogram :  
 $\therefore AB = DE$ , and  $AD = BE$ . I. 34.

But  $AD = AB$  ; *Constr.*  
 $\therefore$  the four straight lines AB, AD, DE, EB are all equal ;  
that is, the figure ADEB is equilateral.

Again, since AB, DE are parallel, and AD meets them,  
 $\therefore$  the angles BAD, ADE together = two right angles ; I. 29.  
but the angle BAD is a right angle ; *Constr.*  
 $\therefore$  also the angle ADE is a right angle.

And the opposite angles of a parallelogram are equal ; I. 34.  
 $\therefore$  each of the angles DEB, EBA is a right angle :  
that is the figure ADEB is rectangular.


Hence it is a square, and it is described on AB.

Q. E. F.

**COROLLARY.** If one angle of a parallelogram is a right angle, all its angles are right angles.

## PROPOSITION 47. THEOREM.

*In a right-angled triangle the square described on the hypotenuse is equal to the sum of the squares described on the other two sides.*



Let  $ABC$  be a right-angled triangle, having the angle  $BAC$  a right angle.

*Then shall the square described on the hypotenuse  $BC$  be equal to the sum of the squares described on  $BA$ ,  $AC$ .*

**Construction.** On  $BC$  describe the square  $BDEC$ ; I. 46.  
and on  $BA$ ,  $AC$  describe the squares  $BAGF$ ,  $ACKH$ .

Through  $A$  draw  $AL$  parallel to  $BD$  or  $CE$ ; I. 31.  
and join  $AD$ ,  $FC$ .

**Proof.** Then because each of the angles  $BAC$ ,  $BAG$  is a right angle,

$\therefore CA$  and  $AG$  are in the same straight line. I. 14.

Now the angle  $CBD =$  the angle  $FBA$ ,  
for each of them is a right angle.

Add to each the angle  $ABC$  :  
then the whole angle  $ABD =$  the whole angle  $FBC$ .

Then in the triangles ABD, FBC,  
 Because  $\left\{ \begin{array}{l} AB = FB, \\ \text{and } BD = BC, \\ \text{also the angle } ABD = \text{the angle } FBC; \\ \therefore \text{ the triangle } ABD = \text{the triangle } FBC. \end{array} \right. \quad \begin{array}{l} \textit{Proved.} \\ \text{I. 4.} \end{array}$

Now the parallelogram BL is double of the triangle ABD, being on the same base BD, and between the same parallels BD, AL. I. 41.

And the square GB is double of the triangle FBC, being on the same base FB, and between the same parallels FB, GC. I. 41.

But doubles of equals are equal : Ax. 6.  
 therefore the parallelogram BL = the square GB.

Similarly, by joining AE, BK it can be shewn that the parallelogram CL = the square CH.

Therefore the whole square BE = the sum of the squares GB, HC :

that is, the square described on the hypotenuse BC is equal to the sum of the squares described on the two sides BA, AC. Q.E.D.

**NOTE.** It is not necessary to the proof of this Proposition that the three squares should be described *external* to the triangle ABC ; and since *each* square may be drawn either *towards* or *away from* the triangle, it may be shewn that there are  $2 \times 2 \times 2$ , or *eight*, possible constructions.

*Obs.* The following properties of a square, though not formally enunciated by Euclid, are employed in subsequent proofs. [See I. 48.]

- (i) *The squares on equal straight lines are equal.*
- (ii) *Equal squares stand upon equal straight lines.*

## EXERCISES ON PROPOSITION 47.

1. In the figure of this Proposition, shew that
  - (i) If  $BG, CH$  are joined, these straight lines are parallel ;
  - (ii) The points  $F, A, K$  are in one straight line ;
  - (iii)  $FC$  and  $AD$  are at right angles to one another ;
  - (iv) If  $GH, KE, FD$  are joined, the triangle  $GAH$  is equal to the given triangle in all respects ; and the triangles  $FBD, KCE$  are each equal in area to the triangle  $ABC$ .  
[See Ex. 9, p. 79.]
2. On the sides  $AB, AC$  of any triangle  $ABC$ , squares  $ABFG, ACKH$  are described both toward the triangle, or both on the side remote from it : shew that the straight lines  $BH$  and  $CG$  are equal.
3. On the sides of any triangle  $ABC$ , equilateral triangles  $BCX, CAY, ABZ$  are described, all externally, or all towards the triangle : shew that  $AX, BY, CZ$  are all equal.
4. The square described on the diagonal of a given square, is double of the given square.
5.  $ABC$  is an equilateral triangle, and  $AX$  is the perpendicular drawn from  $A$  to  $BC$  : shew that the square on  $AX$  is three times the square on  $BX$ .
6. Describe a square equal to the sum of two given squares.
7. From the vertex  $A$  of a triangle  $ABC$ ,  $AX$  is drawn perpendicular to the base : shew that the difference of the squares on the sides  $AB$  and  $AC$ , is equal to the difference of the squares on  $BX$  and  $CX$ , the segments of the base.
8. If from any point  $O$  within a triangle  $ABC$ , perpendiculars  $OX, OY, OZ$  are drawn to the sides  $BC, CA, AB$  respectively : shew that the sum of the squares on the segments  $AZ, BX, CY$  is equal to the sum of the squares on the segments  $AY, CX, BZ$ .
9.  $ABC$  is a triangle right-angled at  $A$  ; and the sides  $AB, AC$  are intersected by a straight line  $PQ$ , and  $BQ, PC$  are joined. Prove that the sum of the squares on  $BQ, PC$  is equal to the sum of the squares on  $BC, PQ$ .
10. In a right-angled triangle four times the sum of the squares on the two medians drawn from the acute angles is equal to five times the square on the hypotenuse.

NOTES ON PROPOSITION 47.

It is believed that Proposition 47 is due to Pythagoras, a Greek philosopher and mathematician, who lived about two centuries before Euclid.


Many experimental proofs of this theorem have been given by means of actual *dissection*: that is to say, it has been shewn how the squares on the sides containing the right angle may be cut up into pieces which, when fitted together in other positions, exactly make up the square on the hypotenuse. Two of these methods of dissection are given below.

I. In the adjoining diagram ABC is the given right-angled triangle, and the figures AF, HK are the squares on AB, AC, placed side by side.

FD is made equal to EH or AC; and the two squares AF, HK are cut along the lines ED, DB.


Then it will be found that the triangle EHD may be placed so as to fill up the space CAB; and the triangle BFD may be made to fill the space CKE.

Hence the two squares AF, HK may be fitted together so as to form the single figure CBDE, which will be found to be a perfect square, namely the square on the hypotenuse BC.



II. In the figure of 1. 47, let DB and EC be produced to meet FG and AH in L and N respectively; and let LM be drawn parallel to BC.


Then it will be found that the several parts of the two squares FA, AK can be fitted together (in the places bearing corresponding numbers) so as exactly to fill up the square DC.





## PROPOSITION 48. THEOREM.

If the square described on one side of a triangle be equal to the sum of the squares described on the other two sides, then the angle contained by these two sides shall be a right angle.



Let ABC be a triangle ; and let the square described on BC be equal to the sum of the squares described on BA, AC.

Then shall the angle BAC be a right angle.

**Construction.** From A draw AD at right angles to AC; I. 11.  
and make AD equal to AB. I. 3.  
Join DC.

**Proof.** Then, because AD = AB, *Constr.*  
∴ the square on AD = the square on AB.  
To each of these add the square on CA ;  
then the sum of the squares on CA, AD = the sum of the squares on CA, AB.

But, because the angle DAC is a right angle, *Constr.*  
∴ the square on DC = the sum of the squares on CA, AD. I. 47.  
And, by hypothesis, the square on BC = the sum of the squares on CA, AB ;

∴ the square on DC = the square on BC :  
∴ also the side DC = the side BC.

Then in the triangles DAC, BAC,  
Because { DA = BA, *Constr.*  
and AC is common to both ;  
also the third side DC = the third side BC; *Proved.*  
∴ the angle DAC = the angle BAC. I. 8.  
But DAC is a right angle. *Constr.*  
Therefore also BAC is a right angle. Q.E.D.

# THEOREMS AND EXAMPLES ON BOOK I.

## INTRODUCTORY.

### HINTS TOWARDS THE SOLUTION OF GEOMETRICAL EXERCISES.

#### ANALYSIS. SYNTHESIS.

It is commonly found that exercises in Pure Geometry present to a beginner far more difficulty than examples in any other branch of Elementary Mathematics. This seems to be due to the following causes :

(i) The variety of such exercises is practically unlimited ; and it is impossible to lay down for their treatment any definite methods, such for example as the rules of Elementary Arithmetic and Algebra.

(ii) The arrangement of Euclid's Propositions, though perhaps the most *convincing* of all forms of argument, affords in most cases little clue as to the way in which the proof or construction *was discovered*.

Euclid's propositions are arranged **synthetically** : that is to say, starting from the hypothesis or data, they first give a construction in accordance with postulates, and problems already solved ; then by successive steps based on known theorems, they prove what was required in the enunciation.

Thus Geometrical Synthesis is a *building up* of *known* results, in order to obtain a *new* result.

But as this is not the way in which constructions or proofs are usually discovered, we draw the student's attention to the following hints.


Begin by *assuming* the result it is desired to establish ; then by working backwards, trace the consequences of the assumption, and try to ascertain its dependence on some simpler theorem which is already known to be true, or on some condition which suggests the necessary construction. If this attempt is successful, the steps of the argument may in general be re-arranged in reverse order, and the construction and proof presented in a synthetic form.

This unravelling of a proposition in order to trace it back to some earlier principle on which it depends, is called **geometrical analysis** : it is the natural way of attacking many theorems, and it is especially useful in solving *problems*.

Although the above directions do not amount to a *method*, they often furnish a mode of *searching for a suggestion*. Geometrical Analysis however can only be used with success when a thorough grasp of the chief propositions of Euclid has been gained.

The practical application of the foregoing hints is illustrated by the following examples.

1. *Construct an isosceles triangle having given the base, and the sum of one of the equal sides and the perpendicular drawn from the vertex to the base.*



Let  $AB$  be the given base, and  $K$  the sum of one side and the perpendicular drawn from the vertex to the base.

**ANALYSIS.** *Suppose  $ABC$  to be the required triangle.*

From  $C$  draw  $CX$  perpendicular to  $AB$  :  
then  $AB$  is bisected at  $X$ .

I. 26.

Now if we produce  $XC$  to  $H$ , making  $XH$  equal to  $K$ ,  
it follows that  $CH = CA$  ;  
and if  $AH$  is joined,

we notice that the angle  $CAH =$  the angle  $CHA$ . I. 5.

Now the straight lines  $XH$  and  $AH$  can be drawn *before the position of  $C$  is known* ;

Hence we have the following construction, which we arrange synthetically.

**SYNTHESIS.** Bisect  $AB$  at  $X$  :  
from  $X$  draw  $XH$  perpendicular to  $AB$ , making  $XH$  equal to  $K$ .  
Join  $AH$ .

At the point  $A$  in  $HA$ , make the angle  $HAC$  equal to the angle  $AHX$ .  
Join  $CB$ .

*Then  $ACB$  shall be the triangle required.*

First the triangle is isosceles, for  $AC = BC$ . I. 4.

Again, since the angle  $HAC =$  the angle  $AHC$ , *Constr.*

$\therefore HC = AC$ . I. 6.


To each add  $CX$  ;

then the sum of  $AC, CX =$  the sum of  $HC, CX =$   $HX$ .

That is, the sum of  $AC, CX = K$ .

Q. E. F.

2. To divide a given straight line so that the square on one part may be double of the square on the other.



Let AB be the given straight line.

**ANALYSIS.** Suppose AB to be divided as required at X: that is, suppose the square on AX to be double of the square on XB.

Now we remember that in an isosceles right-angled triangle, the square on the hypotenuse is double of the square on either of the equal sides.

This suggests to us to draw BC perpendicular to AB, to make BC equal to BX, and to join XC.

Then the square on XC is double of the square on XB; I. 47.  
 $\therefore XC = AX$ .

Hence when we join AC, we notice that  
 the angle XAC = the angle XCA. I. 5.

Thus the exterior angle CXB is double of the angle XAC. I. 32.

But the angle CXB is half of a right angle: I. 32.  
 $\therefore$  the angle XAC is one-fourth of a right angle.

This supplies the clue to the following construction:—

**SYNTHESIS.** From B draw BD perpendicular to AB;  
 and from A draw AC, making BAC one-fourth of a right angle.

From C, the intersection of AC and BD, draw CX, making the angle ACX equal to the angle BAC. I. 23.

Then AB shall be divided as required at X.

For since the angle XCA = the angle XAC,  
 $\therefore XA = XC$ . I. 6.

And because the angle BXC = the sum of the angles BAC, ACX, I. 32.  
 $\therefore$  the angle BXC is half a right angle.

And the angle at B is a right angle;  
 $\therefore$  the angle BCX is half a right angle; I. 32.

$\therefore$  the angle BXC = the angle BCX;

$\therefore BX = BC$ .

Hence the square on XC is double of the square on XB: I. 47.  
 that is, the square on AX is double of the square on XB. Q.E.F.

## I. ON THE IDENTICAL EQUALITY OF TRIANGLES.

See Propositions 4, 8, 26.

1. If in a triangle the perpendicular from the vertex on the base bisects the base, then the triangle is isosceles.

2. If the bisector of the vertical angle of a triangle is also perpendicular to the base, the triangle is isosceles.

3. If the bisector of the vertical angle of a triangle also bisects the base, the triangle is isosceles.

[Produce the bisector, and complete the construction after the manner of I. 16.]

4. If in a triangle a pair of straight lines drawn from the extremities of the base, making equal angles with the remaining sides, are equal, the triangle is isosceles.

5. If in a triangle the perpendiculars drawn from the extremities of the base to the opposite sides are equal, the triangle is isosceles.

6. Two triangles  $ABC$ ,  $ABD$  on the same base  $AB$ , and on opposite sides of it, are such that  $AC$  is equal to  $AD$ , and  $BC$  is equal to  $BD$ : shew that the line joining the points  $C$  and  $D$  is perpendicular to  $AB$ .

7. If from the extremities of the base of an isosceles triangle perpendiculars are drawn to the opposite sides, shew that the straight line joining the vertex to the intersection of these perpendiculars bisects the vertical angle.


8.  $ABC$  is a triangle in which the vertical angle  $BAC$  is bisected by the straight line  $AX$ : from  $B$  draw  $BD$  perpendicular to  $AX$ , and produce it to meet  $AC$ , or  $AC$  produced, in  $E$ ; then shew that  $BD$  is equal to  $DE$ .

9. In a quadrilateral  $ABCD$ ,  $AB$  is equal to  $AD$ , and  $BC$  is equal to  $DC$ : shew that the diagonal  $AC$  bisects each of the angles which it joins.

10. In a quadrilateral  $ABCD$  the opposite sides  $AD$ ,  $BC$  are equal, and also the diagonals  $AC$ ,  $BD$  are equal: if  $AC$  and  $BD$  intersect at  $K$ , shew that each of the triangles  $AKB$ ,  $DKC$  is isosceles.

11. If one angle of a triangle be equal to the sum of the other two, the greatest side is double of the distance of its middle point from the opposite angle.

12. *Two right-angled triangles which have their hypotenuses equal, and one side of one equal to one side of the other, are equal in all respects.*



Let  $ABC$ ,  $DEF$  be two  $\triangle^s$  right-angled at  $B$  and  $E$ , having  $AC$  equal to  $DF$ , and  $AB$  equal to  $DE$ .

*Then shall the  $\triangle ABC$  be equal to the  $\triangle DEF$  in all respects.*

For apply the  $\triangle ABC$  to the  $\triangle DEF$ , so that  $AB$  may coincide with the equal line  $DE$ , and  $C$  may fall on the side of  $DE$  remote from  $F$ . Let  $C'$  be the point on which  $C$  falls.

Then  $DEC'$  represents the  $\triangle ABC$  in its new position.

Now each of the  $\angle^s$   $DEF$ ,  $DEC'$  is a rt.  $\angle$ ; *Hyp.*  
 $\therefore EF$  and  $EC'$  are in one st. line. I. 14.

Then in the  $\triangle C'DF$ , because  $DF = DC'$  (*i.e.*  $AC$ ), *Hyp.*  
 $\therefore$  the  $\angle DFC' =$  the  $\angle DC'F$ . I. 5.

Hence in the two  $\triangle^s$   $DEF$ ,  $DEC'$ ,  
 Because  $\left\{ \begin{array}{l} \text{the } \angle DEF = \text{the } \angle DEC', \text{ being rt. } \angle^s; \\ \text{and the } \angle DFE = \text{the } \angle DC'E; \\ \text{also the side } DE \text{ is common to both;} \end{array} \right.$  *Proved.*  
 $\therefore$  the  $\triangle^s$   $DEF$ ,  $DEC'$  are equal in all respects; I. 26.  
 that is, the  $\triangle^s$   $DEF$ ,  $ABC$  are equal in all respects. Q.E.D.

*Alternative Proof.* Since the  $\angle ABC$  is a rt. angle;  
 $\therefore$  the sq. on  $AC =$  the sqq. on  $AB$ ,  $BC$ . I. 47.


Similarly, the sq. on  $DF =$  the sqq. on  $DE$ ,  $EF$ ; I. 47.

But the sq. on  $AC =$  the sq. on  $DF$ , since  $AC = DF$ ;  
 $\therefore$  the sqq. on  $AB$ ,  $BC =$  the sqq. on  $DE$ ,  $EF$ .

And of these, the sq. on  $AB =$  the sq. on  $DE$ , since  $AB = DE$ ;  
 $\therefore$  the sq. on  $BC =$  the sq. on  $EF$ ; *Ax.* 3.  
 $\therefore BC = EF$ .

Hence the three sides of the  $\triangle ABC$  are respectively equal to the three sides of the  $\triangle DEF$ ;  
 $\therefore$  the  $\triangle ABC =$  the  $\triangle DEF$  in all respects. I. 8.

13. *If two triangles have two sides of the one equal to two sides of the other, each to each, and have likewise the angles opposite to one pair of equal sides equal, then the angles opposite to the other pair of equal sides shall be either equal or supplementary, and in the former case the triangles shall be equal in all respects.*



Let  $ABC, DEF$  be two triangles, in which  
the side  $AB =$  the side  $DE$ ,  
the side  $AC =$  the side  $DF$ ,  
and the  $\angle ABC =$  the  $\angle DEF$ .

*Then shall the  $\angle^s ACB, DFE$  be either equal (as in Figs. 1 and 2) or supplementary (as in Figs. 1 and 3); and in the former case the triangles shall be equal in all respects.*

If the  $\angle BAC =$  the  $\angle EDF$ . [Figs. 1 and 2.]  
then the  $\angle ACB =$  the  $\angle DFE$ , and the triangles are equal in all respects. I. 4.

But if the  $\angle BAC$  be not equal to the  $\angle EDF$ , [Figs. 1 and 3.]  
let the  $\angle EDF$  be greater than the  $\angle BAC$ .

At  $D$  in  $ED$  make the  $\angle EDF'$  equal to the  $\angle BAC$ .  
Then the  $\triangle^s BAC, EDF'$  are equal in all respects. I. 26.

$\therefore AC = DF'$ ;  
but  $AC = DF$ ;  
 $\therefore DF = DF'$ , Hyp.

$\therefore$  the angle  $DFF' =$  the  $\angle DF'F$ . I. 5.

But the  $\angle^s DF'F, DF'E$  are supplementary, I. 13.  
 $\therefore$  the  $\angle^s DFF', DF'E$  are supplementary:  
that is, the  $\angle^s DFE, ACB$  are supplementary.

Q.E.D.

**COROLLARIES.** Three cases of this theorem deserve special attention.

It has been proved that if the angles  $ACB, DFE$  are not supplementary they are equal:

Hence, in addition to the hypothesis of this theorem,

- (i) If the angles  $ACB$ ,  $DFE$  opposite to the two equal sides  $AB$ ,  $DE$  are both acute or both obtuse they cannot be supplementary, and are therefore equal; or if one of them is a right angle, the other must also be a right angle (whether considered as supplementary or equal to it):

in either case the triangles are equal in all respects.

- (ii) If the two given angles are right angles or obtuse angles, it follows that the angles  $ACB$ ,  $DFE$  must be both acute, and therefore equal, by (i):

so that the triangles are equal in all respects.

- (iii) If in each triangle the side opposite the given angle is not less than the other given side; that is, if  $AC$  and  $DF$  are not less than  $AB$  and  $DE$  respectively) then the angles  $ACB$ ,  $DFE$  cannot be greater than the angles  $ABC$ ,  $DEF$  respectively;

therefore the angles  $ACB$ ,  $DFE$  are both acute;

hence, as above, they are equal;

and the triangles  $ABC$ ,  $DEF$  are equal in all respects.

## II. ON INEQUALITIES.

See Propositions 16, 17, 18, 19, 20, 21, 24, 25.

1. In a triangle  $ABC$ , if  $AC$  is not greater than  $AB$ , shew that any straight line drawn through the vertex  $A$ , and terminated by the base  $BC$ , is less than  $AB$ .

2.  $ABC$  is a triangle, and the vertical angle  $BAC$  is bisected by a straight line which meets the base  $BC$  in  $X$ ; shew that  $BA$  is greater than  $BX$ , and  $CA$  greater than  $CX$ . Hence obtain a proof of 1. 20.

3. The perpendicular is the shortest straight line that can be drawn from a given point to a given straight line; and of others, that which is nearer to the perpendicular is less than the more remote; and two, and only two equal straight lines can be drawn from the given point to the given straight line, one on each side of the perpendicular.

4. The sum of the distances of any point from the three angular points of a triangle is greater than half its perimeter.

5. The sum of the distances of any point within a triangle from its angular points is less than the perimeter of the triangle.



6. The perimeter of a quadrilateral is greater than the sum of its diagonals.

7. The sum of the diagonals of a quadrilateral is less than the sum of the four straight lines drawn from the angular points to any given point. Prove this, and point out the exceptional case.

8. *In a triangle any two sides are together greater than twice the median which bisects the remaining side.* [See Def. p. 79.]

[Produce the median, and complete the construction after the manner of I. 16.]

9. *In any triangle the sum of the medians is less than the perimeter.*

10. In a triangle an angle is acute, obtuse, or a right angle, according as the median drawn from it is greater than, less than, or equal to half the opposite side. [See Ex. 4, p. 65.]

11. The diagonals of a rhombus are unequal.

12. *If the vertical angle of a triangle is contained by unequal sides, and if from the vertex the median and the bisector of the angle are drawn, then the median lies within the angle contained by the bisector and the longer side.*

Let  $ABC$  be a  $\triangle$ , in which  $AB$  is greater than  $AC$ ; let  $AX$  be the median drawn from  $A$ , and  $AP$  the bisector of the vertical  $\angle BAC$ .

Then shall  $AX$  lie between  $AP$  and  $AB$ .

Produce  $AX$  to  $K$ , making  $XK$  equal to  $AX$ . Join  $KC$ .

Then the  $\triangle$ 's  $BXA$ ,  $CXK$  may be shewn to be equal in all respects; I. 4.  
hence  $BA = CK$ , and the  $\angle BAX =$  the  $\angle CKX$ .

But since  $BA$  is greater than  $AC$ , *Hyp.*


$\therefore CK$  is greater than  $AC$ ;

$\therefore$  the  $\angle CAK$  is greater than the  $\angle CKA$ : I. 18.

that is, the  $\angle CAX$  is greater than the  $\angle BAX$ ;

$\therefore$  the  $\angle CAX$  must be more than half the vert.  $\angle BAC$ ;

hence  $AX$  lies within the angle  $BAP$ . Q.E.D.



13. *If the vertical angle of a triangle is contained by two unequal sides, and if from the vertex there are drawn the bisector of the vertical angle, the median, and the perpendicular to the base, the first of these lines is intermediate in position and magnitude to the other two.*

## III. ON PARALLELS.

See Propositions 27—31.

1. If a straight line meets two parallel straight lines, and the two interior angles on the same side are bisected; shew that the bisectors meet at right angles. [I. 29, I. 32.]

2. The straight lines drawn from any point in the bisector of an angle parallel to the arms of the angle, and terminated by them, are equal; and the resulting figure is a rhombus.

3.  $AB$  and  $CD$  are two straight lines intersecting at  $D$ , and the adjacent angles so formed are bisected: if through any point  $X$  in  $DC$  a straight line  $YXZ$  be drawn parallel to  $AB$  and meeting the bisectors in  $Y$  and  $Z$ , shew that  $XY$  is equal to  $XZ$ .

4. If two straight lines are parallel to two other straight lines, each to each; and if the acute angles contained by each pair are bisected; shew that the bisecting lines are parallel.

5. The middle point of any straight line which meets two parallel straight lines, and is terminated by them, is equidistant from the parallels.

6. A straight line drawn between two parallels and terminated by them, is bisected; shew that any other straight line passing through the middle point and terminated by the parallels, is also bisected at that point.

7. If through a point equidistant from two parallel straight lines, two straight lines are drawn cutting the parallels, the portions of the latter thus intercepted are equal.

## PROBLEMS.

8.  $AB$  and  $CD$  are two given straight lines, and  $X$  is a given point in  $AB$ : find a point  $Y$  in  $AB$  such that  $YX$  may be equal to the perpendicular distance of  $Y$  from  $CD$ .

9.  $ABC$  is an isosceles triangle: required to draw a straight line  $DE$  parallel to the base  $BC$ , and meeting the equal sides in  $D$  and  $E$ , so that  $BD$ ,  $DE$ ,  $EC$  may be all equal.

10.  $ABC$  is any triangle: required to draw a straight line  $DE$  parallel to the base  $BC$ , and meeting the other sides in  $D$  and  $E$ , so that  $DE$  may be equal to the sum of  $BD$  and  $CE$ .

11.  $ABC$  is any triangle: required to draw a straight line parallel to the base  $BC$ , and meeting the other sides in  $D$  and  $E$ , so that  $DE$  may be equal to the difference of  $BD$  and  $CE$ .

## IV. ON PARALLELOGRAMS.

See Propositions 33, 34, and the deductions from these Props. given on page 70.


1. *The straight line drawn through the middle point of a side of a triangle parallel to the base, bisects the remaining side.*

Let  $ABC$  be a  $\triangle$ , and  $Z$  the middle point of the side  $AB$ . Through  $Z$ ,  $ZY$  is drawn  $\text{par}^1$  to  $BC$ .

Then shall  $Y$  be the middle point of  $AC$ .

Through  $Z$  draw  $ZX$   $\text{par}^1$  to  $AC$ . I. 31.

Then in the  $\triangle^s$   $AZY$ ,  $ZBX$ ,  
because  $ZY$  and  $BC$  are  $\text{par}^1$ ,  
 $\therefore$  the  $\angle AZY =$  the  $\angle ZBX$ ; I. 29.  
and because  $ZX$  and  $AC$  are  $\text{par}^1$ ,  
 $\therefore$  the  $\angle ZAY =$  the  $\angle BZX$ ; I. 29.  
also  $AZ = ZB$ : Hyp.  
 $\therefore AY = ZX$ . I. 26.



But  $ZXCY$  is a  $\text{par}^m$  by construction;

$\therefore ZX = YC$ . I. 34.

Hence  $AY = YC$ ;

that is,  $AC$  is bisected at  $Y$ . Q. E. D.

2. *The straight line which joins the middle points of two sides of a triangle, is parallel to the third side.*

Let  $ABC$  be a  $\triangle$ , and  $Z$ ,  $Y$  the middle points of the sides  $AB$ ,  $AC$ .

Then shall  $ZY$  be  $\text{par}^1$  to  $BC$ .

Produce  $ZY$  to  $V$ , making  $YV$  equal to  $ZY$ .

Join  $CV$ .

Then in the  $\triangle^s$   $AYZ$ ,  $CYV$ ,


Because  $\begin{cases} AY = CY, & \text{Hyp.} \\ \text{and } YZ = YV, & \text{Constr.} \\ \text{and the } \angle AYZ = \text{the vert. opp. } \angle CYV; & \text{I. 15.} \\ \therefore AZ = CV, & \text{I. 4.} \\ \text{and the } \angle ZAY = \text{the } \angle VCY; & \text{I. 27.} \\ \text{hence } CV \text{ is } \text{par}^1 \text{ to } AZ. & \text{Hyp.} \end{cases}$

But  $CV$  is equal to  $AZ$ , that is, to  $BZ$ ;

$\therefore CV$  is equal and  $\text{par}^1$  to  $BZ$ ;

$\therefore ZV$  is equal and  $\text{par}^1$  to  $BC$ ;

that is,  $ZY$  is  $\text{par}^1$  to  $BC$ . Q. E. D.



[A second proof of this proposition may be derived from I. 38, 39.]

3. *The straight line which joins the middle points of two sides of a triangle is equal to half the third side.*

4. *Shew that the three straight lines which join the middle points of the sides of a triangle, divide it into four triangles which are identically equal.*

5. *Any straight line drawn from the vertex of a triangle to the base is bisected by the straight line which joins the middle points of the other sides of the triangle.*

6. Given the three middle points of the sides of a triangle, construct the triangle.

7.  $AB, AC$  are two given straight lines, and  $P$  is a given point between them; required to draw through  $P$  a straight line terminated by  $AB, AC$ , and bisected by  $P$ .

8.  $ABCD$  is a parallelogram, and  $X, Y$  are the middle points of the opposite sides  $AD, BC$ : shew that  $BX$  and  $DY$  trisect the diagonal  $AC$ .

9. *If the middle points of adjacent sides of any quadrilateral are joined, the figure thus formed is a parallelogram.*



10. Shew that the straight lines which join the middle points of opposite sides of a quadrilateral, bisect one another.

11. The straight line which joins the middle points of the oblique sides of a trapezium, is parallel to the two parallel sides, and passes through the middle points of the diagonals.

12. *The straight line which joins the middle points of the oblique sides of a trapezium is equal to half the sum of the parallel sides; and the portion intercepted between the diagonals is equal to half the difference of the parallel sides.*


## DEFINITION.

If from the extremities of one straight line perpendiculars are drawn to another, the portion of the latter intercepted between the perpendiculars is said to be the **Orthogonal Projection** of the first line upon the second.



Thus in the adjoining figures, if from the extremities of the straight line  $AB$  the perpendiculars  $AX, BY$  are drawn to  $PQ$ , then  $XY$  is the *orthogonal projection* of  $AB$  on  $PQ$ .

13. *A given straight line AB is bisected at C; shew that the projections of AC, CB on any other straight line are equal.*



Let XZ, ZY be the projections of AC, CB on any straight line PQ.  
Then XZ and ZY shall be equal.

Through A draw a straight line parallel to PQ, meeting CZ, BY or these lines produced in H, K. I. 31.

Now AX, CZ, BY are parallel, for they are perp. to PQ; I. 28.

$\therefore$  the figures XH, HY are par<sup>ms</sup>;

$\therefore$  AH = XZ, and HK = ZY. I. 34.

But through C, the middle point of AB, a side of the  $\triangle$  ABK, CH has been drawn parallel to the side BK;

$\therefore$  CH bisects AK:

Ex. 1, p. 104.

that is, AH = HK;

$\therefore$  XZ = ZY.

Q. E. D.

14. *If three parallel straight lines make equal intercepts on a fourth straight line which meets them, they will also make equal intercepts on any other straight line which meets them.*

15. *Equal and parallel straight lines have equal projections on any other straight line.*

16. *AB is a given straight line bisected at O; and AX, BY are perpendiculars drawn from A and B on any other straight line: shew that OX is equal to OY.*

17. *AB is a given straight line bisected at O: and AX, BY and OZ are perpendiculars drawn to any straight line PQ, which does not pass between A and B: shew that OZ is equal to half the sum of AX, BY.*

[OZ is said to be the **Arithmetic Mean** between AX and BY.]

18. *AB is a given straight line bisected at O; and through A, B and O parallel straight lines are drawn to meet a given straight line PQ in X, Y, Z: shew that OZ is equal to half the sum, or half the difference of AX and BY, according as A and B lie on the same side or on opposite sides of PQ.*

19. *To divide a given finite straight line into any number of equal parts.*


[For example: required to divide the straight line AB into five equal parts.

From A draw AC, a straight line of unlimited length, making any angle with AB.

In AC take any point P; and by marking off successive parts PQ, QR, RS, ST each equal to AP, make AT to contain AP five times.

Join BT; and through P, Q, R, S draw parallels to BT.

It may be shewn by Ex. 14, p. 106, that these parallels divide AB into five equal parts.]



20. *If through an angle of a parallelogram any straight line is drawn, the perpendicular drawn to it from the opposite angle is equal to the sum or difference of the perpendiculars drawn to it from the two remaining angles, according as the given straight line falls without the parallelogram, or intersects it.*

[Through the opposite angle draw a straight line parallel to the given straight line, so as to meet the perpendicular from one of the remaining angles, produced if necessary; then apply I. 34, I. 26. Or proceed as in the following example.]

21. From the angular points of a parallelogram perpendiculars are drawn to any straight line which is without the parallelogram: shew that the sum of the perpendiculars drawn from one pair of opposite angles is equal to the sum of those drawn from the other pair.

[Draw the diagonals, and from their point of intersection let fall a perpendicular upon the given straight line. See Ex. 17, p. 106.]

22. The sum of the perpendiculars drawn from any point in the base of an isosceles triangle to the equal sides is equal to the perpendicular drawn from either extremity of the base to the opposite side.

[It follows that the sum of the distances of any point in the base of an isosceles triangle from the equal sides is constant, that is, the same whatever point in the base is taken.]

23. In the base produced of an isosceles triangle any point is taken: shew that the difference of its perpendicular distances from the equal sides is constant.

24. The sum of the perpendiculars drawn from any point within an equilateral triangle to the three sides is equal to the perpendicular drawn from any one of the angular points to the opposite side, and is therefore constant.

## PROBLEMS.

25. Draw a straight line through a given point, so that the part of it intercepted between two given parallel straight lines may be of given length. When does this problem admit of two solutions, when of only one, and when is it impossible?

26. Draw a straight line parallel to a given straight line, so that the part intercepted between two other given straight lines may be of given length.

27. Draw a straight line equally inclined to two given straight lines that meet, so that the part intercepted between them may be of given length.

28.  $AB, AC$  are two given straight lines, and  $P$  is a given point *without* the angle contained by them. It is required to draw through  $P$  a straight line to meet the given lines, so that the part intercepted between them may be equal to the part between  $P$  and the nearer line.

## V. MISCELLANEOUS THEOREMS AND EXAMPLES.

Chiefly on I. 32.

1.  $A$  is the vertex of an isosceles triangle  $ABC$ , and  $BA$  is produced to  $D$ , so that  $AD$  is equal to  $BA$ ; if  $DC$  is drawn, shew that  $BCD$  is a right angle.

2. The straight line joining the middle point of the hypotenuse of a right-angled triangle to the right angle is equal to half the hypotenuse.

3. From the extremities of the base of a triangle perpendiculars are drawn to the opposite sides (produced if necessary); shew that the straight lines which join the middle point of the base to the feet of the perpendiculars are equal.

4. In a triangle  $ABC$ ,  $AD$  is drawn perpendicular to  $BC$ ; and  $X, Y, Z$  are the middle points of the sides  $BC, CA, AB$  respectively: shew that each of the angles  $ZXY, ZDY$  is equal to the angle  $BAC$ .

5. In a right-angled triangle, if a perpendicular is drawn from the right angle to the hypotenuse, the two triangles thus formed are equiangular to one another.


6. In a right-angled triangle two straight lines are drawn from the right angle, one bisecting the hypotenuse, the other perpendicular to it: shew that they contain an angle equal to the difference of the two acute angles of the triangle. [See above, Ex. 2 and Ex. 5.]

7. In a triangle if a perpendicular is drawn from one extremity of the base to the bisector of the vertical angle, (i) it will make with either of the sides containing the vertical angle an angle equal to half the sum of the angles at the base; (ii) it will make with the base an angle equal to half the difference of the angles at the base.

Let  $ABC$  be the given  $\triangle$ , and  $AH$  the bisector of the vertical  $\angle BAC$ .

Let  $CLK$  meet  $AH$  at right angles.

(i) Then shall each of the  $\angle^s$   $AKC$ ,  $ACK$  be equal to half the sum of the  $\angle^s$   $ABC$ ,  $ACB$ .



In the  $\triangle^s$   $AKL$ ,  $ACL$ ,  
 Because  $\left\{ \begin{array}{l} \text{the } \angle KAL = \text{the } \angle CAL, \\ \text{also the } \angle ALK = \text{the } \angle ALC, \text{ being rt. } \angle^s; \\ \text{and } AL \text{ is common to both } \triangle^s; \end{array} \right.$  Hyp.  
 $\therefore$  the  $\angle AKL = \text{the } \angle ACL$ . I. 26.

Again, the  $\angle AKC = \text{the sum of the } \angle^s$   $KBC$ ,  $KCB$ ; I. 32.  
 $\therefore$  the  $\angle ACK = \text{the sum of the } \angle^s$   $KBC$ ,  $KCB$ .

To each add the  $\angle ACK$ :  
 then twice the  $\angle ACK = \text{the sum of the } \angle^s$   $ABC$ ,  $ACB$ ;  
 $\therefore$  the  $\angle ACK = \text{half the sum of the } \angle^s$   $ABC$ ,  $ACB$ .

(ii) The  $\angle KCB$  shall be equal to half the difference of the  $\angle^s$   $ACB$ ,  $ABC$ .

As before, the  $\angle ACK = \text{the sum of the } \angle^s$   $KBC$ ,  $KCB$ .

To each of these add the  $\angle KCB$ :

then the  $\angle ACB = \text{the } \angle KBC$  together with twice the  $\angle KCB$ .

$\therefore$  twice the  $\angle KCB = \text{the difference of the } \angle^s$   $ACB$ ,  $KBC$ ;

that is, the  $\angle KCB = \text{half the difference of the } \angle^s$   $ACB$ ,  $ABC$ .

**COROLLARY.** If  $X$  is the middle point of the base, and  $XL$  is joined, it may be shewn by Ex. 3, p. 105, that  $XL$  is half  $BK$ ; that is, that  $XL$  is half the difference of the sides  $AB$ ,  $AC$ .

8. In any triangle the angle contained by the bisector of the vertical angle and the perpendicular from the vertex to the base is equal to half the difference of the angles at the base. [See Ex. 3, p. 65.]

9. In a triangle  $ABC$  the side  $AC$  is produced to  $D$ , and the angles  $BAC$ ,  $BCD$  are bisected by straight lines which meet at  $F$ ; shew that they contain an angle equal to half the angle at  $B$ .

10. If in a right-angled triangle one of the acute angles is double of the other, shew that the hypotenuse is double of the shorter side.



11. If in a diagonal of a parallelogram any two points equidistant from its extremities are joined to the opposite angles, the figure thus formed will be also a parallelogram.

12.  $ABC$  is a given equilateral triangle, and in the sides  $BC$ ,  $CA$ ,  $AB$  the points  $X$ ,  $Y$ ,  $Z$  are taken respectively, so that  $BX$ ,  $CY$  and  $AZ$  are all equal.  $AX$ ,  $BY$ ,  $CZ$  are now drawn, intersecting in  $P$ ,  $Q$ ,  $R$ : shew that the triangle  $PQR$  is equilateral.

13. If in the sides  $AB$ ,  $BC$ ,  $CD$ ,  $DA$  of a parallelogram  $ABCD$  four points  $P$ ,  $Q$ ,  $R$ ,  $S$  are taken in order, one in each side, so that  $AP$ ,  $BQ$ ,  $CR$ ,  $DS$  are all equal; shew that the figure  $PQRS$  is a parallelogram.

14. In the figure of *i. 1*, if the circles intersect at  $F$ , and if  $CA$  and  $CB$  are produced to meet the circles in  $P$  and  $Q$  respectively; shew that the points  $P$ ,  $F$ ,  $Q$  are in the same straight line; and shew also that the triangle  $CPQ$  is equilateral.

[Problems marked (\*) admit in general of more than one solution.]

15. Through two given points draw two straight lines forming with a straight line given in position, an equilateral triangle.

\*16. From a given point it is required to draw to two parallel straight lines two equal straight lines at right angles to one another.

\*17. Three given straight lines meet at a point; draw another straight line so that the two portions of it intercepted between the given lines may be equal to one another.

18. From a given point draw three straight lines of given lengths, so that their extremities may be in the same straight line, and intercept equal distances on that line. [See Fig. to *i. 16*.]

19. Use the properties of the equilateral triangle to trisect a given finite straight line.

20. In a given triangle inscribe a rhombus, having one of its angles coinciding with an angle of the triangle.

## VI. ON THE CONCURRENCE OF STRAIGHT LINES IN A TRIANGLE.

DEFINITIONS. (i) Three or more straight lines are said to be **concurrent** when they meet in one point.

(ii) Three or more points are said to be **collinear** when they lie upon one straight line.

*Obs.* We here give some propositions relating to the concurrence of certain groups of straight lines drawn in a triangle: the importance of these theorems will be more fully appreciated when the student is familiar with Books III. and IV.


1. *The perpendiculars drawn to the sides of a triangle from their middle points are concurrent.*

Let  $ABC$  be a  $\triangle$ , and  $X, Y, Z$  the middle points of its sides.

Then shall the perp<sup>s</sup> drawn to the sides from  $X, Y, Z$  be concurrent.

From  $Z$  and  $Y$  draw perp<sup>s</sup> to  $AB, AC$ ; these perp<sup>s</sup>, since they cannot be parallel, will meet at some point  $O$ . Ax. 12.

Join  $OX$ .



*It is required to prove that  $OX$  is perp. to  $BC$ .*

Join  $OA, OB, OC$ .

In the  $\triangle^s OYA, OYC$ ,

$YA = YC$ ,

and  $OY$  is common to both;

also the  $\angle OYA =$  the  $\angle OYC$ , being rt.  $\angle^s$ ;

$\therefore OA = OC$ .

*Hyp.*

*I. 4.*

Similarly, from the  $\triangle^s OZA, OZB$ ,

it may be proved that  $OA = OB$ .

Hence  $OA, OB, OC$  are all equal.

Again, in the  $\triangle^s OXB, OXC$ ,

$BX = CX$ ,

and  $XO$  is common to both;

also  $OB = OC$ ;

$\therefore$  the  $\angle OXB =$  the  $\angle OXC$ ;

but these are adjacent  $\angle^s$ ;

$\therefore$  they are rt.  $\angle^s$ ;

that is,  $OX$  is perp. to  $BC$ .

*Hyp.*

*Proved.*

*I. 8.*

*Def. 10.*

Hence the three perp<sup>s</sup>  $OX, OY, OZ$  meet at the point  $O$ .

*Q. E. D.*

2. *The bisectors of the angles of a triangle are concurrent.*

Let  $ABC$  be a  $\triangle$ . Bisect the  $\angle^s ABC, BCA$ , by str. right lines which must meet at some point  $O$ . Ax. 12.

Join  $AO$ .

*It is required to prove that  $AO$  bisects the  $\angle BAC$ .*

From  $O$  draw  $OP, OQ, OR$  perp. to the sides of the  $\triangle$ .

Then in the  $\triangle^s OBP, OBR$ ,

the  $\angle OBP =$  the  $\angle OBR$ ,


and the  $\angle OPB =$  the  $\angle ORB$ , being rt.  $\angle^s$ ,

and  $OB$  is common;

$\therefore OP = OR$ .


*Constr.*

*I. 26*



Similarly from the  $\triangle^s$  OCP, OCQ,  
it may be shewn that  $OP=OQ$ ,  
 $\therefore OP, OQ, OR$  are all equal.

Again in the  $\triangle^s$  ORA, OQA,  
Because  $\left\{ \begin{array}{l} \text{the } \angle^s \text{ ORA, OQA are rt. } \angle^s, \\ \text{and the hypotenuse OA is} \\ \text{common,} \\ \text{also } OR=OQ; \text{ Proved.} \end{array} \right.$   
 $\therefore$  the  $\angle$  RAO = the  $\angle$  QAO.



Ex. 12, p. 99.

That is, AO is the bisector of the  $\angle$  BAC.

Hence the bisectors of the three  $\angle^s$  meet at the point O.

Q. E. D.

3. *The bisectors of two exterior angles of a triangle and the bisector of the third angle are concurrent.*

Let ABC be a  $\triangle$ , of which the sides AB, AC are produced to any points D and E.


Then shall the bisectors of the  $\angle^s$  DBC, ECB, BAC be concurrent.

Bisect the  $\angle^s$  DBC, ECB by straight lines which must meet at some point O. Ax. 12.

Join AO.

It is required to prove that AO bisects the angle BAC.

From O draw OP, OQ, OR perp. to the sides of the  $\triangle$ .



Constr.

Then in the  $\triangle^s$  OBP, OBR,  
Because  $\left\{ \begin{array}{l} \text{the } \angle \text{ OBP} = \text{the } \angle \text{ OBR,} \\ \text{also the } \angle \text{ OPB} = \text{the } \angle \text{ ORB, being rt. } \angle^s, \\ \text{and OB is common;} \end{array} \right.$   
 $\therefore OP=OR$ .

I. 26.

Similarly from the  $\triangle^s$  OCP, OCQ,  
it may be shewn that  $OP=OQ$ :  
 $\therefore OP, OQ, OR$  are all equal.

Again in the  $\triangle^s$  ORA, OQA,  
Because  $\left\{ \begin{array}{l} \text{the } \angle^s \text{ ORA, OQA are rt. } \angle^s, \\ \text{and the hypotenuse OA is common,} \\ \text{also } OR=OQ; \end{array} \right.$   
 $\therefore$  the  $\angle$  RAO = the  $\angle$  QAO.

Proved.

Ex. 12, p. 99.

That is, AO is the bisector of the  $\angle$  BAC.

$\therefore$  the bisectors of the two exterior  $\angle^s$  DBC, ECB, and of the interior  $\angle$  BAC meet at the point O.

Q. E. D.

4. *The medians of a triangle are concurrent.*

Let  $ABC$  be a  $\triangle$ .

Then shall its three medians be concurrent.

Let  $BY$  and  $CZ$  be two of its medians, and let them intersect at  $O$ .

Join  $AO$ ,


and produce it to meet  $BC$  in  $X$ .

It is required to shew that  $AX$  is the remaining median of the  $\triangle$ .

Through  $C$  draw  $CK$  parallel to  $BY$  :

produce  $AX$  to meet  $CK$  at  $K$ .

Join  $BK$ .



In the  $\triangle AKC$ ,

because  $Y$  is the middle point of  $AC$ , and  $YO$  is parallel to  $CK$ ,

$\therefore O$  is the middle point of  $AK$ . Ex. 1, p. 104.

Again in the  $\triangle ABK$ ,

since  $Z$  and  $O$  are the middle points of  $AB$ ,  $AK$ ,

$\therefore ZO$  is parallel to  $BK$ , Ex. 2, p. 104.

that is,  $OC$  is parallel to  $BK$  :

$\therefore$  the figure  $BKCO$  is a par<sup>m</sup>.

But the diagonals of a par<sup>m</sup> bisect one another, Ex. 5, p. 70.

$\therefore X$  is the middle point of  $BC$ .

That is,  $AX$  is a median of the  $\triangle$ .

Hence the three medians meet at the point  $O$ . Q.E.D.

**COROLLARY.** *The three medians of a triangle cut one another at a point of trisection, the greater segment in each being towards the angular point.*

For in the above figure it has been proved that

$$AO = OK,$$

also that  $OX$  is half of  $OK$  ;

$\therefore OX$  is half of  $OA$  :

that is,  $OX$  is one third of  $AX$ .

Similarly  $OY$  is one third of  $BY$ ,


and  $OZ$  is one third of  $CZ$ .

Q.E.D.

By means of this Corollary it may be shewn that in any triangle the shorter median bisects the greater side.

[The point of intersection of the three medians of a triangle is called the **centroid**. It is shewn in Mechanics that a thin triangular plate will balance in any position about this point: therefore the centroid of a triangle is also its centre of gravity.]

5. *The perpendiculars drawn from the vertices of a triangle to the opposite sides are concurrent.*



Let  $ABC$  be a  $\triangle$ , and  $AD$ ,  $BE$ ,  $CF$  the three perp<sup>s</sup> drawn from the vertices to the opposite sides.

*Then shall the perp<sup>s</sup>  $AD$ ,  $BE$ ,  $CF$  be concurrent.*

Through  $A$ ,  $B$ , and  $C$  draw straight lines  $MN$ ,  $NL$ ,  $LM$  parallel to the opposite sides of the  $\triangle$ .

Then the figure  $BAMC$  is a par<sup>m</sup>.

*Def. 36.*

$\therefore AB = MC$ .

*I. 34.*

Also the figure  $BACL$  is a par<sup>m</sup>.

$\therefore AB = LC$ ,

$\therefore LC = CM$ ;

that is,  $C$  is the middle point of  $LM$ .

So also  $A$  and  $B$  are the middle points of  $MN$  and  $NL$ .

Hence  $AD$ ,  $BE$ ,  $CF$  are the perp<sup>s</sup> to the sides of the  $\triangle LMN$  from their middle points.

*Ex. 3, p. 60.*

But these perp<sup>s</sup> meet in a point :

*Ex. 1, p. 111.*

that is, the perp<sup>s</sup> drawn from the vertices of the  $\triangle ABC$  to the opposite sides meet in a point.

*Q. E. D.*

[For another proof see Theorems and Examples on Book III.]

#### DEFINITIONS.

(i) The intersection of the perpendiculars drawn from the vertices of a triangle to the opposite sides is called its **orthocentre**.

(ii) The triangle formed by joining the feet of the perpendiculars is called the **pedal triangle**.


### VII. ON THE CONSTRUCTION OF TRIANGLES WITH GIVEN PARTS.

*Obs.* No general rules can be laid down for the solution of problems in this section; but in a few typical cases we give constructions, which the student will find little difficulty in adapting to other questions of the same class.

1. Construct a right-angled triangle, having given the hypotenuse and the sum of the remaining sides.

[It is required to construct a rt.-angled  $\Delta$ , having its hypotenuse equal to the given straight line  $K$ , and the sum of its remaining sides equal to  $AB$ .

From  $A$  draw  $AE$  making with  $BA$  an  $\angle$  equal to half a rt.  $\angle$ . From centre  $B$ , with radius equal to  $K$ , describe a circle cutting  $AE$  in the points  $C, C'$ .




From  $C$  and  $C'$  draw perp<sup>s</sup>  $CD, C'D'$  to  $AB$ ; and join  $CB, C'B$ . Then either of the  $\Delta^s$   $CDB, C'D'B$  will satisfy the given conditions.

**NOTE.** If the given hypotenuse  $K$  be greater than the perpendicular drawn from  $B$  to  $AE$ , there will be *two* solutions. If the line  $K$  be equal to this perpendicular, there will be *one* solution; but if less, the problem is *impossible*.]

2. Construct a right-angled triangle, having given the hypotenuse and the difference of the remaining sides.

3. Construct an isosceles right-angled triangle, having given the sum of the hypotenuse and one side.

4. Construct a triangle, having given the perimeter and the angles at the base.



[Let  $AB$  be the perimeter of the required  $\Delta$ , and  $X$  and  $Y$  the  $\angle^s$  at the base.

From  $A$  draw  $AP$ , making the  $\angle BAP$  equal to half the  $\angle X$ .

From  $B$  draw  $BP$ , making the  $\angle ABP$  equal to half the  $\angle Y$ .

From  $P$  draw  $PQ$ , making the  $\angle APQ$  equal to the  $\angle BAP$ .

From  $P$  draw  $PR$ , making the  $\angle BPR$  equal to the  $\angle ABP$ .

Then shall  $PQR$  be the required  $\Delta$ .]

5. Construct a right-angled triangle, having given the perimeter and one acute angle.

6. Construct an isosceles triangle of given altitude, so that its base may be in a given straight line, and its two equal sides may pass through two fixed points. [See Ex. 7, p. 55.]


7. Construct an equilateral triangle, having given the length of the perpendicular drawn from one of the vertices to the opposite side.

8. Construct an isosceles triangle, having given the base, and the difference of one of the remaining sides and the perpendicular drawn from the vertex to the base. [See Ex. 1, p. 96.]

9. Construct a triangle, having given the base, one of the angles at the base, and the sum of the remaining sides.

10. Construct a triangle, having given the base, one of the angles at the base, and the difference of the remaining sides. [Two cases arise, according as the given angle is adjacent to the greater side or the less.]

11. Construct a triangle, having given the base, the difference of the angles at the base, and the difference of the remaining sides.



[Let  $AB$  be the given base,  $X$  the difference of the  $\angle$ 's at the base, and  $K$  the difference of the remaining sides.

Draw  $BE$ , making the  $\angle ABE$  equal to half the  $\angle X$ .

From centre  $A$ , with radius equal to  $K$ , describe a circle cutting  $BE$  in  $D$  and  $D'$ . Let  $D$  be the point of intersection nearer to  $B$ .

Join  $AD$  and produce it to  $C$ .

Draw  $BC$ , making the  $\angle DBC$  equal to the  $\angle BDC$ .

Then shall  $CAB$  be the  $\triangle$  required. Ex. 7, p. 109.

NOTE. This problem is possible only when the given difference  $K$  is greater than the perpendicular drawn from  $A$  to  $BE$ .]

12. Construct a triangle, having given the base, the difference of the angles at the base, and the sum of the remaining sides.

13. Construct a triangle, having given the perpendicular from the vertex on the base, and the difference between each side and the adjacent segment of the base.

14. Construct a triangle, having given two sides and the median which bisects the remaining side. [See Ex. 18, p. 110.]

15. Construct a triangle, having given one side, and the medians which bisect the two remaining sides.

[See Fig. to Ex. 4, p. 113.]

Let BC be the given side. Take two-thirds of each of the given medians; hence construct the triangle BOC. The rest of the construction follows easily.]

16. Construct a triangle, having given its three medians.

[See Fig. to Ex. 4, p. 113.]

Take two-thirds of each of the given medians, and construct the triangle OKC. The rest of the construction follows easily.]

### VIII. ON AREAS.

See Propositions 35—48.

*Obs.* It must be understood that throughout this section the word *equal* as applied to rectilinear figures will be used as denoting equality of area unless otherwise stated.

1. Shew that a parallelogram is bisected by any straight line which passes through the middle point of one of its diagonals.

[I. 29, 26.]

2. Bisect a parallelogram by a straight line drawn through a given point.

3. Bisect a parallelogram by a straight line drawn perpendicular to one of its sides.

4. Bisect a parallelogram by a straight line drawn parallel to a given straight line.

5. ABCD is a trapezium in which the side AB is parallel to DC. Shew that its area is equal to the area of a parallelogram formed by drawing through X, the middle point of BC, a straight line parallel to AD, meeting DC, or DC produced.

[I. 29, 26.]

6. A trapezium is equal to a parallelogram whose base is half the sum of the parallel sides of the given figure, and whose altitude is equal to the perpendicular distance between them.

7. ABCD is a trapezium in which the side AB is parallel to DC; shew that it is double of the triangle formed by joining the extremities of AD to X, the middle point of BC.

8. Shew that a trapezium is bisected by the straight line which joins the middle points of its parallel sides.

[I. 38.]



*Obs.* In the following group of Exercises the proofs depend chiefly on Propositions 37 and 38, and the two converse theorems.

9. If two straight lines  $AB$ ,  $CD$  intersect at  $X$ , and if the straight lines  $AC$  and  $BD$ , which join their extremities are parallel, shew that the triangle  $AXD$  is equal to the triangle  $BXC$ .

10. If two straight lines  $AB$ ,  $CD$  intersect at  $X$ , so that the triangle  $AXD$  is equal to the triangle  $XCB$ , then  $AC$  and  $BD$  are parallel.

11.  $ABCD$  is a parallelogram, and  $X$  any point in the diagonal  $AC$  produced; shew that the triangles  $XBC$ ,  $XDC$  are equal. [See Ex. 13, p. 70.]

12.  $ABC$  is a triangle, and  $R$ ,  $Q$  the middle points of the sides  $AB$ ,  $AC$ ; shew that if  $BQ$  and  $CR$  intersect in  $X$ , the triangle  $BXC$  is equal to the quadrilateral  $AQXR$ . [See Ex. 5, p. 79.]

13. If the middle points of the sides of a quadrilateral be joined in order, the *parallelogram* so formed [see Ex. 9, p. 105] is equal to half the given figure.

14. Two triangles of equal area stand on the same base but on opposite sides of it: shew that the straight line joining their vertices is bisected by the base, or by the base produced.

15. The straight line which joins the middle points of the diagonals of a trapezium is parallel to each of the two parallel sides.

16. (i) *A triangle is equal to the sum or difference of two triangles on the same base (or on equal bases), if the altitude of the first is equal to the sum or difference of the altitudes of the others.*

(ii) *A triangle is equal to the sum or difference of two triangles of the same altitude, if the base of the first is equal to the sum or difference of the bases of the others.*

Similar statements hold good of parallelograms.

17.  $ABCD$  is a parallelogram, and  $O$  is any point outside it; shew that the sum or difference of the triangles  $OAB$ ,  $OCD$  is equal to half the parallelogram. Distinguish between the two cases.

*Obs.* On the following proposition depends an important theorem in Mechanics: we give a proof of the first case, leaving the second case to be deduced by a similar method.


18. (i)  $ABCD$  is a parallelogram, and  $O$  is any point without the angle  $BAD$  and its opposite vertical angle; shew that the triangle  $OAC$  is equal to the sum of the triangles  $OAD$ ,  $OAB$ .

(ii) If  $O$  is within the angle  $BAD$  or its opposite vertical angle, the triangle  $OAC$  is equal to the difference of the triangles  $OAD$ ,  $OAB$ .

CASE I. If  $O$  is without the  $\angle DAB$  and its opp. vert.  $\angle$ , then  $OA$  is without the par<sup>m</sup>  $ABCD$ : therefore the perp. drawn from  $C$  to  $OA$  is equal to the sum of the perp<sup>s</sup> drawn from  $B$  and  $D$  to  $OA$ . [See Ex. 20, p. 107.]

Now the  $\triangle^s$   $OAC$ ,  $OAD$ ,  $OAB$  are upon the same base  $OA$ ; and the altitude of the  $\triangle$   $OAC$  with respect to this base has been shewn to be equal to the sum of the altitudes of the  $\triangle^s$   $OAD$ ,  $OAB$ .


Therefore the  $\triangle$   $OAC$  is equal to the sum of the  $\triangle^s$   $OAD$ ,  $OAB$ . [See Ex. 16, p. 118.] Q. E. D.



19.  $ABCD$  is a parallelogram, and through  $O$ , any point within it, straight lines are drawn parallel to the sides of the parallelogram; shew that the difference of the parallelograms  $DO$ ,  $BO$  is double of the triangle  $AOC$ . [See preceding theorem (ii).]

20. The area of a quadrilateral is equal to the area of a triangle having two of its sides equal to the diagonals of the given figure, and the included angle equal to either of the angles between the diagonals.

21.  $ABC$  is a triangle, and  $D$  is any point in  $AB$ ; it is required to draw through  $D$  a straight line  $DE$  to meet  $BC$  produced in  $E$ , so that the triangle  $DBE$  may be equal to the triangle  $ABC$ .



[Join  $DC$ . Through  $A$  draw  $AE$  parallel to  $DC$ . I. 31.  
Join  $DE$ .

The  $\triangle$   $EBD$  shall be equal to the  $\triangle$   $ABC$ .] I. 37.

22. On a base of given length describe a triangle equal to a given triangle and having an angle equal to an angle of the given triangle.

23. Construct a triangle equal in area to a given triangle, and having a given altitude.

24. On a base of given length construct a triangle equal to a given triangle, and having its vertex on a given straight line.

25. On a base of given length describe (i) an isosceles triangle ; (ii) a right-angled triangle, equal to a given triangle.

26. Construct a triangle equal to the sum or difference of two given triangles. [See Ex. 16, p. 118.]

27.  $ABC$  is a given triangle, and  $X$  a given point: describe a triangle equal to  $ABC$ , having its vertex at  $X$ , and its base in the same straight line as  $BC$ .

28.  $ABCD$  is a quadrilateral. On the base  $AB$  construct a triangle equal in area to  $ABCD$ , and having the angle at  $A$  common with the quadrilateral.

[Join  $BD$ . Through  $C$  draw  $CX$  parallel to  $BD$ , meeting  $AD$  produced in  $X$ ; join  $BX$ .]

29. Construct a rectilineal figure equal to a given rectilineal figure, and having fewer sides by one than the given figure.

Hence shew how to construct a triangle equal to a given rectilineal figure.

30.  $ABCD$  is a quadrilateral: it is required to construct a triangle equal in area to  $ABCD$ , having its vertex at a given point  $X$  in  $DC$ , and its base in the same straight line as  $AB$ .

31. Construct a rhombus equal to a given parallelogram.

32. Construct a parallelogram which shall have the same area and perimeter as a given triangle.

33. Bisect a triangle by a straight line drawn through one of its angular points.

34. Trisect a triangle by straight lines drawn through one of its angular points. [See Ex. 19, p. 110, and i. 38.]

35. Divide a triangle into any number of equal parts by straight lines drawn through one of its angular points. [See Ex. 19, p. 107, and i. 38.]

36. *Bisect a triangle by a straight line drawn through a given point in one of its sides.*


[Let  $ABC$  be the given  $\triangle$ , and  $P$  the given point in the side  $AB$ .

Bisect  $AB$  at  $Z$ ; and join  $CZ$ ,  $CP$ .  
Through  $Z$  draw  $ZQ$  parallel to  $CP$ .

Join  $PQ$ .

Then shall  $PQ$  bisect the  $\triangle$ .

[See Ex. 21, p. 119.]



37. *Trisect a triangle by straight lines drawn from a given point in one of its sides.*


[Let  $ABC$  be the given  $\triangle$ , and  $X$  the given point in the side  $BC$ .

Trisect  $BC$  at the points  $P$ ,  $Q$ . Ex. 19, p. 107.  
Join  $AX$ , and through  $P$  and  $Q$  draw  $PH$  and  $QK$  parallel to  $AX$ .

Join  $XH$ ,  $XK$ .

These straight lines shall trisect the  $\triangle$ ; as may be shewn by joining  $AP$ ,  $AQ$ .

[See Ex. 21, p. 119.]



38. *Cut off from a given triangle a fourth, fifth, sixth, or any part required by a straight line drawn from a given point in one of its sides.*  
[See Ex. 19, p. 107, and Ex. 21, p. 119.]

39. *Bisect a quadrilateral by a straight line drawn through an angular point.*

[Two constructions may be given for this problem: the first will be suggested by Exercises 28 and 33, p. 120.


The second method proceeds thus.

Let  $ABCD$  be the given quadrilateral, and  $A$  the given angular point.

Join  $AC$ ,  $BD$ , and bisect  $BD$  in  $X$ .  
Through  $X$  draw  $PXQ$  parallel to  $AC$ , meeting  $BC$  in  $P$ ; join  $AP$ .

Then shall  $AP$  bisect the quadrilateral.

Join  $AX$ ,  $CX$ , and use i. 37, 38.]



40. *Cut off from a given quadrilateral a third, a fourth, a fifth, or any part required, by a straight line drawn through a given angular point.*  
[See Exercises 28 and 35, p. 120.]

*Obs.* The following Theorems depend on I. 47.

41. In the figure of I. 47, shew that
- (i) the sum of the squares on AB and AE is equal to the sum of the squares on AC and AD.
  - (ii) the square on EK is equal to the square on AB with four times the square on AC.
  - (iii) the sum of the squares on EK and FD is equal to five times the square on BC.
42. If a straight line is divided into any two parts, the square on the straight line is greater than the sum of the squares on the two parts.
43. If the square on one side of a triangle is less than the squares on the remaining sides, the angle contained by these sides is acute; if greater, obtuse.
44. ABC is a triangle, right-angled at A; the sides AB, AC are intersected by a straight line PQ, and BQ, PC are joined: shew that the sum of the squares on BQ, PC is equal to the sum of the squares on BC, PQ.
45. In a right-angled triangle four times the sum of the squares on the medians which bisect the sides containing the right angle is equal to five times the square on the hypotenuse.
46. Describe a square whose area shall be three times that of a given square.
47. Divide a straight line into two parts such that the sum of their squares shall be equal to a given square.

## IX. ON LOCI.

In many geometrical problems we are required to find the position of a point which satisfies given conditions; and all such problems hitherto considered have been found to admit of a *limited number* of solutions. This, however, will not be the case if *only one* condition is given. For example:

(i) *Required a point which shall be at a given distance from a given point.*

This problem is evidently *indeterminate*, that is to say, it admits of an indefinite number of solutions; for the condition stated is satisfied by *any* point on the circumference of the circle described from the given point as centre, with a radius equal to the given distance. Moreover this condition is satisfied by no other point within or without the circle.

(ii) *Required a point which shall be at a given distance from a given straight line.*

Here again there are an infinite number of such points, and they lie on two parallel straight lines drawn on either side of the given straight line at the given distance from it: further, no point that is not on one or other of these parallels satisfies the given condition.

Hence we see that *one* condition is not sufficient to determine the position of a point absolutely, but it may have the effect of restricting it to some definite line or lines, straight or curved. This leads us to the following definition.

**DEFINITION.** The **Locus** of a point satisfying an assigned condition consists of the line, lines, or part of a line, to which the point is thereby restricted; provided that the condition is satisfied by every point on such line or lines, and by no other.

A locus is sometimes defined as the path traced out by a point which moves in accordance with an assigned law.

Thus the locus of a point, which is always at a given distance from a given point, is a circle of which the given point is the centre: and the locus of a point, which is always at a given distance from a given straight line, is a pair of parallel straight lines.

We now see that in order to infer that a certain line, or system of lines, is the locus of a point under a given condition, it is necessary to prove

(i) that any point which fulfils the given condition is on the supposed locus;

(ii) that every point on the supposed locus satisfies the given condition.

1. *Find the locus of a point which is always equidistant from two given points.*

Let A, B be the two given points.

(a) Let P be any point equidistant from A and B, so that  $AP = BP$ .

Bisect AB at X, and join PX.

Then in the  $\triangle AXP, BXP$ ,


Because  $\left\{ \begin{array}{l} AX = BX, \\ \text{and } PX \text{ is common to both,} \\ \text{also } AP = BP, \end{array} \right.$  *Constr.*

$\therefore$  the  $\angle PXA = \text{the } \angle PXB$ ; *Hyp.*

and they are adjacent  $\angle^s$ ; *I. 8.*

$\therefore$  PX is perp. to AB. *Def. 10.*

$\therefore$  any point which is equidistant from A and B is on the straight line which bisects AB at right angles.



( $\beta$ ) Also every point in this line is equidistant from A and B.

For let Q be any point in this line.

Join AQ, BQ.

Then in the  $\triangle$  AXQ, BXQ,

AX = BX,

and XQ is common to both ;

Because { also the  $\angle$  AXQ = the  $\angle$  BXQ, being rt.  $\angle$ 's ;


$\therefore$  AQ = BQ.

I. 4.

That is, Q is equidistant from A and B.

Hence we conclude that the locus of the point equidistant from two given points A, B is the straight line which bisects AB at right angles.

2. To find the locus of the middle point of a straight line drawn from a given point to meet a given straight line of unlimited length.



Let A be the given point, and BC the given straight line of unlimited length.

(a) Let AX be any straight line drawn through A to meet BC, and let P be its middle point.

Draw AF perp. to BC, and bisect AF at E.

Join EP, and produce it indefinitely.

Since AFX is a  $\triangle$ , and E, P the middle points of the two sides AF, AX,

$\therefore$  EP is parallel to the remaining side FX. Ex. 2, p. 104.

$\therefore$  P is on the straight line which passes through the fixed point E, and is parallel to BC.

( $\beta$ ) Again, every point in EP, or EP produced, fulfils the required condition.

For, in this straight line take any point Q.

Join AQ, and produce it to meet BC in Y.

Then FAY is a  $\triangle$ , and through E, the middle point of the side AF,

EQ is drawn parallel to the side FY ;

$\therefore$  Q is the middle point of AY. Ex. 1, p. 104.

Hence the required locus is the straight line drawn parallel to BC, and passing through E, the middle point of the perp. from A to BC.

3. Find the locus of a point equidistant from two given intersecting straight lines. [See Ex. 3, p. 55.]

4. Find the locus of a point at a given radial distance from the circumference of a given circle.

5. Find the locus of a point which moves so that the sum of its distances from two given intersecting straight lines of unlimited length is constant.

6. Find the locus of a point when the differences of its distances from two given intersecting straight lines of unlimited length is constant.

7. A straight rod of given length slides between two straight rulers placed at right angles to one another: find the locus of its middle point. [See Ex. 2, p. 108.]

8. On a given base as hypotenuse right-angled triangles are described: find the locus of their vertices. [See Ex. 2, p. 108.]

9. AB is a given straight line, and AX is the perpendicular drawn from A to any straight line passing through B: find the locus of the middle point of AX.

10. Find the locus of the vertex of a triangle, when the base and area are given.

11. Find the locus of the intersection of the diagonals of a parallelogram, of which the base and area are given.

12. Find the locus of the intersection of the medians of triangles described on a given base and of given area.

#### X. ON THE INTERSECTION OF LOCI.

It appears from various problems which have already been considered, that we are often required to find a point, the position of which is subject to two given conditions. The method of loci is very useful in solving problems of this kind; for corresponding to each condition there will be a locus on which the required point must lie. Hence all points which are common to these two loci, that is, all the points of intersection of the loci, will satisfy *both* the given conditions.



**EXAMPLE 1.** *To construct a triangle, having given the base, the altitude, and the length of the median which bisects the base.*

Let  $AB$  be the given base, and  $P$  and  $Q$  the lengths of the altitude and median respectively :

then the triangle is known if its vertex is known.

(i) Draw a straight line  $CD$  parallel to  $AB$ , and at a distance from it equal to  $P$  :

then the required vertex must lie on  $CD$ .

(ii) Again, from the middle point of  $AB$  as centre, with radius equal to  $Q$ , describe a circle :

then the required vertex must lie on this circle.

Hence any points which are common to  $CD$  and the circle, satisfy both the given conditions : that is to say, if  $CD$  intersect the circle in  $E, F$  each of the points of intersection might be the vertex of the required triangle. This supposes the length of the median  $Q$  to be greater than the altitude.

**EXAMPLE 2.** *To find a point equidistant from three given points  $A, B, C$ , which are not in the same straight line.*

(i) The locus of points equidistant from  $A$  and  $B$  is the straight line  $PQ$ , which bisects  $AB$  at right angles. Ex. 1, p. 123.

(ii) Similarly the locus of points equidistant from  $B$  and  $C$  is the straight line  $RS$  which bisects  $BC$  at right angles.

Hence the point common to  $PQ$  and  $RS$  must satisfy both conditions : that is to say, the point of intersection of  $PQ$  and  $RS$  will be equidistant from  $A, B$ , and  $C$ .

*Obs.* These principles may also be used to prove the theorems relating to concurrency already given on page 111.

**EXAMPLE.** *To prove that the bisectors of the angles of a triangle are concurrent.*

Let  $ABC$  be a triangle.

Bisect the  $\angle^s$   $ABC, BCA$  by straight lines  $BO, CO$  : these must meet at some point  $O$ . Ax. 12.

Join  $OA$ .

Then shall  $OA$  bisect the  $\angle BAC$ .

Now  $BO$  is the locus of points equidistant from  $BC, BA$ ; Ex. 3, p. 55.


$\therefore OP = OR$ .

Similarly  $CO$  is the locus of points equidistant from  $BC, CA$ .

$\therefore OP = OQ$ ; hence  $OR = OQ$ .

$\therefore O$  is on the locus of points equidistant from  $AB$  and  $AC$  : that is,  $OA$  is the bisector of the  $\angle BAC$ .

Hence the bisectors of the three  $\angle^s$  meet at the point  $O$ .



It may happen that the data of the problem are so related to one another that the resulting loci do not intersect. In this case the problem is impossible.

For example, if in Ex. 1, page 126, the length of the given median *is less than* the given altitude, the straight line CD will not be intersected by the circle, and no triangle can fulfil the conditions of the problem. If the length of the median *is equal* to the given altitude, *one* point is common to the two loci; and consequently *only one* solution of the problem exists: and we have seen that there are two solutions, if the median is greater than the altitude.

In examples of this kind the student should make a point of investigating the relations which must exist among the data, in order that the problem may be possible; and he must observe that if under certain relations *two* solutions are possible, and under other relations no solution exists, there will always be some *intermediate* relation under which *one* and *only one* solution is possible.

#### EXAMPLES.

1. Find a point in a given straight line which is equidistant from two given points.

2. Find a point which is at given distances from each of two given straight lines. How many solutions are possible?

3. *On a given base construct a triangle, having given one angle at the base and the length of the opposite side. Examine the relations which must exist among the data in order that there may be two solutions, one solution, or that the problem may be impossible.*

4. On the base of a given triangle construct a second triangle equal in area to the first, and having its vertex in a given straight line.

5. Construct an isosceles triangle equal in area to a given triangle, and standing on the same base.

6. Find a point which is at a given distance from a given point, and is equidistant from two given parallel straight lines.

When does this problem admit of two solutions, when of one only, and when is it impossible?

## BOOK II.

BOOK II. deals with the areas of rectangles and squares.


A **Rectangle** has been defined (Book I., Def. 37) as a parallelogram which has one of its angles a right angle.

It should be remembered that if a parallelogram has *one* right angle, *all* its angles are right angles. [I. 46, Cor.]

### DEFINITIONS.

1. A rectangle is said to be **contained** by any two of its sides which form a right angle: for it is clear that both the form and magnitude of a rectangle are fully determined when the lengths of two such sides are given.

Thus the rectangle ACDB is said to be contained by AB, AC; or by CD, DB: and if X and Y are two straight lines equal respectively to AB and AC, then the rectangle contained by X and Y is equal to the rectangle contained by AB, AC.




[See Ex. 12, p. 70.]

After Proposition 3, we shall use the abbreviation *rect.* AB, AC to denote the *rectangle contained by AB and AC*.

2. In any parallelogram the figure formed by either of the parallelograms about a diagonal together with the two complements is called a **gnomon**.

Thus the shaded portion of the annexed diagram, consisting of the parallelogram EH together with the complements AK, KC is the *gnomon* AHF.

The other gnomon in the diagram is that which is made up of the figures AK, GF and FH, namely the gnomon AFH.




## INTRODUCTORY.

Before entering upon Book II. the student is reminded of the following arithmetical rule :

**RULE.** *To find the area of a rectangle, multiply the number of units in the length by the number of units in the breadth; the product will be the number of square units in the area.*

For example, if the two sides AB, AD of the rectangle ABCD are respectively *four* and *three* inches long, and if through the points of division parallels are drawn as in the annexed figure, it is seen that the rectangle is divided into *three* rows, each containing *four* square inches, or into *four* columns, each containing *three* square inches.



Hence the whole rectangle contains  $3 \times 4$ , or 12, square inches.


Similarly if AB and AD contain  $m$  and  $n$  units of length respectively, it follows that the rectangle ABCD will contain  $m \times n$  units of area: further, if AB and AD are equal, each containing  $m$  units of length, the rectangle becomes a square, and contains  $m^2$  units of area.

From this we conclude that *the rectangle contained by two straight lines* in Geometry corresponds to *the product of two numbers* in Arithmetic or Algebra; and that *the square described on a straight line* corresponds to *the square of a number*. Accordingly it will be found in the course of Book II. that several theorems relating to the areas of rectangles and squares are analogous to well-known algebraical formulæ.

In view of these principles the rectangle contained by two straight lines AB, BC is sometimes expressed in the form of a product, as  $AB \cdot BC$ , and the square described on AB as  $AB^2$ . This notation, together with the signs + and -, will be employed in the additional matter appended to this book; *but it is not admitted into Euclid's text* because it is desirable in the first instance to emphasize the distinction between *geometrical magnitudes themselves* and the *numerical equivalents* by which they may be expressed arithmetically.

## PROPOSITION 1. THEOREM.

If there are two straight lines, one of which is divided into any number of parts, the rectangle contained by the two straight lines is equal to the sum of the rectangles contained by the undivided straight line and the several parts of the divided line.



Let  $P$  and  $AB$  be two straight lines, and let  $AB$  be divided into any number of parts  $AC$ ,  $CD$ ,  $DB$ .

Then shall the rectangle contained by  $P$ ,  $AB$  be equal to the sum of the rectangles contained by  $P$ ,  $AC$ , by  $P$ ,  $CD$ , and by  $P$ ,  $DB$ .

**Construction.** From  $A$  draw  $AF$  perp. to  $AB$ ; I. 11.  
and make  $AG$  equal to  $P$ . I. 3.  
Through  $G$  draw  $GH$  par<sup>l</sup> to  $AB$ ; I. 31.  
and through  $C$ ,  $D$ ,  $B$  draw  $CK$ ,  $DL$ ,  $BH$  par<sup>l</sup> to  $AG$ .

**Proof.** Now the fig.  $AH$  is made up of the figs.  $AK$ ,  $CL$ ,  $DH$ , and is therefore equal to their sum;  
and of these,

the fig.  $AH$  is the rectangle contained by  $P$ ,  $AB$ ;  
for it is contained by  $AG$ ,  $AB$ ; and  $AG = P$ ;  
and the fig.  $AK$  is the rectangle contained by  $P$ ,  $AC$ ;  
for it is contained by  $AG$ ,  $AC$ ; and  $AG = P$ ;  
also the fig.  $CL$  is the rectangle contained by  $P$ ,  $CD$ ;  
for it is contained by  $CK$ ,  $CD$ ;  
and  $CK =$  the opp. side  $AG$ , and  $AG = P$ . I. 34.

Similarly the fig.  $DH$  is the rectangle contained by  $P$ ,  $DB$ .

$\therefore$  the rectangle contained by  $P$ ,  $AB$  is equal to the sum of the rectangles contained by  $P$ ,  $AC$ , by  $P$ ,  $CD$ , and by  $P$ ,  $DB$ . Q.E.D.

## CORRESPONDING ALGEBRAICAL FORMULA.

In accordance with the principles explained on page 129, the result of this proposition may be written thus :

$$P \cdot AB = P \cdot AC + P \cdot CD + P \cdot DB.$$

Now if the line  $P$  contains  $p$  units of length, and if  $AC$ ,  $CD$ ,  $DB$  contain  $a$ ,  $b$ ,  $c$  units respectively,

$$\text{then } AB = a + b + c;$$

hence the statement

$$P \cdot AB = P \cdot AC + P \cdot CD + P \cdot DB$$

becomes


$$p(a + b + c) = pa + pb + pc.$$

[NOTE. It must be understood that the rule given on page 129, for expressing the area of a rectangle as the product of the lengths of two adjacent sides, implies that those sides are **commensurable**, that is, that they can be expressed *exactly* in terms of some common unit.

This however is not always the case. Two straight lines may be so related that it is impossible to divide either of them into equal parts, *of which the other contains an exact number*. Such lines are said to be **incommensurable**. Hence if the adjacent sides of a rectangle are incommensurable, we cannot choose any linear unit in terms of which these sides may be *exactly* expressed; and thus it will be impossible to subdivide the rectangle into squares of unit area, as illustrated in the figure of page 129. We do not here propose to enter further into the subject of incommensurable quantities: it is sufficient to point out that further knowledge of them will convince the student that the area of a rectangle may be expressed *to any required degree of accuracy* by the product of the lengths of two adjacent sides, whether those lengths are commensurable or not.]

## PROPOSITION 2. THEOREM.

If a straight line is divided into any two parts, the square on the whole line is equal to the sum of the rectangles contained by the whole line and each of the parts.



Let the straight line AB be divided at C into the two parts AC, CB.

Then shall the square on AB be equal to the sum of the rectangles contained by AB, AC, and by AB, BC.

**Construction.** On AB describe the square ADEB. I. 46.  
Through C draw CF par<sup>l</sup> to AD. I. 31.

**Proof.** Now the fig. AE is made up of the figs. AF, CE :  
and of these,

the fig. AE is the sq. on AB : *Constr.*

and the fig. AF is the rectangle contained by AB, AC ;  
for it is contained by AD, AC ; and AD = AB :

also the fig. CE is the rectangle contained by AB, BC ;  
for it is contained by BE, BC ; and BE = AB.

∴ the sq. on AB = the sum of the rectangles contained  
by AB, AC, and by AB, BC. Q.E.D.

## CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AB^2 = AB \cdot AC + AB \cdot BC.$$

Let AC contain  $a$  units of length, and let CB contain  $b$  units,


then  $AB = a + b$  units ;

and we have

$$(a + b)^2 = (a + b)a + (a + b)b.$$

PROPOSITION 3. THEOREM.

*If a straight line is divided into any two parts, the rectangle contained by the whole and one of the parts is equal to the square on that part together with the rectangle contained by the two parts.*



Let the straight line AB be divided at C into the two parts AC, CB.

*Then shall the rectangle contained by AB, AC be equal to the square on AC together with the rectangle contained by AC, CB.*

**Construction.** On AC describe the square AFDC. I. 46.  
Through B draw BE par<sup>l</sup> to AF, meeting FD produced in E.  
I. 31.

**Proof.** Now the fig. AE is made up of the figs. AD, CE ;  
and of these,

the fig. AE is the rectangle contained by AB, AC ;  
for AF = AC ;

and the fig. AD is the sq. on AC ; *Constr.*

also the fig. CE is the rectangle contained by AC, CB ;  
for CD = AC.

∴ the rectangle contained by AB, AC is equal to the sq. on AC together with the rectangle contained by AC, CB.

Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

This result may be written  $AB \cdot AC = AC^2 + AC \cdot CB$ .

Let AC, CB contain  $a$  and  $b$  units of length respectively,  
then  $AB = a + b$  units ;


and we have  $(a + b)a = a^2 + ab$ .

**NOTE.** It should be observed that Props. 2 and 3 are *special cases* of Prop. 1.



## PROPOSITION 4. THEOREM.

*If a straight line is divided into any two parts, the square on the whole line is equal to the sum of the squares on the two parts together with twice the rectangle contained by the two parts.*



Let the straight line  $AB$  be divided at  $C$  into the two parts  $AC$ ,  $CB$ .

Then shall the sq. on  $AB$  be equal to the sum of the sqq. on  $AC$ ,  $CB$ , together with twice the rect.  $AC$ ,  $CB$ .

**Construction.** On  $AB$  describe the square  $ADEB$ . I. 46.  
Join  $BD$ .

Through  $C$  draw  $CF$  par<sup>l</sup> to  $BE$ , meeting  $BD$  in  $G$ . I. 31.  
Through  $G$  draw  $HGK$  par<sup>l</sup> to  $AB$ .

It is first required to shew that the fig.  $CK$  is the sq. on  $CB$ .

**Proof.** Because  $CF$  and  $AD$  are par<sup>l</sup>, and  $BD$  meets them,  
 $\therefore$  the ext. angle  $CGB =$  the int. opp. angle  $ADB$ . I. 29.

And since  $AB = AD$ , being sides of a square ;

$\therefore$  the angle  $ADB =$  the angle  $ABD$  ; I. 5.

$\therefore$  the angle  $CGB =$  the angle  $CBG$ .

$\therefore CB = CG$ . I. 6.

And the opp. sides of the par<sup>m</sup>  $CK$  are equal ; I. 34.

$\therefore$  the fig.  $CK$  is equilateral ;

also the angle  $CBK$  is a right angle ; Def. 30.

$\therefore CK$  is a square, and it is described on  $CB$ . I. 46, Cor.

Similarly, the fig.  $HF$  is the sq. on  $HG$ , that is, the sq. on  $AC$  ;

for  $HG =$  the opp. side  $AC$ . I. 34.

Again, the complement  $AG =$  the complement  $GE$ ; I. 43.  
 and the fig.  $AG =$  the rect.  $AC, CB$ ; for  $CG = CB$ .  
 $\therefore$  the two figs.  $AG, GE =$  twice the rect.  $AC, CB$ .

\* Now the sq. on  $AB =$  the fig.  $AE$   
 $=$  the figs.  $HF, CK, AG, GE$   
 $=$  the sqq. on  $AC, CB$  together with  
 twice the rect.  $AC, CB$ .  
 $\therefore$  the sq. on  $AB =$  the sum of the sqq. on  $AC, CB$  with  
 twice the rect.  $AC, CB$ . Q.E.D.

**COROLLARY 1.** *Parallelograms about the diagonals of a square are themselves squares.*

**COROLLARY 2.** *If a straight line is bisected, the square on the whole line is four times the square on half the line.*

\* For the purpose of oral work, this step of the proof may conveniently be arranged as follows :

Now the sq. on  $AB$  is equal to the fig.  $AE$ ,  
 that is, to the figs.  $HF, CK, AG, GE$  ;  
 that is, to the sqq. on  $AC, CB$  together  
 with twice the rect.  $AC, CB$ .

#### CORRESPONDING ALGEBRAICAL FORMULA.

The result of this important Proposition may be written thus :

$$AB^2 = AC^2 + CB^2 + 2AC \cdot CB.$$


Let  $AC = a$ , and  $CB = b$  ;  
 then  $AB = a + b$  ;

hence the statement  $AB^2 = AC^2 + CB^2 + 2AC \cdot CB$

becomes  $(a + b)^2 = a^2 + b^2 + 2ab$ .

## PROPOSITION 5. THEOREM.

*If a straight line is divided equally and also unequally, the rectangle contained by the unequal parts, together with the square on the line between the points of section, is equal to the square on half the line.*



Let the straight line  $AB$  be divided equally at  $P$ , and unequally at  $Q$ .

*Then the rect.  $AQ, QB$  together with the sq. on  $PQ$  shall be equal to the sq. on  $PB$ .*

**Construction.** On  $PB$  describe the square  $PCDB$ . I. 46.  
Join  $BC$ .

Through  $Q$  draw  $QE$  par<sup>l</sup> to  $BD$ , cutting  $BC$  in  $F$ . I. 31.

Through  $F$  draw  $LFHG$  par<sup>l</sup> to  $AB$ .

Through  $A$  draw  $AG$  par<sup>l</sup> to  $BD$ .

**Proof.**

Now the complement  $PF =$  the complement  $FD$ : I. 43.  
to each add the fig.  $QL$ ;

then the fig.  $PL =$  the fig.  $QD$ .

But the fig.  $PL =$  the fig.  $AH$ , for they are par<sup>ms</sup> on equal bases and between the same par<sup>ls</sup>; I. 36.

$\therefore$  the fig.  $AH =$  the fig.  $QD$ .

To each add the fig.  $PF$ ;

then the fig.  $AF =$  the gnomon  $PLE$ .

Now the fig.  $AF$  is the rect.  $AQ, QB$ ; for  $QF = QB$ ;

$\therefore$  the rect.  $AQ, QB =$  the gnomon  $PLE$ .

To each add the sq. on  $PQ$ , that is, the fig.  $HE$ ; II. 4.  
then the rect.  $AQ, QB$  with the sq. on  $PQ$

$=$  the gnomon  $PLE$  with the fig.  $HE$

$=$  the whole fig.  $PD$ ,


which is the sq. on  $PB$ .

That is, the rect.  $AQ, QB$  together with the square on  $PQ$  is equal to the sq. on  $PB$ . Q.E.D.

**COROLLARY.** From this Proposition it follows that *the difference of the squares on two straight lines is equal to the rectangle contained by their sum and difference.*

For let  $X$  and  $Y$  be the given st. lines, of which  $X$  is the greater.

Draw  $AP$  equal to  $X$ , and produce it to  $B$ , making  $PB$  equal to  $AP$ , that is to  $X$ .



From  $PB$  cut off  $PQ$  equal to  $Y$ .

Then  $AQ$  is equal to the sum of  $X$  and  $Y$ , and  $QB$  is equal to the difference of  $X$  and  $Y$ .

Now because  $AB$  is divided equally at  $P$  and unequally at  $Q$ ,  
 $\therefore$  the rect.  $AQ, QB$  with sq. on  $PQ$  = the sq. on  $PB$ ; II. 5.  
 that is, the difference of the sqq. on  $PB, PQ$  = the rect.  $AQ, QB$ .  
 or, the difference of the sqq. on  $X$  and  $Y$  = the rectangle contained by the sum and the difference of  $X$  and  $Y$ .

**CORRESPONDING ALGEBRAICAL FORMULA.**

This result may be written

$$AQ \cdot QB + PQ^2 = PB^2.$$

Let  $AB = 2a$ ; and let  $PQ = b$ ;

then  $AP$  and  $PB$  each =  $a$ .

Also  $AQ = a + b$ ; and  $QB = a - b$ .

Hence the statement  $AQ \cdot QB + PQ^2 = PB^2$

becomes  $(a + b)(a - b) + b^2 = a^2$ ,


or  $(a + b)(a - b) = a^2 - b^2$ .

**EXERCISE.**

*In the above figure shew that  $AP$  is half the sum of  $AQ$  and  $QB$ ; and that  $PQ$  is half their difference.*

## PROPOSITION 6. THEOREM.

If a straight line is bisected and produced to any point, the rectangle contained by the whole line thus produced and the part of it produced, together with the square on half the line bisected, is equal to the square on the straight line made up of the half and the part produced.



Let the straight line  $AB$  be bisected at  $P$ , and produced to  $Q$ .

Then the rect.  $AQ, QB$  together with the sq. on  $PB$  shall be equal to the sq. on  $PQ$ .

Construction. On  $PQ$  describe the square  $PCDQ$ . I. 46.  
Join  $QC$ .

Through  $B$  draw  $BE$  par<sup>l</sup> to  $QD$ , meeting  $QC$  in  $F$ . I. 31.

Through  $F$  draw  $LFHG$  par<sup>l</sup> to  $AQ$ .

Through  $A$  draw  $AG$  par<sup>l</sup> to  $QD$ .

Proof. Now the complement  $PF =$  the complement  $FD$ . I. 43.

But the fig.  $PF =$  the fig.  $AH$ ; for they are par<sup>ms</sup> on equal bases and between the same par<sup>ls</sup>. I. 36.

$\therefore$  the fig.  $AH =$  the fig.  $FD$ .

To each add the fig.  $PL$ ;

then the fig.  $AL =$  the gnomon  $PLE$ .

Now the fig.  $AL$  is the rect.  $AQ, QB$ ; for  $QL = QB$ ;

$\therefore$  the rect.  $AQ, QB =$  the gnomon  $PLE$ .

To each add the sq. on  $PB$ , that is, the fig.  $HE$ ;

then the rect.  $AQ, QB$  with the sq. on  $PB$   
 $=$  the gnomon  $PLE$  with the fig.  $HE$   
 $=$  the whole fig.  $PD$ ,  
 which is the square on  $PQ$ .

That is, the rect.  $AQ, QB$  together with the sq. on  $PB$  is equal to the sq. on  $PQ$ . Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

This result may be written

$$AQ \cdot QB + PB^2 = PQ^2.$$

Let  $AB = 2a$ ; and let  $PQ = b$ ;

then  $AP$  and  $PB$  each  $= a$ .

Also  $AQ = a + b$ ; and  $QB = b - a$ .

Hence the statement  $AQ \cdot QB + PB^2 = PQ^2$   
 becomes  $(a + b)(b - a) + a^2 = b^2$ ,  
 or  $(b + a)(b - a) = b^2 - a^2$ .

**DEFINITION.** If a point  $X$  is taken in a straight line  $AB$ , or in  $AB$  produced, the distances of the point of section from the extremities of  $AB$  are said to be the **segments** into which  $AB$  is divided at  $X$ .



In the former case  $AB$  is divided **internally**, in the latter case **externally**.

Thus in *each* of the annexed figures, the segments into which  $AB$  is divided at  $X$  are the lines  $AX$  and  $XB$ .

This definition enables us to include Props. 5 and 6 in a single Enunciation.

*If a straight line is bisected, and also divided (internally or externally) into two unequal segments, the rectangle contained by the unequal segments is equal to the difference of the squares on half the line, and on the line between the points of section.*

EXERCISE.


Shew that the Enunciations of Props. 5 and 6 may take the following form :

*The rectangle contained by two straight lines is equal to the difference of the squares on half their sum and on half their difference.*

[See Ex., p. 137.]

## PROPOSITION 7. THEOREM.

*If a straight line is divided into any two parts, the sum of the squares on the whole line and on one of the parts is equal to twice the rectangle contained by the whole and that part, together with the square on the other part.*



Let the straight line  $AB$  be divided at  $C$  into the two parts  $AC$ ,  $CB$ .

*Then shall the sum of the sqq. on  $AB$ ,  $BC$  be equal to twice the rect.  $AB$ ,  $BC$  together with the sq. on  $AC$ .*

**Construction.** On  $AB$  describe the square  $ADEB$ . I. 46.  
Join  $BD$ .

Through  $C$  draw  $CF$  par<sup>l</sup> to  $BE$ , meeting  $BD$  in  $G$ . I. 31.  
Through  $G$  draw  $HGK$  par<sup>l</sup> to  $AB$ .

**Proof.** Now the complement  $AG =$  the complement  $GE$ ; I. 43.  
to each add the fig.  $CK$  :

then the fig.  $AK =$  the fig.  $CE$ .

But the fig.  $AK$  is the rect.  $AB$ ,  $BC$ ; for  $BK = BC$ ;

$\therefore$  the two figs  $AK$ ,  $CE =$  twice the rect.  $AB$ ,  $BC$ .

But the two figs.  $AK$ ,  $CE$  make up the gnomon  $AKF$  and the fig.  $CK$  :

$\therefore$  the gnomon  $AKF$  with the fig.  $CK =$  twice the rect.  $AB$ ,  $BC$ .

To each add the fig.  $HF$ , which is the sq. on  $AC$  :

then the gnomon  $AKF$  with the figs.  $CK$ ,  $HF$

$=$  twice the rect.  $AB$ ,  $BC$  with the sq. on  $AC$ .

But the gnomon  $AKF$  with the figs.  $CK$ ,  $HF$  make up the figs.  $AE$ ,  $CK$ , that is to say, the sqq. on  $AB$ ,  $BC$ ;

$\therefore$  the sqq. on  $AB$ ,  $BC =$  twice the rect.  $AB$ ,  $BC$  with the sq. on  $AC$ . Q.E.D.

## CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AB^2 + BC^2 = 2AB \cdot BC + AC^2.$$

Let  $AB = a$ , and  $BC = b$ ; then  $AC = a - b$ .

Hence the statement

$$AB^2 + BC^2 = 2AB \cdot BC + AC^2$$

becomes

$$a^2 + b^2 = 2ab + (a - b)^2,$$

or

$$(a - b)^2 = a^2 - 2ab + b^2.$$

Comparing this result with that obtained from Prop. 4, we see that

(i) *The square on the sum of two straight lines is greater than the sum of the squares on those lines by twice the rectangle contained by them.* [Prop. 4.]

(ii) *The square on the difference of two straight lines is less than the sum of the squares on those lines by twice the rectangle contained by them.* [Prop. 7.]

## ALTERNATIVE PROOFS OF PROPOSITIONS 4, 5, 6, 7.

The following alternative proofs are recommended for purposes of revision, as affording useful exercise on the enunciations of preceding propositions, and illustrating the way in which many examples on Book II. may be solved. The beginner however should not adopt these proofs until he has thoroughly mastered those given in the text, where the rectangles and squares are actually represented in the diagrams.

## PROPOSITION 4.

Let the straight line  $AB$  be divided at  $C$  into two parts  $AC$ ,  $CB$ .

*Then shall the sq. on  $AB$  be equal to the sum of the sqq. on  $AC$ ,  $CB$  with twice the rect.  $AC$ ,  $CB$ .*



Now the sq. on  $AB =$  the rect.  $AB$ ,  $AC$  with the rect.  $AB$ ,  $CB$ . II. 2.  
But the rect.  $AB$ ,  $AC =$  the sq. on  $AC$  with the rect.  $AC$ ,  $CB$ ; II. 3.  
and the rect.  $AB$ ,  $CB =$  the sq. on  $CB$  with the rect.  $AC$ ,  $CB$ . II. 3.

Hence the sq. on  $AB =$  the sum of the sqq. on  $AC$ ,  $CB$  with twice the rect.  $AC$ ,  $CB$ .



## PROPOSITION 5.

Let the straight line  $AB$  be divided equally at  $P$ , and unequally at  $Q$ .

Then shall the rect.  $AQ, QB$  with the sq. on  $PQ$  be equal to the sq. on  $PB$ .



Now the rect.  $AQ, QB$  = the rect.  $AP, QB$  with the rect.  $PQ, QB$ , II. 1.  
= the rect.  $PB, QB$  with the rect.  $PQ, QB$ .

But the rect.  $PB, QB$  = the sq. on  $QB$  with the rect.  $PQ, QB$ ; II. 3.  
∴ the rect.  $AQ, QB$  = the sq. on  $QB$  with twice the rect.  $PQ, QB$ .

To each of these equals add the sq. on  $PQ$ .

Then the rect.  $AQ, QB$  with the sq. on  $PQ$   
= the sqq. on  $PQ, QB$  with twice the rect.  $PQ, QB$   
= the sq. on  $PB$ . II. 4.

## PROPOSITION 6.

Let the straight line  $AB$  be bisected at  $P$ , and produced to  $Q$ .

Then shall the rect.  $AQ, QB$  with the sq. on  $PB$  be equal to the sq. on  $PQ$ .



Now the rect.  $AQ, QB$  = the rect.  $AP, BQ$  with the rect.  $PQ, BQ$ , II. 1.  
= the rect.  $PB, BQ$  with the rect.  $PQ, BQ$ .

But the rect.  $PQ, BQ$  = the sq. on  $BQ$  with the rect.  $PB, BQ$ , II. 3.  
∴ the rect.  $AQ, QB$  = the sq. on  $BQ$  with twice the rect.  $PB, BQ$ .

To each of these equals add the sq. on  $PB$ .

Then the rect.  $AQ, QB$  with the sq. on  $PB$   
= the sqq. on  $PB, BQ$  with twice the rect.  $PB, BQ$   
= the sq. on  $PQ$ . II. 4.

## PROPOSITION 7.

Let the straight line  $AB$  be divided at any point  $C$ .

Then shall the sum of the sqq. on  $AB, BC$  be equal to twice the rect.  $AB, BC$  with the sq. on  $AC$ .



Now the sq. on  $AB$  = the sqq. on  $AC, CB$  with twice the rect.  $AC, CB$ , II. 4.

To each of these equals add the sq. on  $BC$ .

Then the sqq. on  $AB, BC$  = the sq. on  $AC$  with twice the sq. on  $BC$   
and twice the rect.  $AC, CB$ .

But twice the sq. on  $BC$  with twice the rect.  $AC, CB$   
= twice the rect.  $AB, BC$ . II. 3.  
∴ the sqq. on  $AB, BC$  = the sq. on  $AC$  with twice the rect.  $AB, BC$ .

*Obs.* The following proposition being little used, we merely give the figure and the leading points of Euclid's proof.

PROPOSITION 8. THEOREM.

*If a straight line is divided into any two parts, four times the rectangle contained by the whole line and one of the parts, together with the square on the other part, is equal to the square on the straight line which is made up of the whole and the first named part.*

Let AB be divided at C.  
Produce AB to D, making BD equal to BC.

Then shall four times the rect. AB, BC with the sq. on AC = the sq. on AD.

On AD describe the square AEFD; and complete the construction as indicated in the figure.

Euclid then proves (i) that the figs. CK, BN, GR, KO are all equal:


(ii) that the figs. AG, MP, PL, RF are all equal.

Hence the eight figures named above are together four times the sum of the figs. AG, CK; that is, four times the fig. AK; that is, four times the rect. AB, BC.


But the whole fig. AF, namely the sq. on AD, is made up of these eight figures, together with the fig. XH, which is the sq. on AC:

hence the sq. on AD = four times the rect. AB, BC, together with the sq. on AC.

Q. E. D.



The accompanying figure will suggest a less cumbrous proof, which we leave as an Exercise to the student.



CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$4AB \cdot BC + AC^2 = AD^2.$$


Let  $AB = a$ , and  $BC = b$ ; then  $AC = a - b$ , and  $AD = a + b$ .

Hence we have  $4ab + (a - b)^2 = (a + b)^2$ ;

or  $(a + b)^2 - (a - b)^2 = 4ab.$

## PROPOSITION 9. THEOREM. [EUCLID'S PROOF.]

If a straight line is divided equally and also unequally, the sum of the squares on the two unequal parts is twice the sum of the squares on half the line and on the line between the points of section.



Let the straight line AB be divided equally at P, and unequally at Q.

Then shall the sum of the sqq. on AQ, QB be twice the sum of the sqq. on AP, PQ.

**Construction.** At P draw PC at rt. angles to AB; I. 11.  
and make PC equal to AP or PB. I. 3.  
Join AC, BC.

Through Q draw QD par<sup>l</sup> to PC; I. 31.  
and through D draw DE par<sup>l</sup> to AB.  
Join AD.

**Proof.** Then since PA = PC, *Constr.*  
 $\therefore$  the angle PAC = the angle PCA. I. 5.

And since, in the triangle APC, the angle APC is a rt. angle, *Constr.*

$\therefore$  the sum of the angles PAC, PCA is a rt. angle: I. 32.  
hence each of the angles PAC, PCA is half a rt. angle.

So also, each of the angles PBC, PCB is half a rt. angle.  
 $\therefore$  the whole angle ACB is a rt. angle.

Again, the ext. angle CED = the int. opp. angle CPB; I. 29.  
 $\therefore$  the angle CED is a rt. angle:

and the angle ECD is half a rt. angle. *Proved.*

$\therefore$  the remaining angle EDC is half a rt. angle; I. 32.

$\therefore$  the angle ECD = the angle EDC;  
 $\therefore$  EC = ED. I. 6.

Again, the ext. angle  $DQB =$  the int. opp. angle  $CPB$  ; I. 29.

$\therefore$  the angle  $DQB$  is a rt. angle.

And the angle  $QBD$  is half a rt. angle ; *Proved.*

$\therefore$  the remaining angle  $QDB$  is half a rt. angle ; I. 32.

$\therefore$  the angle  $QBD =$  the angle  $QDB$  ;

$\therefore QD = QB.$

Now the sq. on  $AP =$  the sq. on  $PC$  ; for  $AP = PC.$  *Constr.*

And since the angle  $APC$  is a rt. angle,

$\therefore$  the sq. on  $AC =$  the sum of the sqq. on  $AP, PC$  ; I. 47.

$\therefore$  the sq. on  $AC$  is twice the sq. on  $AP.$

Similarly, the sq. on  $CD$  is twice the sq. on  $ED$ , that is, twice the sq. on the opp. side  $PQ.$  I. 34.

Now the sqq. on  $AQ, QB =$  the sqq. on  $AQ, QD$  *Proved.*

$=$  the sq. on  $AD$ , for  $AQD$  is a rt. angle ; I. 47.

$=$  the sum of the sqq. on  $AC, CD$ , for  $ACD$  is a rt. angle ; I. 47.

$=$  twice the sq. on  $AP$  with twice the sq. on  $PQ.$  *Proved.*

That is,

the sum of the sqq. on  $AQ, QB =$  twice the sum of the sqq. on  $AP, PQ.$  Q.E.D.

CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AQ^2 + QB^2 = 2(AP^2 + PQ^2).$$

Let  $AB = 2a$  ; and  $PQ = b$  ;

then  $AP$  and  $PB$  each  $= a.$

Also  $AQ = a + b$  ; and  $QB = a - b.$

Hence the statement

$$AQ^2 + QB^2 = 2(AP^2 + PQ^2)$$


becomes

$$(a + b)^2 + (a - b)^2 = 2(a^2 + b^2).$$

[NOTE. For alternative proofs of this proposition, see page 148.]

## PROPOSITION 10. THEOREM. [EUCLID'S PROOF.]

If a straight line is bisected and produced to any point, the sum of the squares on the whole line thus produced, and on the part produced, is twice the sum of the squares on half the line bisected and on the line made up of the half and the part produced.



Let the st. line  $AB$  be bisected at  $P$ , and produced to  $Q$ .  
Then shall the sum of the sqq. on  $AQ$ ,  $QB$  be twice the sum of the sqq. on  $AP$ ,  $PQ$ .

**Construction.** At  $P$  draw  $PC$  at right angles to  $AB$ ; I. 11.  
and make  $PC$  equal to  $PA$  or  $PB$ . I. 3.  
Join  $AC$ ,  $BC$ .

Through  $Q$  draw  $QD$  par<sup>l</sup> to  $PC$ , to meet  $CB$  produced in  $D$ ; I. 31.  
and through  $D$  draw  $DE$  par<sup>l</sup> to  $AB$ , to meet  $CP$  produced in  $E$ .

Join  $AD$ .

**Proof.** Then since  $PA = PC$ , *Constr.*  
 $\therefore$  the angle  $PAC =$  the angle  $PCA$ . I. 5.  
And since, in the triangle  $APC$ , the angle  $APC$  is a rt. angle,  
 $\therefore$  the sum of the angles  $PAC$ ,  $PCA$  is a rt. angle. I. 32.  
Hence each of the angles  $PAC$ ,  $PCA$  is half a rt. angle.  
So also, each of the angles  $PBC$ ,  $PCB$  is half a rt. angle.  
 $\therefore$  the whole angle  $ACB$  is a rt. angle.  
Again, the ext. angle  $CPB =$  the int. opp. angle  $CED$ : I. 29.  
 $\therefore$  the angle  $CED$  is a rt. angle:  
and the angle  $ECD$  is half a rt. angle; *Proved.*  
 $\therefore$  the remaining angle  $EDC$  is half a rt. angle. I. 32.  
 $\therefore$  the angle  $ECD =$  the angle  $EDC$ ;  
 $\therefore EC = ED$ . I. 6.

Again, the angle  $DQB =$  the alt. angle  $CPB$ ; I. 29.

$\therefore$  the angle  $DQB$  is a rt. angle.

Also the angle  $QBD =$  the vert. opp. angle  $CBP$ : I. 15.

that is, the angle  $QBD$  is half a rt. angle.

$\therefore$  the remaining angle  $QDB$  is half a rt. angle: I. 32.

$\therefore$  the angle  $QBD =$  the angle  $QDB$ ;

$\therefore QB = QD$ .

I. 6.

Now the sq. on  $AP =$  the sq. on  $PC$ ; for  $AP = PC$ . *Constr.*

And since the angle  $APC$  is a rt. angle,

$\therefore$  the sq. on  $AC =$  the sum of the sqq. on  $AP, PC$ ; I. 47.

$\therefore$  the sq. on  $AC$  is twice the sq. on  $AP$ .

Similarly, the sq. on  $CD$  is twice the sq. on  $ED$ , that is, twice

the sq. on the opp. side  $PQ$ .

I. 34.

Now the sqq. on  $AQ, QB =$  the sqq. on  $AQ, QD$  *Proved.*

$=$  the sq. on  $AD$ , for  $AQD$  is a rt.

angle;

I. 47.

$=$  the sum of the sqq. on  $AC, CD$ ,

for  $ACD$  is a rt. angle; I. 47.

$=$  twice the sq. on  $AP$  with twice

the sq. on  $PQ$ .

*Proved.*

That is,

the sum of the sqq. on  $AQ, QB$  is twice the sum of the sqq.

on  $AP, PQ$ .

Q.E.D.

#### CORRESPONDING ALGEBRAICAL FORMULA.

The result of this proposition may be written

$$AQ^2 + QB^2 = 2(AP^2 + PQ^2).$$

Let  $AB = 2a$ ; and  $PQ = b$ ;

then  $AP$  and  $PB$  each  $= a$ .

Also  $AQ = a + b$ ; and  $QB = b - a$ .

Hence we have

$$(a + b)^2 + (b - a)^2 = 2(a^2 + b^2).$$

[NOTE. For alternative proofs of this proposition, see page 149.]

## PROPOSITION 9. [ALTERNATIVE PROOF.]

If a straight line is divided equally and also unequally, the sum of the squares on the two unequal parts is twice the sum of the squares on half the line and on the line between the points of section.



Let the straight line AB be divided equally at P and unequally at Q.

Then shall the sum of the sqq. on AQ, QB be twice the sum of the sqq. on AP, PQ.

*Proof.*

The sq. on AQ = the sum of the sqq. on AP, PQ with twice the rect. AP, PQ II. 4.  
 = the sum of the sqq. on AP, PQ with twice the rect. PB, PQ; for PB = AP.

To each of these equals add the sq. on QB.

Then the sqq. on AQ, QB = the sum of the sqq. on AP, PQ with twice the rect. PB, PQ and the sq. on QB.

But twice the rect. PB, PQ and the sq. on QB = the sum of the sqq. on PB, PQ. II. 7.

∴ the sqq. on AQ, QB = the sum of the sqq. on AP, PQ with the sum of the sqq. on PB, PQ = twice the sum of the sqq. on AP, PQ. Q.E.D.

NOTE. The following concise proof, obtained from II. 4 and II. 5, is useful as an exercise, but it is hardly admissible as a formal demonstration owing to its algebraical use of the negative sign.

We have  $AQ^2 + QB^2 = AB^2 - 2AQ \cdot QB$  II. 4.  
 $= 4PB^2 - 2AQ \cdot QB$  II. 4, Cor. 2.  
 $= 4PB^2 - 2(PB^2 - PQ^2)$  II. 5.  
 $= 2PB^2 + 2PQ^2.$

PROPOSITION 10. [ALTERNATIVE PROOF.]

*If a straight line is bisected and produced to any point, the sum of the squares on the whole line thus produced and on the part produced, is twice the sum of the squares on half the line bisected and on the line made up of the half and the part produced.*



Let the straight line AB be bisected at P, and produced to Q.

*Then shall the sum of the sqq. on AQ, QB be twice the sum of the sqq. on AP, PQ.*

*Proof.*

The sq. on AQ = the sum of the sqq. on AP, PQ with twice the rect. AP, PQ II. 4.  
 = the sum of the sqq. on AP, PQ with twice the rect. PB, PQ; for PB = AP.

To each of these equals add the sq. on QB.

Then the sqq. on AQ, QB = the sum of the sqq. on AP, PQ with twice the rect. PB, PQ and the sq. on QB.

But twice the rect. PB, PQ and the sq. on QB = the sum of the sqq. on PB, PQ. II. 7.

∴ the sqq. on AQ, QB = the sum of the sqq. on AP, PQ with the sum of the sqq. on PB, PQ = twice the sum of the sqq. on AP, PQ. Q.E.D.


**NOTE.** Another proof of this proposition, based on II. 7 and II. 6, is indicated by the following steps:

$$\begin{aligned}
 \text{We have} \quad & \text{AQ}^2 + \text{QB}^2 = 2\text{AQ} \cdot \text{QB} + \text{AB}^2 && \text{II. 7.} \\
 & = 2\text{AQ} \cdot \text{QB} + 4\text{PB}^2 && \text{II. 4, Cor. 2.} \\
 & = 2(\text{PQ}^2 - \text{PB}^2) + 4\text{PB}^2 && \text{II. 6.} \\
 & = 2\text{PB}^2 + 2\text{PQ}^2.
 \end{aligned}$$



## PROPOSITION 11. PROBLEM.

To divide a given straight line into two parts, so that the rectangle contained by the whole and one part may be equal to the square on the other part.



Let  $AB$  be the given straight line.

It is required to divide  $AB$  into two parts, so that the rectangle contained by the whole and one part may be equal to the square on the other part.

**Construction.** On  $AB$  describe the square  $ACDB$ . I. 46.

Bisect  $AC$  at  $E$ . I. 10.

Join  $EB$ .

Produce  $CA$  to  $F$ , making  $EF$  equal to  $EB$ . I. 3.

On  $AF$  describe the square  $AFGH$ . I. 46.

Then shall  $AB$  be divided at  $H$ , so that the rect.  $AB, BH$  is equal to the sq. on  $AH$ .

Produce  $GH$  to meet  $CD$  in  $K$ .

**Proof.** Because  $CA$  is bisected at  $E$ , and produced to  $F$ ,  
 $\therefore$  the rect.  $CF, FA$  with the sq. on  $EA$  = the sq. on  $EF$  II. 6.  
 = the sq. on  $EB$ . *Constr.*

But the sq. on  $EB$  = the sum of the sqq. on  $EA, AB$ ,  
 for the angle  $EAB$  is a rt. angle. I. 47.

$\therefore$  the rect.  $CF, FA$  with the sq. on  $EA$  = the sum of the sqq. on  $EA, AB$ .

From these equals take the sq. on  $EA$  :  
 then the rect.  $CF, FA$  = the sq. on  $AB$ .

But the rect. CF, FA = the fig. FK ; for FA = FG ;  
and the sq. on AB = the fig. AD. *Constr.*

∴ the fig. FK = the fig. AD.

From these equals take the common fig. AK ;  
then the remaining fig. FH = the remaining fig. HD.


But the fig. HD = the rect. AB, BH ; for BD = AB ;  
and the fig. FH is the sq. on AH.

∴ the rect. AB, BH = the sq. on AH. Q.E.F.

**DEFINITION.** A straight line is said to be divided in **Medial Section** when the rectangle contained by the given line and one of its segments is equal to the square on the other segment.

The student should observe that this division may be *internal* or *external*.

Thus if the straight line AB is divided internally at H, and externally at H', so that

(i)  $AB \cdot BH = AH^2$ ,  $H'$   A H B  
(ii)  $AB \cdot BH' = AH^2$ ,

we shall in either case consider that AB is divided in medial section.

The case of *internal* section is alone given in Euclid II. 11 ; but a straight line may be divided *externally* in medial section by a similar process. See Ex. 21, p. 160.

#### ALGEBRAICAL ILLUSTRATION.

It is required to find a point H in AB, or AB produced, such that  
 $AB \cdot BH = AH^2$ .

Let AB contain  $a$  units of length, and let AH contain  $x$  units ;  
then  $BH = a - x$  :

and  $x$  must be such that  $a(a - x) = x^2$ ,  
or  $x^2 + ax - a^2 = 0$ .

Thus the construction for dividing a straight line in medial section corresponds to the solution of this quadratic equation, the two roots of which indicate the *internal* and *external* points of division.


#### EXERCISES.

In the figure of II. 11, shew that

- (i) if CH is produced to meet BF at L, CL is at right angles to BF ;
- (ii) if BE and CH meet at O, AO is at right angles to CH.
- (iii) the lines BG, DF, AK are parallel :
- (iv) CF is divided in medial section at A.

## PROPOSITION 12. THEOREM.

*In an obtuse-angled triangle, if a perpendicular is drawn from either of the acute angles to the opposite side produced, the square on the side subtending the obtuse angle is greater than the sum of the squares on the sides containing the obtuse angle, by twice the rectangle contained by the side on which, when produced, the perpendicular falls, and the line intercepted without the triangle, between the perpendicular and the obtuse angle.*



Let  $ABC$  be an obtuse-angled triangle, having the obtuse angle at  $C$ ; and let  $AD$  be drawn from  $A$  perp. to the opp. side  $BC$  produced.

*Then shall the sq. on  $AB$  be greater than the sum of the sqq. on  $BC, CA$ , by twice the rect.  $BC, CD$ .*

**Proof.** Because  $BD$  is divided into two parts at  $C$ ,  
 $\therefore$  the sq. on  $BD$  = the sum of the sqq. on  $BC, CD$ , with  
 twice the rect.  $BC, CD$ . II. 4,

To each of these equals add the sq. on  $DA$ .  
 Then the sqq. on  $BD, DA$  = the sum of the sqq. on  $BC, CD,$   
 $DA$ , with twice the rect.  $BC, CD$ .


But the sum of the sqq. on  $BD, DA$  = the sq. on  $AB$ ,  
 for the angle at  $D$  is a rt. angle. I. 47.  
 Similarly, the sum of the sqq. on  $CD, DA$  = the sq. on  $CA$ .

$\therefore$  the sq. on  $AB$  = the sum of the sqq. on  $BC, CA$ , with  
 twice the rect.  $BC, CD$ .

That is, the sq. on  $AB$  is greater than the sum of the  
 sqq. on  $BC, CA$  by twice the rect.  $BC, CD$ . Q.E.D.

## NOTE ON PROP. 12.

A general definition of the **projection of one straight line on another** is given on page 105. The student's attention is here called to a special case of projection which will enable us to simplify the Enunciation of Proposition 12.



In the above diagram, CA is a given straight line drawn from a point C in PQ; and from A a perpendicular AD is drawn to PQ. In this case, CD is said to be the **projection** of CA on PQ.

By applying this definition to the figure of Prop. 12, we see that the statement

*The sq. on AB is greater than the sum of the sqq. on BC, CA by twice the rect. BC, CD*

is the particular form of the following general Enunciation :

*In an obtuse-angled triangle the square on the side opposite the obtuse angle is greater than the sum of the squares on the sides containing the obtuse angle by twice the rectangle contained by one of those sides, and the **projection of the other side upon it.***

The Enunciation of Prop. 12 thus stated should be carefully compared with that of Prop. 13.

## PROPOSITION 13. THEOREM.

*In every triangle, the square on the side subtending an acute angle is less than the sum of the squares on the sides containing that angle, by twice the rectangle contained by either of these sides, and the straight line intercepted between the perpendicular let fall on it from the opposite angle, and the acute angle.*




Fig. 1.




Fig. 2.

Let  $ABC$  be any triangle having the angle at  $C$  an acute angle; and let  $AD$  be the perp. drawn from  $A$  to the opp. side  $BC$ .

*Then shall the sq. on  $AB$  be less than the sum of the sqq. on  $BC$ ,  $CA$ , by twice the rect.  $BC$ ,  $CD$ .*

*Proof.* Now  $AD$  may fall within the triangle  $ABC$ , as in fig. 1, or without it, as in fig. 2.

Because  $\left\{ \begin{array}{l} \text{in fig. 1, } BC \text{ is divided into two parts at } D, \\ \text{in fig. 2, } DC \text{ is divided into two parts at } B, \end{array} \right.$   
 $\therefore$  in both cases

the sum of the sqq. on  $BC$ ,  $CD$  = twice the rect.  $BC$ ,  $CD$  with the sq. on  $BD$ . II. 7.

To each of these equals add the sq. on  $DA$ .

Then the sum of the sqq. on  $BC$ ,  $CD$ ,  $DA$  = twice the rect.  $BC$ ,  $CD$  with the sum of the sqq. on  $BD$ ,  $DA$ .

But the sum of the sqq. on  $CD$ ,  $DA$  = the sq. on  $CA$ , I. 47.  
 for the angle  $ADC$  is a rt. angle.

Similarly, the sum of the sqq. on  $BD$ ,  $DA$  = the sq. on  $AB$ .

$\therefore$  the sum of the sqq. on  $BC$ ,  $CA$  = twice the rect.  $BC$ ,  $CD$  with the sq. on  $AB$ .

That is, the sq. on  $AB$  is less than the sum of the sqq. on  $BC$ ,  $CA$  by twice the rect.  $BC$ ,  $CD$ . Q.E.D.

*Obs.* If the perpendicular AD coincides with AB, that is, if ABC is a right angle, then *twice the rect.* BC, CD becomes *twice the sq. on BC*; and it may be shewn that the proposition merely repeats the result of I. 47.

## NOTES ON PROP. 13.

(i) Remembering the definition of the projection of a straight line given on p. 153, we may enunciate Prop. 13 as follows;

*In every triangle, the square on the side subtending an acute angle is less than the sum of the squares on the sides containing that angle, by twice the rectangle contained by one of these sides and the projection of the other side upon it.*

(ii) Comparing the Enunciations of II. 12, I. 47, II. 13, we see that in the triangle ABC,

if the angle ACB is *obtuse*, we have by II. 12,

$$AB^2 = BC^2 + CA^2 + 2BC \cdot CD;$$

if the angle ACB is a *right angle*, we have by I. 47,

$$AB^2 = BC^2 + CA^2;$$

if the angle ACB is *acute*, we have by II. 13,

$$AB^2 = BC^2 + CA^2 - 2BC \cdot CD.$$

These results may be collected as follows:

*The square on a side of a triangle is greater than, equal to, or less than the sum of the squares on the other sides, according as the angle opposite to the first is obtuse, a right angle, or acute.*

## EXERCISES ON II. 12 AND 13.


1. If from one of the base angles of an isosceles triangle a perpendicular is drawn to the opposite side, then twice the rectangle contained by that side and the segment adjacent to the base is equal to the square on the base.

2. If one angle of a triangle is one-third of two right angles, shew that the square on the opposite side is less than the sum of the squares on the sides forming that angle, by the rectangle contained by these two sides. [See Ex. 10, p. 109.]

3. If one angle of a triangle is two-thirds of two right angles, shew that the square on the opposite side is greater than the sum of the squares on the sides forming that angle, by the rectangle contained by these sides. [See Ex. 10, p. 109.]

## PROPOSITION 14. PROBLEM.

To describe a square that shall be equal to a given rectilinear figure.



Let *A* be the given rectilinear figure.  
It is required to describe a square equal to *A*.

**Construction.** Describe a par<sup>m</sup> *BCDE* equal to the fig. *A*,  
and having the angle *CBE* a right angle. I. 45.

Then if *BC* = *BE*, the fig. *BD* is a square; and what was  
required is done.

But if not, produce *BE* to *F*, making *EF* equal to *ED*; I. 3.  
and bisect *BF* at *G*. I. 10.

With centre *G*, and radius *GF*, describe the semicircle *BHF*;  
produce *DE* to meet the semicircle at *H*.

Then shall the sq. on *EH* be equal to the given fig. *A*.  
Join *GH*.

**Proof.** Because *BF* is divided equally at *G* and unequally  
at *E*,

∴ the rect. *BE*, *EF* with the sq. on *GE* = the sq. on *GF* II. 5.  
= the sq. on *GH*.

But the sq. on *GH* = the sum of the sqq. on *GE*, *EH*;  
for the angle *HEG* is a rt. angle. I. 47.

∴ the rect. *BE*, *EF* with the sq. on *GE* = the sum of the  
sqq. on *GE*, *EH*.

From these equals take the sq. on *GE* :

then the rect. *BE*, *EF* = the sq. on *HE*.

But the rect. *BE*, *EF* = the fig. *BD*; for *EF* = *ED*; *Constr.*  
and the fig. *BD* = the given fig. *A*. *Constr.*

∴ the sq. on *EH* = the given fig. *A*. Q.E.F.

QUESTIONS FOR REVISION ON BOOK II.

1. Explain the phrase, *the rectangle contained by*  $AB, CD$ ; and shew by superposition that if  $AB=PQ$ , and  $CD=RS$ , then the rectangle contained by  $AB, CD$  = the rectangle contained by  $PQ, RS$ .

2. Shew that Prop. 2 is a *special case* of Prop. 1, explaining under what conditions Prop. 1 becomes identical with Prop. 2.

3. What must be the relation between the divided and undivided lines in the enunciation of Prop. 1 in order to give the result proved in Prop. 3?

4. Define the *segments into which a straight line is divided at a point* in such a way as to be applicable to the case when the dividing point is in the given line produced.

Hence frame a statement which includes the enunciations of both II. 5 and II. 6, and find the algebraical formulæ corresponding to these enunciations.

Also combine in a single enunciation the results of I. 9 and II. 10.

5. Compare the results proved in Propositions 4 and 7 by finding the algebraical formulæ corresponding to their enunciations.

6. *The difference of the squares on two straight lines is equal to the rectangle contained by their sum and difference.* Deduce this theorem from Prop. 5.

7. Define *the projection of one straight line on another*.

How may the enunciations of II. 12 and II. 13 be simplified by means of this definition?

8. In the figure of Proposition 14,

(i) If  $BE=8$  inches, and  $ED=2$  inches, find the length of  $EH$ .

(ii) If  $BE=12\cdot5$  inches, and  $EH=2\cdot5$  inches, find the length of  $ED$ .

(iii) If  $BE=9$  inches, and  $EH=3$  inches, find the length of  $GH$ .

9. When is a straight line said to be divided in *medial section*?

If a straight line 8 inches in length is divided internally in medial section, shew that the lengths of the segments are approximately 4·9 inches and 3·1 inches.

[Frame a quadratic equation as explained on page 151, and solve.]



## THEOREMS AND EXAMPLES ON BOOK II.

## ON II. 4 AND 7.

1. *Shew by II. 4 that the square on a straight line is four times the square on half the line.*

[This result is constantly used in solving examples on Book II., especially those which follow from II. 12 and 13.]

2. If a straight line is divided into any three parts, the square on the whole line is equal to the sum of the squares on the three parts together with twice the rectangles contained by each pair of these parts.

Shew that the algebraical formula corresponding to this theorem is

$$(a + b + c)^2 = a^2 + b^2 + c^2 + 2bc + 2ca + 2ab.$$

3. *In a right-angled triangle, if a perpendicular is drawn from the right angle to the hypotenuse, the square on this perpendicular is equal to the rectangle contained by the segments of the hypotenuse.*

4. In an isosceles triangle, if a perpendicular is drawn from one of the angles at the base to the opposite side, shew that the square on the perpendicular is equal to twice the rectangle contained by the segments of that side together with the square on the segment adjacent to the base.

5. Any rectangle is half the rectangle contained by the diagonals of the squares described upon its two sides.

6. In any triangle if a perpendicular is drawn from the vertical angle to the base, the sum of the squares on the sides forming that angle, together with twice the rectangle contained by the segments of the base, is equal to the square on the base together with twice the square on the perpendicular.


## ON II. 5 AND 6.

*Obs.* The student is reminded that these important propositions are both included in the following enunciation :

*The difference of the squares on two straight lines is equal to the rectangle contained by their sum and difference.* [See Cor., p. 137].

7. In a right-angled triangle the square on one of the sides forming the right angle is equal to the rectangle contained by the sum and difference of the hypotenuse and the other side. [I. 47 and II. 5, Cor.]

8. *The difference of the squares on two sides of a triangle is equal to twice the rectangle contained by the base and the intercept between the middle point of the base and the foot of the perpendicular drawn from the vertical angle to the base.*



Let  $ABC$  be a triangle, and let  $P$  be the middle point of the base  $BC$ : let  $AQ$  be drawn perp. to  $BC$ .

*Then shall  $AB^2 - AC^2 = 2BC \cdot PQ$ .*

First, let  $AQ$  fall within the triangle.

$$\text{Now } AB^2 = BQ^2 + QA^2, \quad \text{I. 47.}$$

$$\text{also } AC^2 = QC^2 + QA^2,$$

$$\therefore AB^2 - AC^2 = BQ^2 - QC^2 \quad \text{Ax. 3.}$$

$$= (BQ + QC)(BQ - QC) \quad \text{II. 5, Cor.}$$

$$= BC \cdot 2PQ \quad \text{Ex., p. 137.}$$

$$= 2BC \cdot PQ \quad \text{Q.E.D.}$$

The case in which  $AQ$  falls outside the triangle presents no difficulty.

9. *The square on any straight line drawn from the vertex of an isosceles triangle to the base is less than the square on one of the equal sides by the rectangle contained by the segments of the base.*

10. The square on any straight line drawn from the vertex of an isosceles triangle to the base produced, is greater than the square on one of the equal sides by the rectangle contained by the segments into which the base is divided externally.

11. If a straight line is drawn through one of the angles of an equilateral triangle to meet the opposite side produced, so that the rectangle contained by the segments of the base is equal to the square on the side of the triangle; shew that the square on the line so drawn is double of the square on a side of the triangle.

12. If  $XY$  is drawn parallel to the base  $BC$  of an isosceles triangle  $ABC$ , then the difference of the squares on  $BY$  and  $CY$  is equal to the rectangle contained by  $BC$ ,  $XY$ . [See above, Ex. 8.]

13. In a right-angled triangle, if a perpendicular is drawn from the right angle to the hypotenuse, the square on either side forming the right angle is equal to the rectangle contained by the hypotenuse and the segment of it adjacent to that side.

## ON II. 9 AND 10.

14. Deduce Prop. 9 from Props. 4 and 5, using also the theorem that the square on a straight line is four times the square on half the line.

15. Deduce Prop. 10 from Props. 7 and 6, using also the theorem mentioned in the preceding Exercise.

16. If a straight line is divided equally, and also unequally, the squares on the two unequal segments are together equal to twice the rectangle contained by these segments together with four times the square on the line between the points of section.

## ON II. 11.

17. *If a straight line is divided internally in medial section, and from the greater segment a part be taken equal to the less; shew that the greater segment is also divided in medial section.*

18. If a straight line is divided in medial section, the rectangle contained by the sum and difference of the segments is equal to the rectangle contained by the segments.

19. If  $AB$  is divided at  $H$  in medial section, and if  $X$  is the middle point of the greater segment  $AH$ , shew that a triangle whose sides are equal to  $AH$ ,  $XH$ ,  $BX$  respectively must be right-angled.

20. If a straight line  $AB$  is divided internally in medial section at  $H$ , prove that the sum of the squares on  $AB$ ,  $BH$  is three times the square on  $AH$ .

21. *Divide a straight line externally in medial section.*


[Proceed as in II. 11, but instead of drawing  $EF$ , make  $EF'$  equal to  $EB$  in the direction remote from  $A$ ; and on  $AF'$  describe the square  $AF'G'H'$  on the side remote from  $AB$ . Then  $AB$  will be divided externally at  $H'$  as required.]

## ON II. 12 AND 13.

22. In a triangle  $ABC$  the angles at  $B$  and  $C$  are acute: if  $E$  and  $F$  are the feet of perpendiculars drawn from the opposite angles to the sides  $AC$ ,  $AB$ , shew that the square on  $BC$  is equal to the sum of the rectangles  $AB$ ,  $BF$  and  $AC$ ,  $CE$ .

23.  $ABC$  is a triangle right-angled at  $C$ , and  $DE$  is drawn from a point  $D$  in  $AC$  perpendicular to  $AB$ : shew that the rectangle  $AB$ ,  $AE$  is equal to the rectangle  $AC$ ,  $AD$ .

24. In any triangle the sum of the squares on two sides is equal to twice the square on half the third side together with twice the square on the median which bisects the third side.



Let  $ABC$  be a triangle, and  $AP$  the median bisecting the side  $BC$ .  
Then shall  $AB^2 + AC^2 = 2BP^2 + 2AP^2$ .

Draw  $AQ$  perp. to  $BC$ .

Consider the case in which  $AQ$  falls within the triangle, but does not coincide with  $AP$ .

Now of the angles  $APB$ ,  $APC$ , one must be obtuse, and the other acute: let  $APB$  be obtuse.

Then in the  $\triangle APB$ ,  $AB^2 = BP^2 + AP^2 + 2BP \cdot PQ$ , II. 12.

Also in the  $\triangle APC$ ,  $AC^2 = CP^2 + AP^2 - 2CP \cdot PQ$ . II. 13.

But  $CP = BP$ ,

$\therefore CP^2 = BP^2$ ; and the rect.  $BP, PQ =$  the rect.  $CP, PQ$ ,

Hence adding the above results,

$$AB^2 + AC^2 = 2 \cdot BP^2 + 2 \cdot AP^2.$$

Q. E. D.

The student will have no difficulty in adapting this proof to the cases in which  $AQ$  falls without the triangle, or coincides with  $AP$ .

25. The sum of the squares on the sides of a parallelogram is equal to the sum of the squares on the diagonals.

26. In any quadrilateral the squares on the diagonals are together equal to twice the sum of the squares on the straight lines joining the middle points of opposite sides. [See Ex. 9, p. 105.]

27. If from any point within a rectangle straight lines are drawn to the angular points, the sum of the squares on one pair of the lines drawn to opposite angles is equal to the sum of the squares on the other pair.

28. The sum of the squares on the sides of a quadrilateral is greater than the sum of the squares on its diagonals by four times the square on the straight line which joins the middle points of the diagonals.

29.  $O$  is the middle point of a given straight line  $AB$ , and from  $O$  as centre, any circle is described: if  $P$  be any point on its circumference, shew that the sum of the squares on  $AP$ ,  $BP$  is constant.

30. Given the base of a triangle, and the sum of the squares on the sides forming the vertical angle; find the locus of the vertex.

31.  $ABC$  is an isosceles triangle in which  $AB$  and  $AC$  are equal.  $AB$  is produced beyond the base to  $D$ , so that  $BD$  is equal to  $AB$ . Shew that the square on  $CD$  is equal to the square on  $AB$  together with twice the square on  $BC$ .

32. In a right-angled triangle the sum of the squares on the straight lines drawn from the right angle to the points of trisection of the hypotenuse is equal to five times the square on the line between the points of trisection.

33. Three times the sum of the squares on the sides of a triangle is equal to four times the sum of the squares on the medians.

34.  $ABC$  is a triangle, and  $O$  the point of intersection of its medians: shew that

$$AB^2 + BC^2 + CA^2 = 3(OA^2 + OB^2 + OC^2).$$

35.  $ABCD$  is a quadrilateral, and  $X$  the middle point of the straight line joining the bisections of the diagonals; with  $X$  as centre any circle is described, and  $P$  is any point upon this circle: shew that  $PA^2 + PB^2 + PC^2 + PD^2$  is constant, being equal to

$$XA^2 + XB^2 + XC^2 + XD^2 + 4XP^2.$$

36. The squares on the diagonals of a trapezium are together equal to the sum of the squares on its two oblique sides, with twice the rectangle contained by its parallel sides.

#### PROBLEMS.

37. Construct a rectangle equal to the difference of two squares.

38. Divide a given straight line into two parts so that the rectangle contained by them may be equal to the square described on a given straight line which is less than half the straight line to be divided.

39. Given a square and one side of a rectangle which is equal to the square, find the other side.

40. Produce a given straight line so that the rectangle contained by the whole line thus produced and the part produced, may be equal to the square on another given line.

41. Produce a given straight line so that the rectangle contained by the whole line thus produced and the given line shall be equal to the square on the part produced.


42. Divide a straight line  $AB$  into two parts at  $C$ , such that the rectangle contained by  $BC$  and another line  $X$  may be equal to the square on  $AC$ .

## BOOK III.

Book III. deals with the properties of Circles.

For convenience of reference the following definitions are repeated from Book I.

I. *Def.* 15. A **circle** is a plane figure bounded by one line, which is called the **circumference**, and is such that all straight lines drawn from a certain point within the figure to the circumference are equal to one another: this point is called the **centre** of the circle.



**NOTE.** Circles which have the same centre are said to be **concentric**.

I. *Def.* 16. A **radius** of a circle is a straight line drawn from the centre to the circumference.

I. *Def.* 17. A **diameter** of a circle is a straight line drawn through the centre, and terminated both ways by the circumference.

I. *Def.* 18. A **semicircle** is the figure bounded by a diameter of a circle and the part of the circumference cut off by the diameter.

**NOTE.** From these definitions we draw the following inferences:

(i) The distance of a point from the centre of a circle is less than the radius, if the point is within the circumference: and the distance of a point from the centre is greater than the radius, if the point is without the circumference.

(ii) A point is within a circle if its distance from the centre is less than the radius: and a point is without a circle if its distance from the centre is greater than the radius.

(iii) Circles of equal radius are equal in all respects; that is to say, their areas and circumferences are equal.


(iv) A circle is divided by any diameter into two parts which are equal in all respects.

## DEFINITIONS TO BOOK III.


1. An **arc** of a circle is any part of the circumference.
2. A **chord** of a circle is the straight line which joins any two points on the circumference.

**NOTE.** From these definitions it may be seen that a chord of a circle, which does not pass through the centre, divides the circumference into two unequal arcs; of these, the greater is called the **major arc**, and the less the **minor arc**. Thus the major arc is *greater*, and the minor arc *less* than the semi-circumference.


The major and minor arcs, into which a circumference is divided by a chord, are said to be **conjugate** to one another.




3. Chords of a circle are said to be **equidistant** from the centre, when the perpendiculars drawn to them from the centre are equal : and one chord is said to be **further from the centre** than another, when the perpendicular drawn to it from the centre is greater than the perpendicular drawn to the other.




4. A **secant** of a circle is a straight line of indefinite length, which cuts the circumference in two points.



5. A **tangent** to a circle is a straight line which meets the circumference, but being produced, does not cut it. Such a line is said to **touch** the circle at a point; and the point is called the **point of contact**.




**NOTE.** If a secant, which cuts a circle at the points P and Q, gradually changes its position in such a way that P remains fixed, the point Q will ultimately approach the fixed point P, until at length these points may be made to coincide. When the straight line PQ reaches this limiting position, it becomes the *tangent* to the circle at the point P.




Hence a tangent may be defined as a straight line which passes through *two coincident points* on the circumference.

6. Circles are said to **touch one another** when they meet, but do not cut one another.



**NOTE.** When each of the circles which meet is *outside the other*, they are said to touch one another **externally**, or to have **external contact**: when one of the circles is *within the other*, the first is said to touch the other **internally**, or to have **internal contact** with it.


7. A **segment** of a circle is the figure bounded by a chord and one of the two arcs into which the chord divides the circumference.



**NOTE.** The chord of a segment is sometimes called its base.




8. An angle in a segment is one formed by two straight lines drawn from any point in the arc of the segment to the extremities of its chord.



NOTE. (i) It will be shewn in Proposition 21, that all angles in the same segment of a circle are equal.

NOTE. (ii) The angle *of* a segment (as distinct from the angle *in* a segment) is sometimes defined as that which is contained between the *chord* and the *arc*; but this definition is not required in any proposition of Euclid.


9. An angle at the circumference of a circle is one formed by straight lines drawn from a point on the circumference to the extremities of an arc: such an angle is said to stand upon the arc by which it is subtended.



10. Similar segments of circles are those which contain equal angles.



11. A sector of a circle is a figure bounded by two radii and the arc intercepted between them.




#### SYMBOLS AND ABBREVIATIONS.

In addition to the symbols and abbreviations given on page 11, we shall use the following.

⊙ for circle,  $\bigcirc^{\infty}$  for circumference.

PROPOSITION 1. PROBLEM.

To find the centre of a given circle.



Let ABC be a given circle.

It is required to find the centre of the  $\odot$  ABC.

**Construction.** In the given circle draw any chord AB,  
and bisect AB at D. I. 10.

From D draw DC at right angles to AB; I. 11.  
and produce DC to meet the  $\odot^\infty$  at E and C.

Bisect EC at F. I. 10.

Then shall F be the centre of the  $\odot$  ABC.

**Proof.** First, the centre of the circle must be in EC :  
for if not, suppose the centre to be at a point G outside EC.  
Join AG, DG, BG.

Then in the  $\triangle$  GDA, GDB,

Because  $\left\{ \begin{array}{l} DA = DB, \\ \text{and GD is common;} \\ \text{and GA} = \text{GB, for by supposition they are radii;} \end{array} \right. \quad \text{Constr.}$

$\therefore$  the  $\angle$  GDA = the  $\angle$  GDB; I. 8.

$\therefore$  these angles, being adjacent, are rt. angles.

But the  $\angle$  CDB is a rt. angle; Constr.

$\therefore$  the  $\angle$  GDB = the  $\angle$  CDB, Ax. 11.

the part equal to the whole, which is impossible.

$\therefore$  G is not the centre.

So it may be shewn that no point outside EC is the centre ;


$\therefore$  the centre lies in EC.

$\therefore$  F, the middle point of the diameter EC, must be the  
centre of the  $\odot$  ABC. Q.E.F.

**COROLLARY.** The straight line which bisects a chord of a  
circle at right angles passes through the centre.

## PROPOSITION 2. THEOREM.

*If any two points are taken in the circumference of a circle, the chord which joins them falls within the circle.*



Let  $ABC$  be a circle, and  $A$  and  $B$  any two points on its  $\circ^{\infty}$ .

*Then shall the chord  $AB$  fall within the circle.*

**Construction.** Find  $D$ , the centre of the  $\odot ABC$ ; III. 1.  
and in  $AB$  take any point  $E$ .

Join  $DA$ ,  $DE$ ,  $DB$ .

**Proof.** In the  $\triangle DAB$ , because  $DA = DB$ , I. Def. 15.  
 $\therefore$  the  $\angle DAB =$  the  $\angle DBA$ . I. 5.

But the ext.  $\angle DEB$  is greater than the int. opp.  $\angle DAE$ ;  
I. 16.

$\therefore$  the  $\angle DEB$  is also greater than the  $\angle DBE$ .

$\therefore$  in the  $\triangle DEB$ , the side  $DB$ , which is opposite the greater angle, is greater than  $DE$  which is opposite the less: I. 19.

that is to say,  $DE$  is less than  $DB$ , a radius of the circle;

$\therefore E$  falls within the circle.

Similarly, any other point between  $A$  and  $B$  may be shewn to fall within the circle.

$\therefore AB$  falls within the circle. Q.E.D.


**NOTE.** A part of a curved line is said to be **concave** to a point, when for every chord (taken so as to lie between the point and the curve) all straight lines drawn from the given point to the intercepted arc are cut by the chord: if, when any chord whatever is taken, no straight line drawn from the given point to the intercepted arc is cut by the chord, the curve is said to be **convex** to that point.

Proposition 2 proves that the whole circumference of a circle is *concave to its centre*.

PROPOSITION 3. THEOREM.

If a straight line drawn through the centre of a circle bisects a chord which does not pass through the centre, it shall cut the chord at right angles.

Conversely, if it cuts the chord at right angles, it shall bisect it.



Let ABC be a circle; and let CD be a st. line drawn through the centre, and AB a chord which does not pass through the centre.

*First.* Let CD bisect the chord AB at F.  
Then shall CD cut AB at rt. angles.

**Construction.** Find E the centre of the circle; III. 1.  
and join EA, EB.

**Proof.** Then in the  $\triangle$  AFE, BFE,  
Because  $\left\{ \begin{array}{l} AF = BF, \\ \text{and FE is common;} \\ \text{and AE} = \text{BE, being radii of the circle;} \\ \therefore \text{the } \angle \text{AFE} = \text{the } \angle \text{BFE;} \\ \therefore \text{these angles, being adjacent, are rt. angles;} \\ \text{that is, DC cuts AB at rt. angles.} \end{array} \right. \begin{array}{l} \text{Hyp.} \\ \\ \\ \text{I. 8.} \\ \text{Q.E.D.} \end{array}$


*Conversely.* Let CD cut the chord AB at rt. angles.  
Then shall CD bisect AB at F.

**Construction.** Find E the centre; and join EA, EB.

**Proof.** In the  $\triangle$  EAB, because EA = EB, I. Def. 15.  
 $\therefore$  the  $\angle$  EAB = the  $\angle$  EBA. I. 5.  
Then in the  $\triangle$  EFA, EFB,  
Because  $\left\{ \begin{array}{l} \text{the } \angle \text{EAF} = \text{the } \angle \text{EBF,} \\ \text{and the } \angle \text{EFA} = \text{the } \angle \text{EFB, being rt. angles;} \\ \text{and EF is common;} \\ \therefore \text{AF} = \text{BF;} \\ \text{that is, CD bisects AB at F.} \end{array} \right. \begin{array}{l} \text{Proved.} \\ \text{Hyp.} \\ \\ \text{I. 26.} \\ \text{Q.E.D.} \end{array}$

## PROPOSITION 4. THEOREM.

*If in a circle two chords cut one another, which do not both pass through the centre, they cannot both be bisected at their point of intersection.*



Let  $ABCD$  be a circle, and  $AC$ ,  $BD$  two chords which intersect at  $E$ , but do not both pass through the centre.

*Then  $AC$  and  $BD$  shall not be both bisected at  $E$ .*

**CASE I.** If *one* chord passes through the centre, it is a diameter, and the centre is its middle point ;  
 $\therefore$  it cannot be bisected by the other chord, which by hypothesis does not pass through the centre.

**CASE II.** If neither chord passes through the centre ; then, if possible, suppose  $E$  to be the middle point of *both* ; that is, let  $AE = EC$  ; and  $BE = ED$ .

**Construction.** Find  $F$ , the centre of the circle. III. 1.  
 Join  $EF$ .

**Proof.** Because  $FE$ , which passes through the centre, bisects the chord  $AC$ ,  
 $\therefore$  the  $\angle FEC$  is a rt. angle. *Hyp.* III. 3.

And because  $FE$ , which passes through the centre, bisects the chord  $BD$ ,  
 $\therefore$  the  $\angle FED$  is a rt. angle. *Hyp.* III. 3.  
 $\therefore$  the  $\angle FEC =$  the  $\angle FED$ ,

the whole equal to its part, which is impossible.

$\therefore AC$  and  $BD$  are not *both* bisected at  $E$ . Q.E.D.

## EXERCISES.

## ON PROPOSITION 1.

1. If two circles intersect at the points **A**, **B**, shew that the line which joins their centres bisects their common chord **AB** at right angles.

2. **AB**, **AC** are two equal chords of a circle; shew that the straight line which bisects the angle **BAC** passes through the centre.

3. *Two chords of a circle are given in position and magnitude: find the centre of the circle.*

4. *Describe a circle that shall pass through three given points, which are not in the same straight line.*

5. *Find the locus of the centres of circles which pass through two given points.*

6. Describe a circle that shall pass through two given points, and have a given radius. When is this impossible?

## ON PROPOSITION 2.

7. *A straight line cannot cut a circle in more than two points.*

## ON PROPOSITION 3.

8. Through a given point within a circle draw a chord which shall be bisected at that point.

9. The parts of a straight line intercepted between the circumferences of two concentric circles are equal.

10. The line joining the middle points of two parallel chords of a circle passes through the centre.


11. Find the locus of the middle points of a system of parallel chords drawn in a circle.

12. If two circles cut one another, any two parallel straight lines drawn through the points of intersection to cut the circles, are equal.

13. **PQ** and **XY** are two parallel chords in a circle: shew that the points of intersection of **PX**, **QY**, and of **PY**, **QX**, lie on the straight line which passes through the middle points of the given chords.

## PROPOSITION 5. THEOREM.

*If two circles cut one another, they cannot have the same centre.*



Let the two  $\odot^s$  AGC, BFC cut one another at C.

*Then they shall not have the same centre.*

**Construction.** If possible, let the two circles have the same centre; and let it be called E.

Join EC;

and from E draw any st. line to meet the  $\odot^{ces}$  at F and G.

**Proof.** Because E is the centre of the  $\odot$  AGC, *Hyp.*

$\therefore EG = EC.$

And because E is also the centre of the  $\odot$  BFC, *Hyp.*

$\therefore EF = EC.$

$\therefore EG = EF,$

the whole equal to its part, which is impossible.

Therefore the two circles have not the same centre.

Q.E.D.


## EXERCISES.

## ON PROPOSITIONS 4 AND 5.

1. If a parallelogram can be inscribed in a circle, the point of intersection of its diagonals must be at the centre of the circle.
2. Rectangles are the only parallelograms that can be inscribed in a circle.
3. Two circles, which intersect at one point, must also intersect at another.

PROPOSITION 6. THEOREM.

*If two circles touch one another internally, they cannot have the same centre.*



Let the two  $\odot^s$  ABC, DEC touch one another internally at C.

*Then they shall not have the same centre.*

**Construction.** If possible, let the two circles have the same centre; and let it be called F.

Join FC;

and from F draw any st. line to meet the  $\odot^{oes}$  at E and B.

**Proof.** Because F is the centre of the  $\odot$  ABC, *Hyp.*  
 $\therefore$  FB = FC.

And because F is the centre of the  $\odot$  DEC, *Hyp.*  
 $\therefore$  FE = FC.

$\therefore$  FB = FE,

the whole equal to its part, which is impossible.

Therefore the two circles have not the same centre.

Q.E.D.

**NOTE.** From Propositions 5 and 6 it is seen that circles, whose circumferences have any point in common, cannot be concentric, unless they coincide entirely.

Conversely, the circumferences of concentric circles can have no point in common.




## PROPOSITION 7. THEOREM.

If from any point within a circle which is not the centre, straight lines are drawn to the circumference, then the greatest is that which passes through the centre; and the least is the remaining part of the diameter.

And of all other such lines, that which is nearer to the greatest is always greater than one more remote.

And two equal straight lines, and only two, can be drawn from the given point to the circumference, one on each side of the diameter.



Let ABCD be a circle, and from F, any point within it which is not the centre, let FA, FB, FC, FG, and FD be drawn to the  $\circ^{\infty}$ , of which FA passes through E the centre, and FD is the remaining part of the diameter.

Then of all these st. lines,

- (i) FA shall be the greatest;
- (ii) FD shall be the least;
- (iii) FB, which is nearer to FA, shall be greater than FC, which is more remote;
- (iv) also two, and only two, equal st. lines can be drawn from F to the  $\circ^{\infty}$ .

**Construction.** Join EB, EC.

**Proof.** (i) In the  $\triangle FEB$ , the two sides FE, EB are together greater than the third side FB. I. 20.

But  $EB = EA$ , being radii of the circle;  
 $\therefore$  FE, EA are together greater than FB;  
 that is, FA is greater than FB.

Similarly FA may be shewn to be greater than any other st. line drawn from F to the  $\circ^\infty$ ;

$\therefore$  FA is the greatest of all such lines.

(ii) In the  $\triangle$  EFG, the two sides EF, FG are together greater than EG; I. 20.

and EG = ED, being radii of the circle;

$\therefore$  EF, FG are together greater than ED.

Take away the common part EF;

then FG is greater than FD.

Similarly any other st. line drawn from F to the  $\circ^\infty$  may be shewn to be greater than FD;

$\therefore$  FD is the least of all such lines.

(iii) In the  $\triangle^s$  BEF, CEF, I. Def. 15.

Because  $\left\{ \begin{array}{l} BE = CE, \\ \text{and } EF \text{ is common;} \\ \text{but the } \angle BEF \text{ is greater than the } \angle CEF; \end{array} \right.$

$\therefore$  FB is greater than FC. I. 24.

Similarly it may be shewn that FC is greater than FG.

(iv) Join EG, and at E in FE make the  $\angle FEH$  equal to the  $\angle FEG$ . I. 23.

Join FH.

Then in the  $\triangle^s$  GEF, HEF,

Because  $\left\{ \begin{array}{l} GE = HE, \\ \text{and } EF \text{ is common;} \\ \text{also the } \angle GEF = \text{the } \angle HEF; \end{array} \right.$  I. Def. 15.

$\therefore$  FG = FH. I. 4.

And besides FH no other straight line can be drawn from F to the  $\circ^\infty$  equal to FG.

For, if possible, let FK = FG.

Then, because FH = FG, Proved.

$\therefore$  FK = FH,

that is, a line nearer to FA, the greatest, is equal to a line which is more remote; which is impossible. Proved.


Therefore two, and only two, equal st. lines can be drawn from F to the  $\circ^\infty$ . Q.E.D.

## PROPOSITION 8. THEOREM.

If from any point without a circle straight lines are drawn to the circumference, of those which fall on the concave circumference, the greatest is that which passes through the centre; and of others, that which is nearer to the greatest is always greater than one more remote.

Of those which fall on the convex circumference, the least is that which, when produced, passes through the centre; and of others, that which is nearer to the least is always less than one more remote.

From the given point there can be drawn to the circumference two, and only two, equal straight lines, one on each side of the shortest line.



Let BGD be a circle; and from A, any point outside the circle, let ABD, AEH, AFG, be drawn, of which AD passes through C, the centre, and AH is nearer than AG to AD.

Then of st. lines drawn from A to the concave  $\circ^{\infty}$ ,

(i) AD shall be the greatest, and (ii) AH greater than AG.

And of st. lines drawn from A to the convex  $\circ^{\infty}$ ,

(iii) AB shall be the least, and (iv) AE less than AF.

(v) Also two, and only two, equal st. lines can be drawn from A to the  $\circ^{\infty}$ .

Construction. Join CH, CG, CF, CE.

Proof. (i) In the  $\triangle ACH$ , the two sides AC, CH are together greater than AH: I. 20.

but CH = CD, being radii of the circle;

$\therefore$  AC, CD are together greater than AH:

that is, AD is greater than AH.

Similarly AD may be shewn to be greater than any other st. line drawn from A to the concave  $\circ^{\infty}$ ;

$\therefore$  AD is the greatest of all such lines.

(ii) In the  $\triangle^s$  HCA, GCA,  
 Because  $\left\{ \begin{array}{l} \text{HC} = \text{GC}, \\ \text{and CA is common;} \\ \text{but the } \angle \text{HCA is greater than the } \angle \text{GCA;} \\ \therefore \text{AH is greater than AG.} \end{array} \right. \begin{array}{l} \text{I. Def. 15.} \\ \\ \\ \text{I. 24.} \end{array}$

(iii) In the  $\triangle$  AEC, the two sides AE, EC are together greater than AC;  
 but EC = BC; I. Def. 15.  
 $\therefore$  the remainder AE is greater than the remainder AB.

Similarly any other st. line drawn from A to the convex  $\circ^\infty$  may be shewn to be greater than AB;  
 $\therefore$  AB is the least of all such lines.

(iv) In the  $\triangle$  AFC, because AE, EC are drawn from the extremities of the base to a point E within the triangle,  
 $\therefore$  AF, FC are together greater than AE, EC. I. 21.  
 But FC = EC; I. Def. 15.  
 $\therefore$  the remainder AF is greater than the remainder AE.

(v) At C, in AC, make the  $\angle$  ACM equal to the  $\angle$  ACE.  
 Join AM.

Then in the two  $\triangle^s$  ECA, MCA,  
 Because  $\left\{ \begin{array}{l} \text{EC} = \text{MC}, \\ \text{and CA is common;} \\ \text{also the } \angle \text{ECA} = \text{the } \angle \text{MCA;} \\ \therefore \text{AE} = \text{AM.} \end{array} \right. \begin{array}{l} \text{I. Def. 15.} \\ \\ \text{Constr.} \\ \text{I. 4.} \end{array}$

And besides AM, no st. line can be drawn from A to the  $\circ^\infty$ , equal to AE.

For, if possible, let AK = AE :  
 then because AM = AE, Proved.  
 $\therefore$  AM = AK;


that is, a line nearer to AB, the shortest line, is equal to a line which is more remote; which is impossible. Proved.

Therefore two, and only two, equal st. lines can be drawn from A to the  $\circ^\infty$ . Q.E.D.

EXERCISE. Where are the limits of that part of the circumference which is concave to the point A?

## PROPOSITION 9. THEOREM. [FIRST PROOF.]

If from a point within a circle more than two equal straight lines can be drawn to the circumference, that point is the centre of the circle.



Let  $ABC$  be a circle, and  $D$  a point within it, from which more than two equal st. lines are drawn to the  $\circ^{\circ}$ , namely  $DA$ ,  $DB$ ,  $DC$ .

Then  $D$  shall be the centre of the circle  $ABC$ .

**Construction.** Join  $AB$ ,  $BC$  :  
and bisect  $AB$ ,  $BC$  at  $E$  and  $F$  respectively. I. 10.  
Join  $DE$ ,  $DF$ .

**Proof.** In the  $\triangle^s$   $DEA$ ,  $DEB$ ,  
Because  $\begin{cases} EA = EB, \\ \text{and } DE \text{ is common;} \\ \text{and } DA = DB; \end{cases}$  *Constr.*  
 $\therefore$  the  $\angle DEA = \angle DEB$ ; *Hyp.* I. 8.  
 $\therefore$  these angles, being adjacent, are rt. angles.

Hence  $ED$ , which bisects the chord  $AB$  at rt. angles, must pass through the centre. III. 1. *Cor.*


Similarly it may be shewn that  $FD$  passes through the centre.

$\therefore D$ , which is the only point common to  $ED$  and  $FD$ , must be the centre. Q.E.D.

**NOTE.** Of the two proofs of this proposition given by Euclid the first has the advantage of being *direct*.

PROPOSITION 9. THEOREM. [SECOND PROOF.]

*If from a point within a circle more than two equal straight lines can be drawn to the circumference, that point is the centre of the circle.*



Let ABC be a circle, and D a point within it, from which more than two equal st. lines are drawn to the  $\circ^\infty$ , namely DA, DB, DC.

*Then D shall be the centre of the circle ABC.*

**Construction.** For if not, suppose, if possible, E to be the centre.

Join DE, and produce it to meet the  $\circ^\infty$  at F, G.

**Proof.** Because D is a point within the circle, not the centre, and because DF passes through the centre E ;

$\therefore$  DA, which is nearer to DF, is greater than DB, which is more remote : III. 7.

but this is impossible, since by hypothesis, DA, DB, are equal.

$\therefore$  E is not the centre of the circle.


\* And wherever we suppose the centre E to be, otherwise than at D, two at least of the st. lines DA, DB, DC may be shewn to be unequal, which is contrary to hypothesis.

$\therefore$  D is the centre of the  $\odot$  ABC. Q.E.D.

\* **NOTE.** For example, if the centre E were supposed to be within the angle BDC, then DB would be greater than DA; if within the angle ADB, then DB would be greater than DC; if on one of the three straight lines, as DB, then DB would be greater than both DA and DC.

## PROPOSITION 10. THEOREM. [FIRST PROOF.]

*One circle cannot cut another at more than two points.*



If possible, let  $DABC$ ,  $EABC$  be two circles, cutting one another at more than two points, namely at  $A$ ,  $B$ ,  $C$ .

**Construction.** Join  $AB$ ,  $BC$ .

Draw  $FH$ , bisecting  $AB$  at rt. angles; I. 10, 11.  
and draw  $GH$  bisecting  $BC$  at rt. angles.

**Proof.** Because  $AB$  is a chord of *both* circles, and because  $FH$  bisects  $AB$  at rt. angles,  
 $\therefore$  the centre of both circles lies in  $FH$ . III. 1. *Cor.*

Again, because  $BC$  is a chord of both circles, and because  $GH$  bisects  $BC$  at right angles,  
 $\therefore$  the centre of both circles lies in  $GH$ . III. 1. *Cor.*

Hence  $H$ , the only point common to  $FH$  and  $GH$ , is the centre of both circles;  
which is impossible, for circles which cut one another cannot have a common centre. III. 5.

Therefore one circle cannot cut another at more than two points. Q.E.D.


**COROLLARIES.** (i) *Two circles cannot have three points in common without coinciding entirely.*

(ii) *Two circles cannot have a common arc without coinciding entirely.*

(iii) *Only one circle can be described through three points, which are not in the same straight line.*

PROPOSITION 10. THEOREM. [SECOND PROOF.]

*One circle cannot cut another at more than two points.*



If possible, let  $DABC$ ,  $EABC$  be two circles, cutting one another at more than two points, namely at  $A$ ,  $B$ ,  $C$ .

**Construction.** Find  $H$ , the centre of the  $\odot DABC$ , III. 1. and join  $HA$ ,  $HB$ ,  $HC$ .

**Proof.** Since  $H$  is the centre of the  $\odot DABC$ ,  
 $\therefore HA$ ,  $HB$ ,  $HC$  are all equal. I. Def. 15.

And because  $H$  is a point within the  $\odot EABC$ , from which more than two equal st. lines, namely  $HA$ ,  $HB$ ,  $HC$  are drawn to the  $\odot$ ,

$\therefore H$  is the centre of the  $\odot EABC$ : III. 9.

that is to say, the two circles have a common centre  $H$ ;  
 but this is impossible, since they cut one another. III. 5.

Therefore one circle cannot cut another in more than two points. Q.E.D.


**NOTE.** Both the proofs of Proposition 10 given by Euclid are indirect.

The second of these is imperfect, because it assumes that the centre of the circle  $DABC$  must fall within the circle  $EABC$ ; whereas it may be conceived to fall either without the circle  $EABC$ , or on its circumference. Hence to make the proof complete, two additional cases are required.



## PROPOSITION 11. THEOREM.

If two circles touch one another internally, the straight line which joins their centres, being produced, shall pass through the point of contact.



Let ABC and ADE be two circles which touch one another internally at A; let F be the centre of the  $\odot$  ABC, and G the centre of the  $\odot$  ADE.

Then shall FG produced pass through A.

**Construction.** For if not, suppose, if possible, FG to pass otherwise, as FGEH.

Join FA, GA.

**Proof.** In the  $\triangle$  FGA, the two sides FG, GA are together greater than FA :

but FA = FH, being radii of the  $\odot$  ABC : I. 20.

FG, GA are together greater than FH.

Take away the common part FG :

then GA is greater than GH.

But GA = GE, being radii of the  $\odot$  ADE : Hyp.

$\therefore$  GE is greater than GH,

the part greater than the whole ; which is impossible.

$\therefore$  FG, when produced, must pass through A.

Q.E.D.


## EXERCISES.

1. If the distance between the centres of two circles is equal to the difference of their radii, then the circles must meet in one point, but in no other ; that is, they must touch one another.

2. If two circles whose centres are A and B touch one another internally, and a straight line is drawn through their point of contact, cutting the circumferences at P and Q ; shew that the radii AP and BQ are parallel.

PROPOSITION 12. THEOREM.

*If two circles touch one another externally, the straight line which joins their centres shall pass through the point of contact.*



Let ABC and ADE be two circles which touch one another externally at A; let F be the centre of the  $\odot$  ABC, and G the centre of the  $\odot$  ADE.

*Then shall FG pass through A.*

**Construction.** For if not, suppose, if possible, FG to pass otherwise, as FHKG.

Join FA, GA.

**Proof.** In the  $\triangle$  FAG, the two sides FA, GA are together greater than FG:

but FA = FH, being radii of the  $\odot$  ABC; *Hyp.*

and GA = GK, being radii of the  $\odot$  ADE; *Hyp.*

$\therefore$  FH and GK are together greater than FG;  
which is impossible.

$\therefore$  FG must pass through A.


Q.E.D.

EXERCISES.

1. Find the locus of the centres of all circles which touch a given circle at a given point.
2. Find the locus of the centres of all circles of given radius, which touch a given circle.
3. If the distance between the centres of two circles is equal to the sum of their radii, then the circles meet in one point, but in no other; that is, they touch one another.
4. If two circles whose centres are A and B touch one another externally, and a straight line is drawn through their point of contact cutting the circumferences at P and Q; shew that the radii AP and BQ are parallel.

## PROPOSITION 13. THEOREM.

Two circles cannot touch one another at more than one point, whether internally or externally.



If possible, let ABC, EDF be two circles which touch one another at more than one point, namely at B and D.

**Construction.** Join BD ;  
and draw GF, bisecting BD at rt. angles. I. 10, 11.

**Proof.** Now, whether the circles touch one another internally, as in Fig 1 or externally as in Fig 2,

because B and D are on the  $\circ^{\text{ces}}$  of both circles,

$\therefore$  BD is a chord of both circles :

$\therefore$  the centres of both circles lie in GF, which bisects BD at rt. angles. III. 1. Cor.

Hence GF which joins the centres must pass through a point of contact ; III. 11, and 12.

which is impossible, since B and D are outside GF.

Therefore two circles cannot touch one another at more than one point.

Q.E.D.


**NOTE.** It must be observed that the proof here given applies, by virtue of Propositions 11 and 12, to *both* the above figures : we have therefore omitted the separate discussion of Fig. 2, which finds a place in most editions based on Simson's text.

## EXERCISES ON PROPOSITIONS 1-13.

1. Describe a circle to pass through two given points and have its centre on a given straight line. When is this impossible?
2. All circles which pass through a fixed point, and have their centres on a given straight line, pass also through a second fixed point.
3. Describe a circle of given radius to touch a given circle at a given point. How many solutions will there be? When will there be only one solution?
4. From a given point as centre describe a circle to touch a given circle. How many solutions will there be?
5. Describe a circle to pass through a given point, and touch a given circle at a given point. [See Ex. 1, p. 183, and Ex. 5, p. 171.] When is this impossible?
6. Describe a circle of given radius to touch two given circles. [See Ex. 2, p. 183.] How many solutions will there be?
7. Two parallel chords of a circle are six inches and eight inches in length respectively, and the perpendicular distance between them is one inch: find the radius.
8. If two circles touch one another externally, the straight lines, which join the extremities of parallel diameters towards opposite parts, must pass through the point of contact.
9. Find the greatest and least straight lines which have one extremity on each of two given circles, which do not intersect.
10. In any segment of a circle, of all straight lines drawn at right angles to the chord and intercepted between the chord and the arc, the greatest is that which passes through the middle point of the chord; and of others that which is nearer the greatest is greater than one more remote.
11. If from any point on the circumference of a circle straight lines be drawn to the circumference, the greatest is that which passes through the centre; and of others, that which is nearer to the greatest is greater than one more remote; and from this point there can be drawn to the circumference two, and only two, equal straight lines.

## PROPOSITION 14. THEOREM.

*Equal chords in a circle are equidistant from the centre.  
Conversely, chords which are equidistant from the centre  
are equal.*



Let  $ABC$  be a circle, and let  $AB$  and  $CD$  be chords, of which the perp. distances from the centre are  $EF$  and  $EG$ .

*First.* Let  $AB = CD$ .

*Then shall  $AB$  and  $CD$  be equidistant from the centre  $E$ .*

**Construction.** Join  $EA, EC$ .

**Proof.** Because  $EF$ , which passes through the centre, is perp. to the chord  $AB$ ;

$\therefore EF$  bisects  $AB$ ;

*Hyp.*  
III. 3.

$\therefore AB$  is double of  $FA$ .

For a similar reason,  $CD$  is double of  $GC$ .

But  $AB = CD$ ,

$\therefore FA = GC$ .

*Hyp.*  
*Ax.* 7.

Now  $EA = EC$ , being radii of the circle;

$\therefore$  the sq. on  $EA =$  the sq. on  $EC$ .

But since the  $\angle EFA$  is a rt. angle:

$\therefore$  the sq. on  $EA =$  the sqq. on  $EF, FA$ . I. 47.

And since the  $\angle EGC$  is a rt. angle;

$\therefore$  the sq. on  $EC =$  the sqq. on  $EG, GC$ .

$\therefore$  the sqq. on  $EF, FA =$  the sqq. on  $EG, GC$ .

Now of these, the sq. on  $FA =$  the sq. on  $GC$ ; for  $FA = GC$ .

$\therefore$  the sq. on  $EF =$  the sq. on  $EG$ ;

$\therefore EF = EG$ ;

that is, the chords  $AB, CD$  are equidistant from the centre.

Q.E.D.

*Conversely.* Let AB and CD be equidistant from the centre E ;

that is, let  $EF = EG$ .

*Then shall*  $AB = CD$ .

**Proof.** The same construction being made, it may be shewn as before that AB is double of FA, and CD double of GC ;

and that the sqq. on EF, FA = the sqq. on EG, GC.

Now of these, the sq. on EF = the sq. on EG,

for  $EF = EG$  :

*Hyp.*

$\therefore$  the sq. on FA = the sq. on GC ;

$\therefore$   $FA = GC$  ;

and doubles of these equals are equal ; *Ax. 6.*

that is,  $AB = CD$ .

Q.E.D.


#### EXERCISES.

1. Find the locus of the middle points of equal chords of a circle.
2. If two chords of a circle cut one another, and make equal angles with the straight line which joins their point of intersection to the centre, they are equal.
3. If two equal chords of a circle intersect, shew that the segments of the one are equal respectively to the segments of the other.
4. In a given circle draw a chord which shall be equal to one given straight line (not greater than the diameter) and parallel to another.
5. PQ is a fixed chord in a circle, and AB is any diameter : shew that the sum or difference of the perpendiculars let fall from A and B on PQ is constant, that is, the same for all positions of AB.

## PROPOSITION 15. THEOREM.

*The diameter is the greatest chord in a circle ;  
and of other chords, that which is nearer to the centre is  
greater than one more remote.*

*Conversely, the greater chord is nearer to the centre than  
the less.*



Let ABCD be a circle of which AD is a diameter, and E the centre ; and let BC and FG be any two chords, whose perp. distances from the centre are EH and EK.

*Then* (i) AD shall be greater than BC ;  
(ii) if EH is less than EK, BC shall be greater than FG :  
(iii) if BC is greater than FG, EH shall be less than EK.

(i) **Construction.** Join EB, EC.

**Proof.** In the  $\triangle BEC$ , the two sides BE, EC are together greater than BC ; I. 20.  
but BE = AE, I. Def. 15.  
and EC = ED ;

$\therefore$  AE and ED together are greater than BC ;  
that is, AD is greater than BC.

Similarly AD may be shewn to be greater than any other chord, not a diameter.

(ii) Let EH be less than EK.  
*Then BC shall be greater than FG.*

**Construction.** Join EF.

**Proof.** Since EH, passing through the centre, is perp. to the chord BC,

$\therefore$  EH bisects BC ; III. 3.

$\therefore$  BC is double of HB.

For a similar reason FG is double of FK.

Now EB = EF,

I. Def. 15.

$\therefore$  the sq. on EB = the sq. on EF.

But since the  $\angle$  EHB is a rt. angle,

$\therefore$  the sq. on EB = the sqq. on EH, HB.

I. 47.

And since the  $\angle$  EKF is a rt. angle,

$\therefore$  the sq. on EF = the sqq. on EK, KF;

$\therefore$  the sqq. on EH, HB = the sqq. on EK, KF.

But the sq. on EH is less than the sq. on EK,  
for EH is less than EK;

Hyp.

$\therefore$  the sq. on HB is greater than the sq. on KF;

$\therefore$  HB is greater than KF;

hence BC is greater than FG.

(iii) *Conversely.* Let BC be greater than FG.

*Then EH shall be less than EK.*

*Proof.* The same construction being made, it may be shewn as before that BC is double of BH. and FG double of FK; and that the sqq. on EH, HB = the sqq. on EK, KF.

But since BC is greater than FG,

Hyp.

$\therefore$  HB is greater than KF;

$\therefore$  the sq. on HB is greater than the sq. on KF.

$\therefore$  the sq. on EH is less than the sq. on EK;

$\therefore$  EH is less than EK.

Q.E.D.

#### EXERCISES.

1. *Through a given point within a circle draw the least possible chord.*

2. AB is a fixed chord of a circle, and XY any other chord having its middle point Z on AB; what is the greatest, and what the least length that XY may have?

Shew that XY increases, as Z approaches the middle point of AB.


3. In a given circle draw a chord of given length, having its middle point on a given chord.

When is this problem impossible?



## PROPOSITION 16. THEOREM. [ALTERNATIVE PROOF.]

The straight line drawn at right angles to a diameter of a circle at one of its extremities is a tangent to the circle :  
and every other straight line drawn through this point cuts the circle.



Let  $AKB$  be a circle, of which  $E$  is the centre, and  $AB$  a diameter ; and through  $B$  let the st. line  $CBD$  be drawn at rt. angles to  $AB$ .

Then (i)  $CBD$  shall be a tangent to the circle ;

(ii) any other st. line through  $B$ , such as  $BF$ , shall cut the circle.

(i) Construction. In  $CD$  take any point  $G$ , and join  $EG$ .

Proof. In the  $\triangle EBG$ , the  $\angle EBG$  is a rt. angle ; *Hyp.*

$\therefore$  the  $\angle EGB$  is less than a rt. angle ; I. 17.

$\therefore$  the  $\angle EBG$  is greater than the  $\angle EGB$  ;

$\therefore$   $EG$  is greater than  $EB$  : I. 19.

that is,  $EG$  is greater than a radius of the circle ;

$\therefore$  the point  $G$  is without the circle.

Similarly any other point in  $CD$ , except  $B$ , may be shewn to be outside the circle.

Hence  $CD$  meets the circle at  $B$ , but being produced, does not cut it ;

that is,  $CD$  is a tangent to the circle. III. Def. 5.

(ii) Construction. Draw  $EH$  perp. to  $BF$ . I. 12.

Proof. In the  $\triangle EHB$ , because the  $\angle EHB$  is a rt. angle,

$\therefore$  the  $\angle EBH$  is less than a rt. angle ; I. 17.

$\therefore$   $EB$  is greater than  $EH$  ; I. 19.

that is, EH is less than a radius of the circle :

$\therefore$  H, a point in BF, is within the circle ;


$\therefore$  BF must cut the circle. Q.E.D.

**NOTE.** The above proof of Proposition 16 is not that given by Euclid, but it is preferable as being *direct*. Euclid's proof by *Reductio ad Absurdum* is given below.

PROPOSITION 16. THEOREM. [EUCLID'S PROOF.]

*The straight line drawn at right angles to a diameter of a circle at one of its extremities, is a tangent to the circle :*

*and no other straight line can be drawn through this point so as not to cut the circle.*



Let ABC be a circle, of which D is the centre, and AB a diameter ; let AE be drawn at rt. angles to BA, at its extremity A.

(i) *Then shall AE be a tangent to the circle.*

**Construction.**

For, if possible, suppose AE to cut the circle at C.

Join DC.

**Proof.** Then in the  $\triangle DAC$ , because  $DA=DC$ , I. Def. 15.

$\therefore$  the  $\angle DAC =$  the  $\angle DCA$ .

But the  $\angle DAC$  is a rt. angle ; Hyp.


$\therefore$  the  $\angle DCA$  is a rt. angle ;

that is, two angles of the  $\triangle DAC$  are together equal to two rt. angles ; which is impossible. I. 17.

Hence AE meets the circle at A, but being produced, does not cut it ;

that is, AE is a tangent to the circle. III. Def. 5.

(ii) *Also through A no other straight line but AE can be drawn so as not to cut the circle.*



**Construction.** For, if possible, let AF be another st. line drawn through A so as not to cut the circle.

From D draw DG perp. to AF ; i. 12.  
and let DG meet the  $\circ^{\text{ce}}$  at H.

**Proof.** Then in the  $\triangle$  DAG, because the  $\angle$  DGA is a rt. angle,  
 $\therefore$  the  $\angle$  DAG is less than a rt. angle ; i. 17.

$\therefore$  DA is greater than DG. i. 19.

But DA = DH, i. Def. 15.

$\therefore$  DH is greater than DG,

the part greater than the whole, which is impossible.

Therefore no st. line can be drawn from the point A, so as not to cut the circle, except AE.

**COROLLARY.** (i) *A tangent touches a circle at one point only.*

**COROLLARY.** (ii) *There can be but one tangent to a circle at a given point.*

PROPOSITION 17. PROBLEM.

To draw a tangent to a circle from a given point either on, or without the circumference.

Fig. 1.





Fig. 2.



Let BCD be the given circle, and A the given point.

It is required to draw from A a tangent to the  $\odot$  CDB.

CASE I. When the given point A is on the  $\odot$ .

Construction. Find E, the centre of the circle. III. 1.

Join EA

At A draw AK at rt. angles to EA. I. 11.

Proof. Then AK being perp to a diameter at one of its extremities, is a tangent to the circle. III. 16.

CASE II. When the given point A is without the  $\odot$ .

Construction. Find E, the centre of the circle; III. 1.  
and join AE, cutting the  $\odot$  BCD at D.

With centre E and radius EA, describe the  $\odot$  AFG.

At D, draw GDF at rt. angles to EA, cutting the  $\odot$  AFG at F and G. I. 11.

Join EF, EG, cutting the  $\odot$  BCD at B and C.

Join AB, AC.

Then both AB and AC shall be tangents to the  $\odot$  CDB.

Proof. In the  $\triangle$  AEB, FED,


Because {  $\begin{cases} AE = FE, \text{ being radii of the } \odot \text{ AFG;} \\ \text{and } EB = ED, \text{ being radii of the } \odot \text{ BDC;} \\ \text{and the included angle AEF is common;} \end{cases}$

$\therefore$  the  $\angle$  ABE = the  $\angle$  FDE.

I. 4.

H.S.E.

N



But the  $\angle FDE$  is a rt. angle ; *Constr.*

$\therefore$  the  $\angle ABE$  is a rt. angle.

Hence AB, being drawn at rt. angles to a diameter at one of its extremities, is a tangent to the  $\odot BCD$ . III. 16.

Similarly it may be shewn that AC is a tangent. Q.E.F.

**COROLLARY.** *If two tangents are drawn to a circle from an external point, then (i) they are equal ; (ii) they subtend equal angles at the centre ; (iii) they make equal angles with the straight line which joins the given point to the centre.*

For, in the above figure,

Since ED is perp. to FG, a chord of the  $\odot FAG$ , *Constr.*  
 $\therefore DF = DG$ . III. 3.


Then in the  $\triangle DEF, DEG$ ,  
 Because  $\left\{ \begin{array}{l} DE \text{ is common to both,} \\ \text{and } EF = EG ; \\ \text{and } DF = DG ; \end{array} \right.$  I. Def. 15.  
 $\therefore$  the  $\angle DEF =$  the  $\angle DEG$ . *Proved.*  
I. 8.

Again in the  $\triangle AEB, AEC$ ,  
 Because  $\left\{ \begin{array}{l} AE \text{ is common to both,} \\ \text{and } EB = EC, \\ \text{and the } \angle AEB = \text{the } \angle AEC ; \end{array} \right.$  *Proved.*  
 $\therefore AB = AC$  ; I. 4.  
 and the  $\angle EAB =$  the  $\angle EAC$ . Q. E. D.

**NOTE.** If the given point A is within the circle, no solution is possible. Hence we see that this problem admits of *two* solutions, *one* solution, or *no* solution, according as the given point A is *without*, *on*, or *within* the circumference of a circle. For a simpler method of drawing a tangent to a circle from a given point, see page 218.

PROPOSITION 18. THEOREM.

*The straight line drawn from the centre of a circle to the point of contact of a tangent is perpendicular to the tangent.*



Let ABC be a circle, of which F is the centre ;  
and let the st. line DE touch the circle at C.

*Then shall FC be perp. to DE.*

For, if not, suppose, if possible, FG to be perp. to DE, I. 12.  
and let FG meet the  $\bigcirc^\infty$  at B.

**Proof.**

In the  $\triangle FCG$ , because the  $\angle FGC$  is a rt. angle, *Hyp.*

$\therefore$  the  $\angle FCG$  is less than a rt. angle ; I. 17.

$\therefore$  the  $\angle FGC$  is greater than the  $\angle FCG$  ;

$\therefore$  FC is greater than FG ; I. 19.

but  $FC = FB$  ;

$\therefore$  FB is greater than FG,

the part greater than the whole, which is impossible.

$\therefore$  FC cannot be otherwise than perp. to DE :

that is, FC is perp. to DE.

Q.E.D.

EXERCISES.

1. Draw a tangent to a circle (i) parallel to, (ii) at right angles to a given straight line.

2. Tangents drawn to a circle from the extremities of a diameter are parallel.


3. Circles which touch one another internally or externally have a common tangent at their point of contact.

4. In two concentric circles, any chord of the outer circle which touches the inner, is bisected at the point of contact.

5. In two concentric circles, all chords of the outer circle which touch the inner, are equal.

## PROPOSITION 19. THEOREM.

*The straight line drawn perpendicular to a tangent to a circle from the point of contact passes through the centre.*



Let ABC be a circle, and DE a tangent to it at the point C; and let CA be drawn perp. to DE.

*Then shall CA pass through the centre.*

**Construction.** For if not, suppose, if possible, the centre F to be outside CA.

Join CF.

**Proof.** Because DE is a tangent to the circle, and FC is drawn from the centre F to the point of contact,

$\therefore$  the  $\angle FCE$  is a rt. angle. III. 18.

But the  $\angle ACE$  is a rt. angle; Hyp.

$\therefore$  the  $\angle FCE =$  the  $\angle ACE$ ;

the part equal to the whole, which is impossible.

$\therefore$  the centre cannot be otherwise than in CA;  
that is, CA passes through the centre.

Q.E.D.

## EXERCISES ON THE TANGENT.

## PROPOSITIONS 16, 17, 18, 19.

1. *The centre of any circle which touches two intersecting straight lines must lie on the bisector of the angle between them.*

2. *AB and AC are two tangents to a circle whose centre is O; shew that AO bisects the chord of contact BC at right angles.*

3. If two circles are concentric all tangents drawn from points on the circumference of the outer to the inner circle are equal.

4. The diameter of a circle bisects all chords which are parallel to the tangent at either extremity.

5. Find the locus of the centres of all circles which touch a given straight line at a given point.

6. Find the locus of the centres of all circles which touch each of two parallel straight lines.

7. Find the locus of the centres of all circles which touch each of two intersecting straight lines of unlimited length.

8. Describe a circle of given radius to touch two given straight lines.

9. Through a given point, within or without a circle, draw a chord equal to a given straight line.

In order that the problem may be possible, between what limits must the given line lie, when the given point is (i) without the circle, (ii) within it?

10. Two parallel tangents to a circle intercept on any third tangent a segment which subtends a right angle at the centre.

11. In any quadrilateral circumscribed about a circle, the sum of one pair of opposite sides is equal to the sum of the other pair.

12. Any parallelogram which can be circumscribed about a circle, must be equilateral.

13. If a quadrilateral is described about a circle, the angles subtended at the centre by any two opposite sides are together equal to two right angles.

14. AB is any chord of a circle, AC the diameter through A, and AD the perpendicular on the tangent at B: shew that AB bisects the angle DAC.

15. Find the locus of the extremities of tangents of fixed length drawn to a given circle.

16. In the diameter of a circle produced, determine a point such that the tangent drawn from it shall be of given length.

17. In the diameter of a circle produced, determine a point such that the two tangents drawn from it may contain a given angle.

18. Describe a circle that shall pass through a given point, and touch a given straight line at a given point. [See page 197. Ex. 5.]

19. Describe a circle of given radius, having its centre on a given straight line, and touching another given straight line.

20. Describe a circle that shall have a given radius, and touch a given circle and a given straight line. How many such circles can be drawn?



## PROPOSITION 20. THEOREM.

*The angle at the centre of a circle is double of an angle at the circumference, standing on the same arc.*

Fig. 1.





Fig. 2.



Let  $ABC$  be a circle, of which  $E$  is the centre; and let  $BEC$  be the angle at the centre, and  $BAC$  an angle at the  $\circ^{\circ}$ , standing on the same arc  $BC$ .

*Then shall the  $\angle BEC$  be double of the  $\angle BAC$ .*

**Construction.** Join  $AE$ , and produce it to  $F$ .

**CASE I.** When the centre  $E$  is within the angle  $BAC$ .

**Proof.** In the  $\triangle EAB$ , because  $EA = EB$ , I. Def. 15.

$\therefore$  the  $\angle EAB =$  the  $\angle EBA$ ; I. 5.

$\therefore$  the sum of the  $\angle^s EAB, EBA =$  twice the  $\angle EAB$ .

But the ext.  $\angle BEF =$  the sum of the  $\angle^s EAB, EBA$ ; I. 32.

$\therefore$  the  $\angle BEF =$  twice the  $\angle EAB$ .

Similarly the  $\angle FEC =$  twice the  $\angle EAC$ .

$\therefore$  the sum of the  $\angle^s BEF, FEC =$  twice the sum of  
the  $\angle^s EAB, EAC$ ;

that is, the  $\angle BEC =$  twice the  $\angle BAC$ .

**CASE II.** When the centre  $E$  is without the  $\angle BAC$ .

As before, it may be shewn that the  $\angle FEB =$  twice the  $\angle FAB$ ;


also the  $\angle FEC =$  twice the  $\angle FAC$ ;

$\therefore$  the difference of the  $\angle^s FEC, FEB =$  twice the difference  
of the  $\angle^s FAC, FAB$ ;

that is, the  $\angle BEC =$  twice the  $\angle BAC$ . Q.E.D.


NOTE 1. The case in which the centre E falls on AB or AC needs no proof beyond that given under Case I.

NOTE 2. If the arc BFC, on which the angles stand, is greater than a semi-circumference, the angle BEC at the centre will be reflex: but it may still be shewn as, in Case I., that the reflex  $\angle BEC$  is double of the  $\angle BAC$  at the  $\odot^\infty$ , standing on the same arc BFC.



PROPOSITION 21. THEOREM.

*Angles in the same segment of a circle are equal.*



Let ABCD be a circle, and let BAD, BED be angles in the same segment BAED.

*Then shall the  $\angle BAD =$  the  $\angle BED$ .*

Construction. Find F, the centre of the circle. III. 1.

CASE I. When the segment BAED is greater than a semicircle.

Join BF, DF.


Proof. Because the  $\angle BFD$  is at the centre, and the  $\angle BAD$  at the  $\odot^\infty$ , standing on the same arc BD,

$\therefore$  the  $\angle BFD =$  twice the  $\angle BAD$ . III. 20.

Similarly the  $\angle BFD =$  twice the  $\angle BED$ . III. 20.

$\therefore$  the  $\angle BAD =$  the  $\angle BED$ . Ax. 7.

CASE II. When the segment BAED is not greater than a semicircle.



**Construction.**

Join AF, and produce it to meet the  $\bigcirc^{\infty}$  at C.

Join EC.

**Proof.** Then since AEDC is a semicircle ;

$\therefore$  the segment BAEC is greater than a semicircle :

$\therefore$  the  $\angle$  BAC = the  $\angle$  BEC, in this segment. *Case 1*

Similarly the segment CAED is greater than a semicircle ;

$\therefore$  the  $\angle$  CAD = the  $\angle$  CED, in this segment.

$\therefore$  the  $\angle$  BAC, CAD = the sum of the  $\angle$  BEC, CED .

that is, the  $\angle$  BAD = the  $\angle$  BED.

Q.E.D.

#### EXERCISES.

1. P is any point on the arc of a segment of which AB is the chord. Shew that the sum of the angles PAB, PBA is constant.

2. PQ and RS are two chords of a circle intersecting at X : prove that the triangles PXS, RXQ are equiangular.

3. Two circles intersect at A and B ; and through A any straight line PAQ is drawn terminated by the circumferences : shew that PQ subtends a constant angle at B.

4. Two circles intersect at A and B ; and through A any two straight lines PAQ, XAY are drawn terminated by the circumferences ; shew that the arcs PX, QY subtend equal angles at B.

5. P is any point on the arc of a segment whose chord is AB : and the angles PAB, PBA are bisected by straight lines which intersect at O. Find the locus of the point O.


NOTE. If the extension of Proposition 20, given in Note 2 on page 199, is adopted, a separate treatment of the second case of the present proposition is unnecessary.

For, as in Case I.,

the reflex  $\angle BFD =$  twice the  $\angle BAD$  ;  
 III. 20.

also the reflex  $\angle BFD =$  twice the  $\angle BED$  ;

$\therefore$  the  $\angle BAD =$  the  $\angle BED$ .



Obs. The converse of Prop. 21 is important. For the construction used, viz. To describe a circle about a given triangle, see Book iv., Prop. 5, or Theorems and Examples on Book I, page 111, No. 1.

CONVERSE OF PROPOSITION 21.

Equal angles standing on the same base, and on the same side of it, have their vertices on an arc of a circle, of which the given base is the chord.

Let  $\angle BAC, \angle BDC$  be two equal angles standing on the same base  $BC$ .

Then shall the vertices  $A$  and  $D$  lie upon a segment of a circle having  $BC$  as its chord.

Describe a circle about the  $\triangle BAC$ . iv. 5.

Then this circle shall pass through  $D$ .

For, if not, it must cut  $BD$ , or  $BD$  produced, at some other point  $E$ .

Join  $EC$ .

Then the  $\angle BAC =$  the  $\angle BEC$ , in the same segment : III. 21.

but the  $\angle BAC =$  the  $\angle BDC$ , by hypothesis ;

$\therefore$  the  $\angle BEC =$  the  $\angle BDC$  ;


that is, an ext. angle of a triangle = an int. opp. angle ;

which is impossible. I. 16.

$\therefore$  the circle which passes through  $B, A, C$ , cannot pass otherwise than through  $D$ .

That is, the vertices  $A$  and  $D$  are on an arc of a circle of which the chord is  $BC$ .


Q. E. D.



COROLLARY. The locus of the vertices of triangles drawn on the same base and on the same side of it with equal vertical angles is an arc of a circle.

## PROPOSITION 22. THEOREM.

*The opposite angles of any quadrilateral inscribed in a circle are together equal to two right angles.*



Let ABCD be a quadrilateral inscribed in the  $\odot$  ABC.

Then shall (i) the  $\angle^s$  ADC, ABC together = two rt. angles ;  
 (ii) the  $\angle^s$  BAD, BCD together = two rt. angles.

Construction. Join AC, BD.

Proof.

Since the  $\angle$  ADB = the  $\angle$  ACB, in the segment ADCB; III. 21.  
 and the  $\angle$  CDB = the  $\angle$  CAB, in the segment CDAB;  
 $\therefore$  the  $\angle$  ADC = the sum of the  $\angle^s$  ACB, CAB.

To each of these equals add the  $\angle$  ABC :  
 then the two  $\angle^s$  ADC, ABC together = the three  $\angle^s$  ACB,  
 CAB, ABC.

But the  $\angle^s$  ACB, CAB, ABC, being the angles of a triangle,  
 together = two rt. angles ; I. 32.

$\therefore$  the  $\angle^s$  ADC, ABC together = two rt. angles.

Similarly it may be shewn that

the  $\angle^s$  BAD, BCD together = two rt. angles. Q.E.D.

## EXERCISES.

1. If a circle can be described about a parallelogram, the parallelogram must be rectangular.
2. ABC is an isosceles triangle, and XY is drawn parallel to the base BC cutting the sides in X and Y : shew that the four points B, C, X, Y lie on a circle.
3. If one side of a quadrilateral inscribed in a circle is produced, the exterior angle is equal to the opposite interior angle of the quadrilateral.

PROPOSITION 22. [ALTERNATIVE PROOF.]

Let ABCD be a quadrilateral inscribed in the  $\odot$  ABC.

Then shall the  $\angle^s$  ADC, ABC together = two rt. angles.

Join FA, FC.


Since the  $\angle$  AFC at the centre = twice the  $\angle$  ADC at the  $\odot^\infty$ , standing on the same arc ABC; III. 20.

and the reflex angle AFC at the centre = twice the  $\angle$  ABC at the  $\odot^\infty$ , standing on the same arc ADC; III. 20.

$\therefore$  the  $\angle^s$  ADC, ABC are together half the sum of the  $\angle$  AFC and the reflex angle AFC;

but these make up four rt. angles: I. 15. Cor. 2.

$\therefore$  the  $\angle^s$  ADC, ABC together = two rt. angles. Q.E.D.



DEFINITION. Four or more points through which a circle may be described are said to be **concylic**.

CONVERSE OF PROPOSITION 22.

If a pair of opposite angles of a quadrilateral are together equal to two right angles, its vertices are concyclic.

Let ABCD be a quadrilateral, in which the opposite angles at B and D together = two rt. angles.

Then shall the four points A, B, C, D be concyclic.

Through the three points A, B, C describe a circle. IV. 5.

Then this circle must pass through D.

For, if not, it will cut AD, or AD produced, at some other point E.

Join EC.

Then since the quadrilateral ABCE is inscribed in a circle,

$\therefore$  the  $\angle^s$  ABC, AEC together = two rt. angles. III. 22.

But the  $\angle^s$  ABC, ADC together = two rt. angles; Hyp.

hence the  $\angle^s$  ABC, AEC = the  $\angle^s$  ABC, ADC.

Take from these equals the  $\angle$  ABC;


then the  $\angle$  AEC = the  $\angle$  ADC;

that is, an ext. angle of a triangle = an int. opp. angle;

which is impossible. I. 16.

$\therefore$  the circle which passes through A, B, C cannot pass otherwise than through D:


that is the four vertices A, B, C, D are concyclic. Q.E.D.



**DEFINITION.** **Similar segments** of circles are those which contain equal angles. [Book III., Def. 10.]

**PROPOSITION 23. THEOREM.**

*On the same chord and on the same side of it, there cannot be two similar segments of circles, not coinciding with one another.*



If possible, on the same chord  $AB$ , and on the same side of it, let there be two similar segments of circles  $ACB$ ,  $ADB$ , not coinciding with one another.

Then since the arcs  $ADB$ ,  $ACB$  intersect at  $A$  and  $B$ ,  
 $\therefore$  they cannot cut one another again; III. 10.  
 $\therefore$  one segment falls within the other.

**Construction.** In the inner arc take any point  $C$ .

Join  $AC$ , producing it to meet the outer arc at  $D$  :  
 join  $CB$ ,  $DB$ .

**Proof.** Then because the segments are similar,  
 $\therefore$  the  $\angle ACB =$  the  $\angle ADB$ ; III. Def. 10.  
 that is, an ext. angle of the  $\triangle CDB =$  an int. opp. angle;  
 which is impossible. I. 16.

Hence the two similar segments  $ACB$ ,  $ADB$ , on the same chord  $AB$  and on the same side of it, must coincide. Q.E.D.

**EXERCISES ON PROPOSITION 22.**


1. The straight lines which bisect any angle of a quadrilateral figure inscribed in a circle and the opposite exterior angle, meet on the circumference.

2. A triangle is inscribed in a circle: shew that the sum of the angles in the three segments exterior to the triangle is equal to four right angles.

3. Divide a circle into two segments, so that the angle contained by the one shall be double of the angle contained by the other.

PROPOSITION 24. THEOREM.

*Similar segments of circles on equal chords are equal to one another.*



Let AEB and CFD be similar segments on equal chords AB, CD.

*Then shall the segment AEB = the segment CFD.*

**Proof.** If the segment AEB be applied to the segment CFD, so that A falls on C, and AB falls along CD ;

then since  $AB = CD$ ,

*Hyp.*

$\therefore$  B must coincide with D.

$\therefore$  the segment AEB must coincide with the segment CFD ; for if not, on the same chord and on the same side of it there would be two similar segments of circles, not coinciding with one another : which is impossible. III. 23.

$\therefore$  the segment AEB = the segment CFD. Q.E.D.

EXERCISES.

1. Of two segments standing on the same chord, the greater segment contains the smaller angle.

2. A segment of a circle stands on a chord AB, and P is any point on the same side of AB as the segment : shew that the angle APB is greater or less than the angle in the segment, according as P is within or without the segment.

3. P, Q, R are the middle points of the sides of a triangle, and X is the foot of the perpendicular let fall from one vertex on the opposite side : shew that the four points P, Q, R, X are concyclic.


[See page 104, Ex. 2 : also page 108, Ex. 2.]

4. Use the preceding exercise to shew that the middle points of the sides of a triangle and the feet of the perpendiculars let fall from the vertices on the opposite sides, are concyclic.



## PROPOSITION 25. PROBLEM.

*An arc of a circle being given, to describe the whole circumference of which the given arc is a part.*



Let  $ABC$  be an arc of a circle.

*It is required to describe the whole  $\circ^{\infty}$  of which the arc  $ABC$  is a part.*

**Construction.**

In the given arc take any three points  $A, B, C$ .

Join  $AB, BC$ .

Draw  $DF$  bisecting  $AB$  at rt. angles, i. 10. 11.  
and draw  $EF$  bisecting  $BC$  at rt. angles.

**Proof.**

Then because  $DF$  bisects the chord  $AB$  at rt. angles,

$\therefore$  the centre of the circle lies in  $DF$ . III. 1 *Cor.*

Again, because  $EF$  bisects the chord  $BC$  at rt. angles,

$\therefore$  the centre of the circle lies in  $EF$ . III. 1 *Cor.*


$\therefore$  the centre of the circle is  $F$ , the only point common to  $DF$  and  $EF$ .

Hence the  $\circ^{\infty}$  of a circle described with centre  $F$ , and radius  $FA$ , is that of which the given arc is a part. Q.E.F.

NOTE. Euclid gave to this proposition a somewhat different form, as follows :

PROPOSITION 25. [EUCLID'S METHOD.]

*A segment of a circle being given, to describe the circle of which it is a segment.*



Let ABC be the given segment of a circle, standing on the chord AC.

*It is required to describe the circle of which ABC is a segment.*

**Construction.** Draw DB, bisecting AC at rt. angles, and meeting the  $\text{O}^\infty$  at B.

Join AB.

CASE I. When the  $\angle DAB$  is not equal to the  $\angle ABD$ .

At A, in BA, make the  $\angle BAE$  equal to the  $\angle ABD$ ; I. 23.

and let AE meet BD, or BD produced, at E.

Join EC.

*Then E shall be the centre of the required circle.*

**Proof.** Since the  $\angle EAB = \text{the } \angle EBA$ , Constr.

$\therefore EA = EB$ . I. 6.

And in the  $\triangle^s EDA, EDC$ ,

$DA = DC$ , Constr.

and ED is common ;

Because { also the  $\angle EDA = \text{the } \angle EDC$ , being rt. angles ;

$\therefore EA = EC$ . I. 4.


Hence EA, EB, and EC are all equal ;

$\therefore E$  is the centre of the required circle, and EA, EB, EC are radii.

CASE II. When the  $\angle DAB = \text{the } \angle ABD$ .


In this case it follows that  $DB = DA$ ; I. 6.

$\therefore DB, DA, DC$  are all equal, so that D is the centre of the required circle.



## PROPOSITION 26. THEOREM.

*In equal circles the arcs which subtend equal angles, whether at the centres or at the circumferences, shall be equal.*



Let  $ABC, DEF$  be equal circles ;  
and let the  $\angle^s BGC, EHF$  at the centres be equal,  
and consequently the  $\angle^s BAC, EDF$  at the  $\odot^{\text{ces}}$  equal. III. 20.

*Then shall the arc  $BKC =$  the arc  $ELF$ .*

**Construction.** Join  $BC, EF$ .

**Proof.** Because the  $\odot^s ABC, DEF$  are equal,  
 $\therefore$  their radii are equal.

Hence in the  $\triangle^s BGC, EHF,$

Because  $\left\{ \begin{array}{l} BG = EH, \\ \text{and } GC = HF, \\ \text{and the } \angle BGC = \text{the } \angle EHF ; \\ \therefore BC = EF. \end{array} \right.$

*Hyp.*  
I. 4.

Again, because the  $\angle BAC =$  the  $\angle EDF,$  *Hyp.*  
 $\therefore$  the segment  $BAC$  is similar to the segment  $EDF$  ;  
III. Def. 10.

and these segments are on equal chords  $BC, EF$  ;  
 $\therefore$  the segment  $BAC =$  the segment  $EDF$ . III. 24.


But the whole  $\odot ABC =$  the whole  $\odot DEF$  ;  
 $\therefore$  the remaining segment  $BKC =$  the remaining segment  $ELF$  ;  
 $\therefore$  the arc  $BKC =$  the arc  $ELF$ .

Q.E.D.

[For Exercises and an Alternative Proof see pp. 212, 213.]

PROPOSITION 27. THEOREM.

*In equal circles the angles, whether at the centres or the circumferences, which stand on equal arcs, shall be equal.*



Let  $ABC, DEF$  be equal circles ;  
and let the arc  $BC =$  the arc  $EF$ .

*Then shall the  $\angle BGC =$  the  $\angle EHF$ , at the centres ;  
and also the  $\angle BAC =$  the  $\angle EDF$ , at the  $\odot^{ces}$ .*

**Construction.** If the  $\angle^s BGC, EHF$  are not equal, one must be the greater.

If possible, let the  $\angle BGC$  be the greater.

At  $G$ , in  $BG$ , make the  $\angle BGK$  equal to the  $\angle EHF$ . I. 23.

**Proof.** In the equal  $\odot^s ABC, DEF$ ,  
because the  $\angle BGK =$  the  $\angle EHF$ , at the centres ; *Constr.*  
 $\therefore$  the arc  $BK =$  the arc  $EF$ . III. 26.

But the arc  $BC =$  the arc  $EF$  ; *Hyp.*  
 $\therefore$  the arc  $BK =$  the arc  $BC$ ,  
a part equal to the whole, which is impossible.

$\therefore$  the  $\angle BGC$  is not unequal to the  $\angle EHF$  ;  
that is, the  $\angle BGC =$  the  $\angle EHF$ .


And since the  $\angle BAC$  at the  $\odot^{ce}$  is half the  $\angle BGC$  at the centre, III. 20.

and likewise the  $\angle EDF$  is half the  $\angle EHF$ ,  
 $\therefore$  the  $\angle BAC =$  the  $\angle EDF$ . *Ax. 7.*

**Q.E.D.**

## PROPOSITION 28. THEOREM.

*In equal circles the arcs, which are cut off by equal chords, shall be equal, the major arc equal to the major arc, and the minor to the minor.*



Let  $ABC, DEF$  be equal circles ;  
and let the chord  $BC =$  the chord  $EF$ .

*Then shall the major arc  $BAC =$  the major arc  $EDF$  ;  
and the minor arc  $BGC =$  the minor arc  $EHF$ .*

**Construction.**

Find  $K$  and  $L$  the centres of the  $\odot^s ABC, DEF$  ; III. 1.  
and join  $BK, KC, EL, LF$ .

**Proof.** Because the  $\odot^s ABC, DEF$  are equal,  
 $\therefore$  their radii are equal.

Hence in the  $\triangle^s BKC, ELF$ ,

Because  $\left\{ \begin{array}{l} BK = EL, \\ KC = LF, \\ \text{and } BC = EF; \end{array} \right.$

$\therefore$  the  $\angle BKC =$  the  $\angle ELF$ .

*Hyp.*  
I. 8.

$\therefore$  the arc  $BGC =$  the arc  $EHF$  ;


for these arcs subtend equal angles at the centre ; III. 26.  
and they are the minor arcs.

But the whole  $\circ^\circ ABGC =$  the whole  $\circ^\circ DEHF$  ; *Hyp.*  
 $\therefore$  the remaining arc  $BAC =$  the remaining arc  $EDF$  ;  
and these are the major arcs. Q.E.D.

[For Exercises see p. 212.]

PROPOSITION 29. THEOREM.

*In equal circles the chords, which cut off equal arcs, shall be equal.*



Let ABC, DEF be equal circles ;  
and let the arc BGC = the arc EHF.  
Then shall the chord BC = the chord EF.

**Construction.** Find K, L the centres of the circles.  
Join BK, KC, EL, LF.

**Proof.** In the equal  $\odot^s$  ABC, DEF,  
because the arc BGC = the arc EHF,  
 $\therefore$  the  $\angle$  BKC = the  $\angle$  ELF, at the centres. III. 27.

Hence in the  $\triangle^s$  BKC, ELF,  
BK = EL, being radii of equal circles ;  
KC = LF, for the same reason,  
and the  $\angle$  BKC = the  $\angle$  ELF ;  
 $\therefore$  BC = EF. *Proved.*  
I. 4.  
Q.E.D.

EXERCISES.

ON PROPOSITIONS 26, 27.

1. *If two chords of a circle are parallel, they intercept equal arcs.*
2. *The straight lines, which join the extremities of two equal arcs of a circle towards the same parts, are parallel.*
3. *In a circle, or in equal circles, sectors are equal if their angles at the centres are equal.*

4. If two chords of a circle intersect at right angles, the opposite arcs are together equal to a semi-circumference.

5. If two chords intersect within a circle, they form an angle equal to that subtended at the circumference by the sum of the arcs they cut off.

6. If two chords intersect without a circle, they form an angle equal to that subtended at the circumference by the difference of the arcs they cut off.

7. If  $AB$  is a fixed chord of a circle, and  $P$  any point on one of the arcs cut off by it, then the bisector of the angle  $APB$  cuts the conjugate arc in the same point, whatever be the position of  $P$ .

8. Two circles intersect at  $A$  and  $B$ ; and through these points straight lines are drawn from any point  $P$  on the circumference of one of the circles: shew that when produced they intercept on the other circumference an arc which is constant for all positions of  $P$ .

9. A triangle  $ABC$  is inscribed in a circle, and the bisectors of the angles meet the circumference at  $X, Y, Z$ . Find each angle of the triangle  $XYZ$  in terms of those of the original triangle.

#### ON PROPOSITIONS 28, 29.

10. The straight lines which join the extremities of parallel chords in a circle (i) towards the same parts, (ii) towards opposite parts, are equal.

11. Through  $A$ , a point of intersection of two equal circles, two straight lines  $PAQ, XAY$  are drawn: shew that the chord  $PX$  is equal to the chord  $QY$ .

12. Through the points of intersection of two circles two parallel straight lines are drawn terminated by the circumferences: shew that the straight lines which join their extremities towards the same parts are equal.

13. Two equal circles intersect at  $A$  and  $B$ ; and through  $A$  any straight line  $PAQ$  is drawn terminated by the circumferences: shew that  $BP = BQ$ .

14.  $ABC$  is an isosceles triangle inscribed in a circle, and the bisectors of the base angles meet the circumference at  $X$  and  $Y$ . Shew that the figure  $BXAYC$  must have four of its sides equal.


What relation must subsist among the angles of the triangle  $ABC$ , in order that the figure  $BXAYC$  may be equilateral?

NOTE. We have given Euclid's demonstrations of Propositions 26, 27; but it should be noticed that these propositions also admit of proof by the method of *superposition*.

To illustrate this method we will apply it to Proposition 26.

PROPOSITION 26. [ALTERNATIVE PROOF.]

*In equal circles, the arcs which subtend equal angles, whether at the centres or circumferences, shall be equal.*



Let  $ABC, DEF$  be equal circles ;  
 and let the  $\angle^s BGC, EHF$  at the centres be equal,  
 and consequently the  $\angle^s BAC, EDF$  at the  $\circ^{ces}$  equal. III. 20.  
*Then shall the arc  $BKC =$  the arc  $ELF$ .*

**Proof.** For if the  $\odot ABC$  be applied to the  $\odot DEF$ , so that the centre  $G$  may fall on the centre  $H$ ,

then because the circles are equal, *Hyp.*

$\therefore$  their  $\circ^{ces}$  must coincide ;

hence by revolving the upper circle about its centre, the lower circle remaining fixed,

$B$  may be made to coincide with  $E$ ,  
 and consequently  $GB$  with  $HE$ .

And because the  $\angle BGC =$  the  $\angle EHF$ , *Hyp.*

$\therefore GC$  must coincide with  $HF$  :

and since  $GC = HF$ , *Hyp.*

$\therefore C$  must fall on  $F$ .

Now  $B$  coincides with  $E$ , and  $C$  with  $F$ ,  
 and the  $\circ^{ce}$  of the  $\odot ABC$  with the  $\circ^{ce}$  of the  $\odot DEF$  ;

$\therefore$  the arc  $BKC$  must coincide with the arc  $ELF$ .


$\therefore$  the arc  $BKC =$  the arc  $ELF$ .

Q. E. D.



## PROPOSITION 30. PROBLEM.

To bisect a given arc.



Let ADB be the given arc.

It is required to bisect the arc ADB.

**Construction.** Join AB; and bisect AB at C. I. 10.  
At C draw CD at rt. angles to AB, meeting the given arc at D. I. 11.

Then shall the arc ADB be bisected at D.  
Join AD, BD.

**Proof.** In the  $\triangle$  ACD, BCD,  
Because  $\left\{ \begin{array}{l} AC = BC, \\ \text{and } CD \text{ is common;} \\ \text{and the } \angle ACD = \text{the } \angle BCD, \text{ being rt. angles;} \end{array} \right.$  *Constr.*  
 $\therefore AD = BD.$  I. 4.

And since in the  $\odot$  ADB, the chords AD, BD are equal,  
 $\therefore$  the arcs cut off by them are equal, the minor arc equal to the minor, and the major arc to the major: III. 28.  
and the arcs AD, BD are both minor arcs,  
for each is less than a semi-circumference, since DC, bisecting the chord AB at rt. angles, must pass through the centre of the circle. III. 1. *Cor.*

$\therefore$  the arc AD = the arc BD :  
that is, the arc ADB is bisected at D. Q.E.F.

## EXERCISES.

1. If a tangent to a circle is parallel to a chord, the point of contact will bisect the arc cut off by the chord.
2. Trisect a quadrant, or the fourth part of the circumference, of a circle.

**NOTE.** The following alternative proof of Proposition 30 removes the necessity of distinguishing between the major and minor arcs cut off by the chords AD, BD.

PROPOSITION 30. [ALTERNATIVE PROOF.]

The construction being made as before, we may proceed thus :

Proof.	In the $\triangle^s$ ACD, BCD,	
Because {	AC = BC,	<i>Constr.</i>
	and CD is common ;	
	and the $\angle$ ACD = the $\angle$ BCD, being rt. angles :	
	$\therefore$ the $\angle$ DAC = the $\angle$ DBC :	I. 4.
	that is, the $\angle$ DAB = the $\angle$ DBA.	

But these are angles at the  $\circ^e$  subtended by the arcs DB, DA ;

$\therefore$  the arc DB = the arc DA : III. 26.  
that is, the arc ADB is bisected at D. Q.E.F.

QUESTIONS FOR REVISION.


1. When is a straight line said (i) *to meet*, (ii) *to cut*, (iii) *to touch*, the circumference of a circle?
2. When are circles said *to touch* one another? Distinguish between *internal* and *external* contact.
3. What theorems have been so far proved by Euclid regarding (i) *circles which cut one another*, (ii) *circles which touch one another*?
4. If two unequal circles are concentric, shew that one must lie wholly within the other.
5. Shew how to divide the circumference of a circle into *three*, *four*, or *six* equal parts.
6. Enunciate the propositions so far proved by Euclid relating to the properties of a *tangent to a circle*.

## PROPOSITION 31. THEOREM.

*The angle in a semicircle is a right angle.*

*The angle in a segment greater than a semicircle is less than a right angle.*

*The angle in a segment less than a semicircle is greater than a right angle.*



Let ABCD be a circle, of which BC is a diameter, and E the centre; and let AC be a chord dividing the circle into the segments ABC, ADC, of which the segment ABC is greater, and the segment ADC is less than a semicircle.

Then (i) the angle in the semicircle BAC shall be a right angle;

(ii) the angle in the segment ABC shall be less than a rt. angle;

(iii) the angle in the segment ADC shall be greater than a rt. angle.

**Construction.** In the arc ADC take any point D;  
Join BA, AD, DC, AE; and produce BA to F.

**Proof.** (i) Because EA = EB, I. Def. 15.

$\therefore$  the  $\angle$  EAB = the  $\angle$  EBA. I. 5.

And because EA = EC,

$\therefore$  the  $\angle$  EAC = the  $\angle$  ECA.

$\therefore$  the whole  $\angle$  BAC = the sum of the  $\angle$ 's EBA, ECA;

but the ext.  $\angle$  FAC = the sum of the two int.  $\angle$ 's CBA, BCA;

$\therefore$  the  $\angle$  BAC = the  $\angle$  FAC;

$\therefore$  these angles, being adjacent, are rt. angles.

$\therefore$  the  $\angle$  BAC, in the semicircle BAC, is a rt. angle.

(ii) In the  $\triangle ABC$ , because the sum of the  $\angle^s ABC, BAC$  is less than two rt. angles ; I. 17.

and of these, the  $\angle BAC$  is a rt. angle ; *Proved.*

$\therefore$  the  $\angle ABC$ , which is the angle in the segment  $ABC$ , is less than a rt. angle.

(iii) Because  $ABCD$  is a quadrilateral inscribed in the  $\odot ABC$ ,

$\therefore$  the opp.  $\angle^s ABC, ADC$  together = two rt. angles ; III. 22.  
and of these, the  $\angle ABC$  is less than a rt. angle : *Proved.*

$\therefore$  the  $\angle ADC$ , which is the angle in the segment  $ADC$ , is greater than a rt. angle. Q.E.D.

## EXERCISES.

1. A circle described on the hypotenuse of a right-angled triangle as diameter, passes through the opposite angular point.

2. A system of right-angled triangles is described upon a given straight line as hypotenuse ; find the locus of the opposite angular points.

3. A straight rod of given length slides between two straight rulers placed at right angles to one another ; find the locus of its middle point.

4. Two circles intersect at  $A$  and  $B$  ; and through  $A$  two diameters  $AP, AQ$  are drawn, one in each circle : shew that the points  $P, B, Q$  are collinear. [See Def. p. 110.]

5. A circle is described on one of the equal sides of an isosceles triangle as diameter. Shew that it passes through the middle point of the base.

6. Of two circles which have internal contact, the diameter of the inner is equal to the radius of the outer. Shew that any chord of the outer circle, drawn from the point of contact, is bisected by the circumference of the inner circle.

7. Circles described on any two sides of a triangle as diameters intersect on the third side, or the third side produced.

8. Find the locus of the middle points of chords of a circle drawn through a fixed point. Distinguish between the cases when the given point is within, on, or without the circumference.

9. Describe a square equal to the difference of two given squares.

10. Through one of the points of intersection of two circles draw a chord of one circle which shall be bisected by the other.


11. On a given straight line as base a system of equilateral four-sided figures is described : find the locus of the intersection of their diagonals.

## NOTES ON PROPOSITION 31.

NOTE 1. The extension of Proposition 20 to *straight and reflex* angles furnishes a simple alternative proof of the first theorem contained in Proposition 31, namely,

*The angle in a semicircle is a right angle.*

For, in the adjoining figure, the angle at the centre, standing on the arc BHC, is double the angle BAC at the  $\odot^e$ , standing on the same arc.



Now the angle at the centre is the *straight angle* BEC ;  
 $\therefore$  the  $\angle$  BAC is half of the *straight angle* BEC :  
 and a straight angle = two rt. angles ;  
 $\therefore$  the  $\angle$  BAC = one half of two rt. angles,  
 = one rt. angle.

Q. E. D.

NOTE 2. From Proposition 31 we may derive a simple practical solution of Proposition 17, namely,

*To draw a tangent to a circle from a given external point.*

Let BCD be the given circle, and A the given external point.

*It is required to draw from A a tangent to the  $\odot$  BCD.*

Find E, the centre of the given circle, and join AE.

On AE describe the semicircle ABE, to cut the given circle at B.

Join AB.

*Then AB shall be a tangent to the  $\odot$  BCD.*

For the  $\angle$  ABE, being in a semicircle, is a rt. angle. III. 31.


$\therefore$  AB is drawn at rt. angles to the radius EB, from its extremity B ;

$\therefore$  AB is a tangent to the circle.

III. 16.

Q. E. F.

Since the semicircle might be described on either side of AE, it is clear that there will be a second solution of the problem, as shewn by the dotted lines of the figure.




## QUESTIONS FOR REVISION AND NUMERICAL EXERCISES.

1. Define an *arc*, a *chord*, a *segment* of a circle. When are segments of circles said to be *similar* to one another ?
2. Enunciate propositions which give the properties of *chords* of a circle in relation to the *centre*.
3. Prove that in a circle whose diameter is 34 inches, a chord 30 inches in length is at a distance of 8 inches from the centre.
4. In a circle a chord 2 feet in length stands at a distance of 5 inches from the centre : shew that the diameter of the circle is 2 inches longer than the chord.
5. What must be the length of a chord which is 1 foot distant from the centre of a circle, if the diameter is 2 yards 2 inches ?
6. Two parallel chords of a circle, whose diameter is 13 inches, are respectively 5 inches and 1 foot in length : shew that the distance between them is  $8\frac{1}{2}$  inches, or  $3\frac{1}{2}$  inches.
7. Two circles, whose radii are respectively 26 inches and 25 inches, intersect at two points which are 4 feet apart. Shew that the distance between their centres is 17 inches.
8. The diameters of two concentric circles are respectively 50 inches and 48 inches : shew that any chord of the outer circle which touches the inner must be 14 inches in length.
9. Of two concentric circles the diameter of the greater is 74 inches, and any chord of it which touches the smaller circle is 70 inches in length : shew that the diameter of the smaller circle is 2 feet.
10. Two circles of diameters 74 and 40 inches respectively have a common chord 2 feet in length : shew that the distance between their centres is 51 inches.
11. The chord of an arc is 24 inches in length, and the height of the arc is 8 inches ; shew that the diameter of the circle is 26 inches.
12. AB is a line 20 inches in length, and C is its middle point. On AB, AC, CB semicircles are described. Shew that if a circle is inscribed in the space enclosed by the three semicircles its radius must be  $3\frac{1}{2}$  inches.

## PROPOSITION 32. THEOREM.

If a straight line touches a circle, and from the point of contact a chord is drawn, the angles which this chord makes with the tangent shall be equal to the angles in the alternate segments of the circle.



Let  $EF$  touch the given  $\odot ABC$  at  $B$ , and let  $BD$  be a chord drawn from  $B$ , the point of contact.

Then shall

- (i) the  $\angle DBF =$  the angle in the alternate segment  $BAD$  :
- (ii) the  $\angle DBE =$  the angle in the alternate segment  $BCD$ .

**Construction.** From  $B$  draw  $BA$  perp. to  $EF$ . I. 11.  
 Take any point  $C$  in the arc  $BD$  ;  
 and join  $AD, DC, CB$ .

(i) **Proof.** Because  $BA$  is drawn perp. to the tangent  $EF$ , at its point of contact  $B$ ,

$\therefore BA$  passes through the centre of the circle : III. 19.

$\therefore$  the  $\angle ADB$ , being in a semicircle, is a rt. angle : III. 31.

$\therefore$  in the  $\triangle ABD$ , the other  $\angle^s ABD, BAD$  together = a rt. angle ; I. 32.

that is, the  $\angle^s ABD, BAD$  together = the  $\angle ABF$ .

From these equals take the common  $\angle ABD$  ;

$\therefore$  the  $\angle DBF =$  the  $\angle BAD$ , which is in the alternate segment.

(ii) Because ABCD is a quadrilateral inscribed in a circle,

$\therefore$  the opp.  $\angle^s$  BCD, BAD together = two rt. angles : III. 22.

but the  $\angle^s$  DBE, DBF together = two rt. angles ; I. 13.

$\therefore$  the  $\angle^s$  DBE, DBF together = the  $\angle^s$  BCD, BAD ;

and of these the  $\angle$  DBF = the  $\angle$  BAD ; *Proved.*

$\therefore$  the  $\angle$  DBE = the  $\angle$  BCD, which is in the alternate segment.

Q.E.D.

#### EXERCISES.

1. State and prove the converse of Proposition 32.

2. Use this proposition to shew that the tangents drawn to a circle from an external point are equal.

3. If two circles touch one another, any straight line drawn through the point of contact cuts off similar segments.

Prove this for (i) internal, (ii) external contact.

4. If two circles touch one another, and from A, the point of contact, two chords APQ, AXY are drawn : then PX and QY are parallel.

Prove this for (i) internal, (ii) external contact.

5. Two circles intersect at the points A, B : and one of them passes through O, the centre of the other : prove that OA bisects the angle between the common chord and the tangent to the first circle at A.


6. Two circles intersect at A and B ; and through P, any point on the circumference of one of them, straight lines PAC, PBD are drawn to cut the other circle at C and D : shew that CD is parallel to the tangent at P.

7. If from the point of contact of a tangent to a circle, a chord is drawn, the perpendiculars dropped on the tangent and chord from the middle point of either arc cut off by the chord are equal.



## PROPOSITION 33. PROBLEM.

On a given straight line to describe a segment of a circle which shall contain an angle equal to a given angle.



Let  $AB$  be the given st. line, and  $C$  the given angle.

It is required to describe on  $AB$  a segment of a circle which shall contain an angle equal to  $C$ .

Construction.

At  $A$  in  $BA$ , make the  $\angle BAD$  equal to the  $\angle C$ . I. 23.

From  $A$  draw  $AE$  at rt. angles to  $AD$ . I. 11.

Bisect  $AB$  at  $F$ . I. 10.

From  $F$  draw  $FG$  at rt. angles to  $AB$ , cutting  $AE$  at  $G$ .

Join  $GB$ .

Then in the  $\triangle AFG, BFG$ ,

Because  $\left\{ \begin{array}{l} AF = BF, \\ \text{and } FG \text{ is common,} \\ \text{and the } \angle AFG = \text{the } \angle BFG, \text{ being rt. angles;} \end{array} \right. \quad \text{Constr.}$

$\therefore GA = GB$ : I. 4.

$\therefore$  the circle described with centre  $G$ , and radius  $GA$ , will pass through  $B$ .

Describe this circle, and call it  $ABH$ .

Then the segment  $AHB$  shall contain an angle equal to  $C$ .

Proof. Because  $AD$  is drawn at rt. angles to the radius  $GA$  from its extremity  $A$ ,

$\therefore AD$  is a tangent to the circle; III. 16.

and from  $A$ , its point of contact, a chord  $AB$  is drawn;


$\therefore$  the  $\angle BAD =$  the angle in the alt. segment  $AHB$ . III. 32.

But the  $\angle BAD =$  the  $\angle C$ : Constr.

$\therefore$  the angle in the segment  $AHB =$  the  $\angle C$ .

$\therefore AHB$  is the segment required. Q.E.F.

**NOTE.** In the particular case when the given angle  $C$  is a rt. angle, the segment required will be the semicircle described on the given st. line  $AB$ ; for the angle in a semicircle is a rt. angle.



III. 31.

**EXERCISES.**

[The following exercises depend on the corollary to the Converse of Proposition 21 given on page 201, namely

*The locus of the vertices of triangles which stand on the same base and have a given vertical angle, is the arc of the segment standing on this base, and containing an angle equal to the given angle.*

Exercises 1 and 2 afford good illustrations of the method of finding required points by the *Intersection of Loci*. See page 125.]

1. Describe a triangle on a given base, having a given vertical angle, and having its vertex on a given straight line.

2. Construct a triangle, having given the base, the vertical angle and

- (i) one other side.
- (ii) the altitude.
- (iii) the length of the median which bisects the base.
- (iv) the point at which the perpendicular from the vertex meets the base.

3. Construct a triangle having given the base, the vertical angle, and the point at which the base is cut by the bisector of the vertical angle.

[Let  $AB$  be the base,  $X$  the given point in it, and  $K$  the given angle. On  $AB$  describe a segment of a circle containing an angle equal to  $K$ ; complete the  $\odot^\infty$  by drawing the arc  $APB$ . Bisect the arc  $APB$  at  $P$ : join  $PX$ , and produce it to meet the  $\odot^\infty$  at  $C$ . Then  $ABC$  shall be the required triangle.]


4. Construct a triangle having given the base, the vertical angle, and the sum of the remaining sides.

[Let  $AB$  be the given base,  $K$  the given angle, and  $H$  the given line equal to the sum of the sides. On  $AB$  describe a segment containing an angle equal to  $K$ , also another segment containing an angle equal to half the  $\angle K$ . From centre  $A$ , with radius  $H$ , describe a circle cutting the arc of the last drawn segment at  $X$  and  $Y$ . Join  $AX$  (or  $AY$ ) cutting the arc of the first segment at  $C$ . Then  $ABC$  shall be the required triangle.]

5. Construct a triangle having given the base, the vertical angle, and the difference of the remaining sides.

## PROPOSITION 34. PROBLEM.

*From a given circle to cut off a segment which shall contain an angle equal to a given angle.*



Let  $ABC$  be the given circle, and  $D$  the given angle.

*It is required to cut off from the  $\odot ABC$  a segment which shall contain an angle equal to  $D$ .*

**Construction.** Take any point  $B$  on the  $\odot^{\circ}$ ,  
and at  $B$  draw the tangent  $EBF$ . III. 17.

At  $B$ , in  $FB$ , make the  $\angle FBC$  equal to the  $\angle D$ . I. 23.

*Then the segment  $BAC$  shall contain an angle equal to  $D$ .*

**Proof.** Because  $EF$  is a tangent to the circle, and from  $B$ , its point of contact, a chord  $BC$  is drawn,

$\therefore$  the  $\angle FBC =$  the angle in the alternate segment  $BAC$ .

III. 32.

But the  $\angle FBC =$  the  $\angle D$ ;

*Constr.*

$\therefore$  the angle in the segment  $BAC =$  the  $\angle D$ .

Hence from the given  $\odot ABC$  a segment  $BAC$  has been cut off, containing an angle equal to  $D$ . Q.E.F.

## EXERCISES.

1. The chord of a given segment of a circle is produced to a fixed point: on this straight line so produced draw a segment of a circle similar to the given segment.

2. Through a given point without a circle draw a straight line that will cut off a segment capable of containing an angle equal to a given angle.

QUESTIONS FOR REVISION.

1. Enunciate the propositions from which we infer that a straight line and a circle must either

- (i) intersect in two points ; or
- (ii) touch at one point ; or
- (iii) have no point in common.

2. Give two independent constructions for drawing a tangent to a circle from an external point.

Shew that the two tangents so drawn

- (i) are equal ;
- (ii) subtend equal angles at the centre ;
- (iii) make equal angles with the straight line which joins the given point to the centre.

3. Enunciate propositions relating to

- (i) angles in a segment of a circle ;
- (ii) similar segments of circles.

4. What are *conjugate arcs* of a circle ?

*The angles in conjugate segments of a circle are supplementary.*

How does Euclid enunciate this theorem ? State and prove its converse.

5. Explain what is meant by a *reflex angle*. What simplifications may be made in the proofs of Third Book Propositions if reflex angles are admitted ?

6. If the circumference of a circle is divided into six equal arcs, shew that the chords joining successive points of division are all equal to the radius of the circle.


7. Find the locus of the centres of all circles

- (i) which pass through two given points ;
- (ii) which touch a given circle at a given point ;
- (iii) which are of given radius, and touch a given circle ;
- (iv) which are of given radius, and pass through a given point ;
- (v) which touch a given straight line at a given point ;
- (vi) which touch each of two parallel straight lines ;
- (vii) which touch each of two intersecting straight lines of unlimited length.

8. If a system of triangles stand on the same base and on the same side of it, and have equal vertical angles, shew that the locus of their vertices is the arc of a circle. Prove this theorem, having first enunciated the proposition of which it is the converse.

## PROPOSITION 35. THEOREM.

If two chords of a circle cut one another, the rectangle contained by the segments of one shall be equal to the rectangle contained by the segments of the other.



Let AB, CD, two chords of the  $\odot$  ACBD, cut one another at E.

Then shall the rect. AE, EB = the rect. CE, ED.

Construction. Find F, the centre of the  $\odot$  ACB; III. 1.  
From F draw FG, FH perp. respectively to AB, CD. I. 12.  
Join FA, FE, FD.

Proof. Because FG is drawn from the centre F perp. to AB,  
 $\therefore$  AB is bisected at G. III. 3.

For a similar reason CD is bisected at H.

Again, because AB is divided equally at G, and unequally at E,  
 $\therefore$  the rect. AE, EB with the sq. on EG = the sq. on AG. II. 5.

To each of these equals add the sq. on GF;  
then the rect. AE, EB with the sq. on EG, GF = the sum of  
the sqq. on AG, GF.

But the sqq. on EG, GF = the sq. on FE; I. 47.  
and the sqq. on AG, GF = the sq. on AF;  
for the angles at G are rt. angles.

$\therefore$  the rect. AE, EB with the sq. on FE = the sq. on AF.

Similarly it may be shewn that

the rect. CE, ED with the sq. on FE = the sq. on FD.

But the sq. on AF = the sq. on FD; for AF = FD.

$\therefore$  the rect. AE, EB with the sq. on FE = the rect. CE, ED  
with the sq. on FE.

From these equals take the sq. on FE:  
then the rect. AE, EB = the rect. CE, ED. Q.E.D.

**COROLLARY.** *If through a fixed point within a circle any number of chords are drawn, the rectangles contained by their segments are all equal.*

**NOTE.** The following special cases of this proposition deserve notice :

- (i) when the given chords both pass through the centre :
- (ii) when one chord passes through the centre, and cuts the other at right angles :
- (iii) when one chord passes through the centre, and cuts the other obliquely.

In each of these cases the general proof requires some modification, which may be left as an exercise to the student. .

#### EXERCISES.

1. Two straight lines  $AB$ ,  $CD$  intersect at  $E$ , so that the rectangle  $AE$ ,  $EB$  is equal to the rectangle  $CE$ ,  $ED$  ; shew that the four points  $A$ ,  $B$ ,  $C$ ,  $D$  are concyclic.

2. The rectangle contained by the segments of any chord drawn through a given point within a circle is equal to the square on half the shortest chord which may be drawn through that point.

3.  $ABC$  is a triangle right-angled at  $C$  ; and from  $C$  a perpendicular  $CD$  is drawn to the hypotenuse : shew that the square on  $CD$  is equal to the rectangle  $AD$ ,  $DB$ .

4.  $ABC$  is a triangle ; and  $AP$ ,  $BQ$ , the perpendiculars dropped from  $A$  and  $B$  on the opposite sides, intersect at  $O$  : shew that the rectangle  $AO$ ,  $OP$  is equal to the rectangle  $BO$ ,  $OQ$ .

5. Two circles intersect at  $A$  and  $B$ , and through any point in  $AB$  their common chord two chords are drawn, one in each circle ; shew that their four extremities are concyclic.

6.  $A$  and  $B$  are two points within a circle such that the rectangle contained by the segments of any chord drawn through  $A$  is equal to the rectangle contained by the segments of any chord through  $B$  : shew that  $A$  and  $B$  are equidistant from the centre.


7. If through  $E$ , a point without a circle, two secants,  $EAB$ ,  $ECD$  are drawn ; shew that the rectangle  $EA$ ,  $EB$  is equal to the rectangle  $EC$ ,  $ED$ .

[Proceed as in III. 35, using II. 6.]

8. Through  $A$ , a point of intersection of two circles, two straight lines  $CAE$ ,  $DAF$  are drawn, each passing through a centre and terminated by the circumferences : shew that the rectangle  $CA$ ,  $AE$  is equal to the rectangle  $DA$ ,  $AF$ .

## PROPOSITION 36. THEOREM.

If from any point without a circle a tangent and a secant are drawn, then the rectangle contained by the whole secant and the part of it without the circle shall be equal to the square on the tangent.



Let ABC be a circle; and from D, a point without it, let there be drawn the secant DCA, and the tangent DB.

Then the rect. DA, DC shall be equal to the sq. on DB.

**Construction.** Find E, the centre of the  $\odot$  ABC: III. 1.  
and from E, draw EF perp. to AD. I. 12.  
Join EB, EC, ED.

**Proof.** Because EF, passing through the centre, is perp. to the chord AC,

$\therefore$  AC is bisected at F. III. 3.

And since AC is bisected at F and produced to D,  
 $\therefore$  the rect. DA, DC with the sq. on FC = the sq. on FD. II. 6.

To each of these equals add the sq. on EF:  
then the rect. DA, DC with the sqq. on EF, FC = the sqq. on EF, FD.

But the sqq. on EF, FC = the sq. on EC; for EFC is a rt. angle;  
= the sq. on EB.

And the sqq. on EF, FD = the sq. on ED; for EFD is a rt. angle;  
= the sqq. on EB, BD; for EBD is a  
rt. angle. III. 18.

$\therefore$  the rect. DA, DC with the sq. on EB = the sqq. on EB, BD.


From these equals take the sq. on EB:

then the rect. DA, DC = the sq. on DB. Q.E.D.

**NOTE.** This proof may easily be adapted to the case when the secant passes through the centre of the circle.

**COROLLARY.** *If from a given point without a circle any number of secants are drawn, the rectangles contained by the whole secants and the parts of them without the circle are all equal; for each of these rectangles is equal to the square on the tangent drawn from the given point to the circle.*

For instance, in the adjoining figure, each of the rectangles PB, PA and PD, PC and PF, PE is equal to the square on the tangent PQ:



$$\begin{aligned} \therefore \text{ the rect. PB, PA} \\ &= \text{ the rect. PD, PC} \\ &= \text{ the rect. PF, PE.} \end{aligned}$$

**NOTE.** Remembering that the segments into which the chord AB is divided at P, are the lines PA, PB, (see Def., page 139) we are enabled to include the corollaries of Propositions 35 and 36 in a single enunciation.

*If any number of chords of a circle are drawn through a given point within or without a circle, the rectangles contained by the segments of the chords are equal.*


**EXERCISES.**

1. Use this proposition to shew that tangents drawn to a circle from an external point are equal.
2. If two circles intersect, tangents drawn to them from any point in their common chord produced are equal.
3. If two circles intersect at A and B, and PQ is a tangent to both circles; shew that AB produced bisects PQ.
4. If P is any point on the straight line AB produced, shew that the tangents drawn from P to all circles which pass through A and B are equal.
5. ABC is a triangle right-angled at C, and from any point P in AC, a perpendicular PQ is drawn to the hypotenuse: shew that the rectangle AC, AP is equal to the rectangle AB, AQ.
6. ABC is a triangle right-angled at C, and from C a perpendicular CD is drawn to the hypotenuse: shew that the rect. AB, AD is equal to the square on AC.



## PROPOSITION 37. THEOREM.

If from a point without a circle there are drawn two straight lines, one of which cuts the circle, and the other meets it, and if the rectangle contained by the whole line which cuts the circle and the part of it without the circle is equal to the square on the line which meets the circle, then the line which meets the circle shall be a tangent to it.



Let ABC be a circle; and from D, a point without it, let there be drawn two st. lines DCA and DB, of which DCA cuts the circle at C and A, and DB meets it; and let the rect. DA, DC = the sq. on DB.

Then shall DB be a tangent to the circle.

**Construction.** From D draw DE to touch the  $\odot$  ABC: III. 17.  
let E be the point of contact.

Find the centre F, and join FB, FD, FE. III. 1.

**Proof.** Since DCA is a secant, and DE a tangent to the circle,  
 $\therefore$  the rect. DA, DC = the sq. on DE, III. 36.

But, by hypothesis, the rect. DA, DC = the sq. on DB;

$\therefore$  the sq. on DE = the sq. on DB;

$\therefore$  DE = DB.

Hence in the  $\triangle$  DBF, DEF,

Because  $\begin{cases} \text{DB} = \text{DE}, \\ \text{and } \text{BF} = \text{EF}; \\ \text{and } \text{DF} \text{ is common;} \end{cases}$  *Proved.*  
I. Def. 15.

$\therefore$  the  $\angle$  DBF = the  $\angle$  DEF. I. 8.

But DEF is a rt. angle, for DE is a tangent; III. 18.

$\therefore$  DBF is also a rt. angle;

and since BF is a radius,

$\therefore$  DB touches the  $\odot$  ABC at the point B. Q.E.D.

NOTE ON THE METHOD OF LIMITS AS APPLIED TO TANGENCY.

Euclid defines a tangent to a circle as a straight line which meets the circumference, but being produced, does not cut it: and from this definition he deduces the fundamental theorem that a tangent is perpendicular to the radius drawn to the point of contact. III. Prop. 16.

But this result may also be established by the Method of Limits, which regards the tangent as the ultimate position of a secant when its two points of intersection with the circumference are brought into coincidence [See Note on page 165]: and it may be shewn that every theorem relating to the tangent may be derived from some more general proposition relating to the secant, by considering the ultimate case when the two points of intersection coincide.

1. To prove by the Method of Limits that a tangent to a circle is at right angles to the radius drawn to the point of contact.

Let  $ABD$  be a circle, whose centre is  $C$ ; and  $PABQ$  a secant cutting the  $\bigcirc^{\infty}$  in  $A$  and  $B$ ; and let  $P'AQ'$  be the limiting position of  $PQ$  when the point  $B$  is brought into coincidence with  $A$ .

Then shall  $CA$  be perp. to  $P'Q'$ .


Bisect  $AB$  at  $E$  and join  $CE$ :

then  $CE$  is perp. to  $PQ$ . III. 3.

Now let the secant  $PABQ$  change its position in such a way that while the point  $A$  remains fixed, the point  $B$  continually approaches  $A$ , and ultimately coincides with it;

then, however near  $B$  approaches to  $A$ , the st. line  $CE$  is always perp. to  $PQ$ , since it joins the centre to the middle point of the chord  $AB$ .

But in the limiting position, when  $B$  coincides with  $A$ , and the secant  $PQ$  becomes the tangent  $P'Q'$ , it is clear that the point  $E$  will also coincide with  $A$ ; and the perpendicular  $CE$  becomes the radius  $CA$ . Hence  $CA$  is perp. to the tangent  $P'Q'$  at its point of contact  $A$ . Q. E. D.



NOTE. It follows from Proposition 2 that a straight line cannot cut the circumference of a circle at more than two points. Now when the two points in which a secant cuts a circle move towards coincidence, the secant ultimately becomes a tangent to the circle: we infer therefore that a tangent cannot meet a circle otherwise than at its point of contact. Thus Euclid's definition of a tangent may be deduced from that given by the Method of Limits.

2. *By this method Proposition 32 may be derived as a special case from Proposition 21.*

For let  $A$  and  $B$  be two points on the  $\odot^{\infty}$  of the  $\odot ABC$ ;

and let  $\angle BCA$ ,  $\angle BPA$  be any two angles in the segment  $BCPA$ ;

then the  $\angle BPA = \angle BCA$ . III. 21.

Produce  $PA$  to  $Q$ .

Now let the point  $P$  continually approach the fixed point  $A$ , and ultimately coincide with it;

then, however near  $P$  may approach to  $A$ , the  $\angle BPQ = \angle BCA$ . III. 21.


But in the limiting position when  $P$  coincides with  $A$ ,

and the secant  $PAQ$  becomes the tangent  $AQ'$ ,

it is clear that  $BP$  will coincide with  $BA$ ,

and the  $\angle BPQ$  becomes the  $\angle BAQ'$ .

Hence the  $\angle BAQ' = \angle BCA$ , in the alternate segment. Q.E.D.



The contact of circles may be treated in a similar manner by adopting the following definition.

**DEFINITION.** If one or other of two intersecting circles alters its position in such a way that the two points of intersection continually approach one another, and ultimately coincide; in the limiting position they are said to touch one another, and the point in which the two points of intersection ultimately coincide is called the point of contact.

#### EXAMPLES ON LIMITS.

1. Deduce Proposition 19 from the Corollary of Proposition 1 and Proposition 3.

2. Deduce Propositions 11 and 12 from Ex. 1, page 171.

3. Deduce Proposition 6 from Proposition 5.

4. Deduce Proposition 13 from Proposition 10.

5. Shew that a straight line cuts a circle in two different points, two coincident points, or not at all, according as its distance from the centre is less than, equal to, or greater than a radius.

6. Deduce Proposition 32 from Ex. 3, page 202.

7. Deduce Proposition 36 from Ex. 7, page 227.

8. *The angle in a semi-circle is a right angle.*

To what Theorem is this statement reduced, when the vertex of the right angle is brought into coincidence with an extremity of the diameter?

9. From Ex. 1, page 204, deduce the corresponding property of a triangle inscribed in a circle.


## THEOREMS AND EXAMPLES ON BOOK III.

## I. ON THE CENTRE AND CHORDS OF A CIRCLE.

[See Propositions 1, 3, 14, 15, 25.]

1. *All circles which pass through a fixed point, and have their centres on a given straight line, pass also through a second fixed point.*

Let  $AB$  be the given st. line, and  $P$  the given point.



From  $P$  draw  $PR$  perp. to  $AB$ ;  
and produce  $PR$  to  $P'$ , making  $RP'$  equal to  $PR$ .

*Then all circles which pass through  $P$ , and have their centres on  $AB$ , shall pass also through  $P'$ .*

For let  $C$  be the centre of *any* one of these circles.

Join  $CP, CP'$ .

Then in the  $\triangle^s$   $CRP, CRP'$ ,

Because  $\left\{ \begin{array}{l} CR \text{ is common,} \\ \text{and } RP = RP', \\ \text{and the } \angle CRP = \text{the } \angle CRP', \text{ being rt. angles;} \end{array} \right. \quad \begin{array}{l} \text{Constr.} \\ \text{I. 4.} \end{array}$

$\therefore CP = CP'$ ;

$\therefore$  the circle whose centre is  $C$ , and which passes through  $P$ , must pass also through  $P'$ .

But  $C$  is the centre of *any* circle of the system;

$\therefore$  all circles, which pass through  $P$ , and have their centres in  $AB$ , pass also through  $P'$ . Q. E. D.

2. *Describe a circle that shall pass through three given points not in the same straight line.*

3. Describe a circle that shall pass through two given points and have its centre in a given straight line. When is this impossible?

4. Describe a circle of given radius to pass through two given points. When is this impossible?

5.  $ABC$  is an isosceles triangle; and from the vertex  $A$ , as centre, a circle is described cutting the base, or the base produced, at  $X$  and  $Y$ . Shew that  $BX = CY$ .

6. If two circles which intersect are cut by a straight line parallel to the common chord, shew that the parts of it intercepted between the circumferences are equal.

7. If two circles cut one another, any two straight lines drawn through a point of section, making equal angles with the common chord, and terminated by the circumferences, are equal.

[Ex. 12, p. 171.]

8. If two circles cut one another, of all straight lines drawn through a point of section and terminated by the circumferences, the greatest is that which is parallel to the line joining the centres.

9. Two circles, whose centres are  $C$  and  $D$ , intersect at  $A$ ,  $B$ ; and through  $A$  a straight line  $PAQ$  is drawn terminated by the circumferences: if  $PC$ ,  $QD$  intersect at  $X$ , shew that the angle  $PXQ$  is equal to the angle  $CAD$ .

10. Through a point of section of two circles which cut one another draw a straight line terminated by the circumferences and bisected at the point of section.

11.  $AB$  is a fixed diameter of a circle, whose centre is  $C$ ; and from  $P$ , any point on the circumference,  $PQ$  is drawn perpendicular to  $AB$ ; shew that the bisector of the angle  $CPQ$  always intersects the circle in one or other of two fixed points.

12. Circles are described on the sides of a quadrilateral as diameters: shew that the common chord of any two consecutive circles is parallel to the common chord of the other two.

[Ex. 9, p. 105.]

13. Two equal circles touch one another externally, and through the point of contact two chords are drawn, one in each circle, at right angles to each other: shew that the straight line joining their other extremities is equal to the diameter of either circle.

14. Straight lines are drawn from a given external point to the circumference of a circle: find the locus of their middle points. [Ex. 3, p. 105.]

15. Two equal segments of circles are described on opposite sides of the same chord  $AB$ ; and through  $O$ , the middle point of  $AB$ , any straight line  $POQ$  is drawn, intersecting the arcs of the segments at  $P$  and  $Q$ : shew that  $OP = OQ$ .

## II. ON THE TANGENT AND THE CONTACT OF CIRCLES.

[See Propositions 11, 12, 16, 17, 18, 19.]

1. All equal chords placed in a given circle touch a fixed concentric circle.

2. If from an external point two tangents are drawn to a circle, the angle contained by them is double the angle contained by the chord of contact and the diameter drawn through one of the points of contact.

3. Two circles touch one another externally, and through the point of contact a straight line is drawn terminated by the circumferences: shew that the tangents at its extremities are parallel.

4. Two circles intersect, and through one point of section any straight line is drawn terminated by the circumferences: shew that the angle between the tangents at its extremities is equal to the angle between the tangents at the point of section.

5. Shew that two parallel tangents to a circle intercept on any third tangent a segment which subtends a right angle at the centre.

6. Two tangents are drawn to a given circle from a fixed external point  $A$ , and any third tangent cuts them produced at  $P$  and  $Q$ : shew that  $PQ$  subtends a constant angle at the centre of the circle.

7. *In any quadrilateral circumscribed about a circle, the sum of one pair of opposite sides is equal to the sum of the other pair.*

8. *If the sum of one pair of opposite sides of a quadrilateral is equal to the sum of the other pair, shew that a circle may be inscribed in the figure.*

[Bisect two adjacent angles of the figure, and so describe a circle to touch three of its sides. Then prove indirectly by means of the last exercise that this circle must also touch the fourth side.]

9. Two circles touch one another internally, the centre of the outer being within the inner circle: shew that of all chords of the outer circle which touch the inner, the greatest is that which is perpendicular to the straight line joining the centres.

10. In any triangle, if a circle is described from the middle point of one side as centre and with a radius equal to half the sum of the other two sides, it will touch the circles described on these sides as diameters.

11. Through a given point, draw a straight line to cut a circle, so that the part intercepted by the circumference may be equal to a given straight line.

In order that the problem may be possible, between what limits must the given line lie, when the given point is (i) without the circle, (ii) within it?

12. A series of circles touch a given straight line at a given point: shew that the tangents to them at the points where they cut a given parallel straight line all touch a fixed circle, whose centre is the given point.

13. If two circles touch one another internally, and any third circle be described touching one internally and the other externally; then the sum of the distances of the centre of this third circle from the centres of the two given circles is constant.

14. Find the locus of points such that the pairs of tangents drawn from them to a given circle contain a constant angle.

15. Find a point such that the tangents drawn from it to two given circles may be equal to two given straight lines. When is this impossible?

16. If three circles touch one another two and two; prove that the tangents drawn to them at the three points of contact are concurrent and equal.

#### THE COMMON TANGENTS TO TWO CIRCLES.


17. To draw a common tangent to two circles.

*First.* When the given circles are external to one another, or when they intersect.

Let  $A$  be the centre of the greater circle, and  $B$  the centre of the less.

From  $A$ , with radius equal to the diff<sup>ce</sup> of the radii of the given circles, describe a circle: and from  $B$  draw  $BC$  to touch the last drawn circle. Join  $AC$ , and produce it to meet the greater of the given circles at  $D$ .

Through  $B$  draw the radius  $BE$  par<sup>l</sup> to  $AD$ , and in the same direction.



Join  $DE$ .

*Then*  $DE$  shall be a common tangent to the two given circles.

For since  $AC =$  the diff<sup>ce</sup> between  $AD$  and  $BE$ ,

*Constr.*

$\therefore CD = BE$ ;

and  $CD$  is par<sup>l</sup> to  $BE$ ;

*Constr.*

$\therefore DE$  is equal and par<sup>l</sup> to  $CB$ .

I. 33.

But since  $BC$  is a tangent to the circle at  $C$ ,

$\therefore$  the  $\angle ACB$  is a rt. angle;

III. 18.

hence each of the angles at  $D$  and  $E$  is a rt. angle:

I. 29.

$\therefore DE$  is a tangent to both circles.

Q.E.F.

It follows from hypothesis that the point B is outside the circle used in the construction :

$\therefore$  two tangents such as BC may always be drawn to it from B ;  
hence two common tangents may always be drawn to the given circles by the above method. These are called the **direct common tangents**.

*Secondly.* When the given circles are external to one another and do not intersect, two more common tangents may be drawn.

For, from centre A, with a radius equal to the *sum* of the radii of the given circles, describe a circle.

From B draw a tangent to this circle ;  
and proceed as before, but draw BE in the direction *opposite* to AD.

It follows from hypothesis that B is external to the circle used in the construction ;

$\therefore$  two tangents may be drawn to it from B.

Hence two more common tangents may be drawn to the given circles : these will be found to pass between the given circles, and are called the **transverse common tangents**.

Thus, in general, *four* common tangents may be drawn to two given circles.

The student should investigate for himself the number of common tangents which may be drawn in the following special cases, noting in each case where the general construction fails, or is modified :—

- (i) When the given circles intersect :
- (ii) When the given circles have external contact :
- (iii) When the given circles have internal contact :
- (iv) When one of the given circles is wholly within the other.

18. *Draw the direct common tangents to two equal circles.*

19. If the two direct, or the two transverse, common tangents are drawn to two circles, the parts of the tangents intercepted between the points of contact are equal.

20. If four common tangents are drawn to two circles external to one another ; shew that the two direct, and also the two transverse, tangents intersect on the straight line which joins the centres of the circles.

21. Two given circles have external contact at A, and a direct common tangent is drawn to touch them at P and Q : shew that PQ subtends a right angle at the point A.

22. Two circles have external contact at A, and a direct common tangent is drawn to touch them at P and Q : shew that a circle described on PQ as diameter is touched at A by the straight line which joins the centres of the circles.



23. Two circles whose centres are  $C$  and  $C'$  have external contact at  $A$ , and a direct common tangent is drawn to touch them at  $P$  and  $Q$ : shew that the bisectors of the angles  $PCA$ ,  $QC'A$  meet at right angles in  $PQ$ . And if  $R$  is the point of intersection of the bisectors, shew that  $RA$  is also a common tangent to the circles.

24. Two circles have external contact at  $A$ , and a direct common tangent is drawn to touch them at  $P$  and  $Q$ : shew that the square on  $PQ$  is equal to the rectangle contained by the diameters of the circles.

25. Draw a tangent to a given circle, so that the part of it intercepted by another given circle may be equal to a given straight line. When is this impossible?

26. Draw a secant to two given circles, so that the parts of it intercepted by the circumferences may be equal to two given straight lines.

#### PROBLEMS ON TANGENCY.

*Obs.* The following exercises are solved by the Method of Intersection of Loci, explained on page 125.

The student should begin by making himself familiar with the following loci.

(i) *The locus of the centres of circles which pass through two given points.*

(ii) *The locus of the centres of circles which touch a given straight line at a given point.*

(iii) *The locus of the centres of circles which touch a given circle at a given point.*

(iv) *The locus of the centres of circles which touch a given straight line, and have a given radius.*

(v) *The locus of the centres of circles which touch a given circle, and have a given radius.*

(vi) *The locus of the centres of circles which touch two given straight lines.*

In each exercise the student should investigate the limits and relations among the data, in order that the problem may be possible.

27. Describe a circle to touch three given straight lines.

28. Describe a circle to pass through a given point, and touch a given straight line at a given point.

29. Describe a circle to pass through a given point, and touch a given circle at a given point.

30. Describe a circle of given radius to pass through a given point, and touch a given straight line.
31. Describe a circle of given radius to touch two given circles.
32. Describe a circle of given radius to touch two given straight lines.
33. Describe a circle of given radius to touch a given circle and a given straight line.
34. Describe two circles of given radii to touch one another and a given straight line, on the same side of it.
35. If a circle touches a given circle and a given straight line, shew that the points of contact and an extremity of the diameter of the given circle at right angles to the given line are collinear.

36. *To describe a circle to touch a given circle, and also to touch a given straight line at a given point.*


Let  $DEB$  be the given circle,  $PQ$  the given straight line, and  $A$  the given point in it.

*It is required to describe a circle to touch the  $\odot DEB$ , and also to touch  $PQ$  at  $A$ .*

At  $A$  draw  $AF$  perp. to  $PQ$ : I. 11. then the centre of the required circle must lie in  $AF$ . III. 19.

Find  $C$ , the centre of the  $\odot DEB$ , III. 1.

and draw a diameter  $BD$  perp. to  $PQ$ : join  $A$  to one extremity  $D$ , cutting the  $\odot$  at  $E$ .



Join  $CE$ , and produce it to cut  $AF$  at  $F$ .

*Then  $F$  shall be the centre, and  $FA$  the radius of the required circle.*

[Supply the proof: and shew that a second solution is obtained by joining  $AB$ , and producing it to meet the  $\odot$  at  $E$ .

Also distinguish between the nature of the contact of the circles, when  $PQ$  cuts, touches, or is without the given circle.]

37. Describe a circle to touch a given straight line, and to touch a given circle at a given point.
38. Describe a circle to touch a given circle, have its centre in a given straight line, and pass through a given point in that straight line.

[For other problems of the same class see page 253.]

## ORTHOGONAL CIRCLES.

**DEFINITION.** Circles which intersect at a point, so that the two tangents at that point are at right angles to one another, are said to be **orthogonal**, or to **cut one another orthogonally**.

39. In two intersecting circles the angle between the tangents at one point of intersection is equal to the angle between the tangents at the other.

40. *If two circles cut one another orthogonally, the tangent to each circle at a point of intersection will pass through the centre of the other circle.*

41. *If two circles cut one another orthogonally, the square on the distance between their centres is equal to the sum of the squares on their radii.*

42. Find the locus of the centres of all circles which cut a given circle orthogonally at a given point.

43. Describe a circle to pass through a given point and cut a given circle orthogonally at a given point.

### III. ON ANGLES IN SEGMENTS, AND ANGLES AT THE CENTRES AND CIRCUMFERENCES OF CIRCLES.

[See Propositions 20, 21, 22 ; 26, 27, 28, 29 ; 31, 32, 33, 34.]

1. *If two chords intersect within a circle, they form an angle equal to that at the centre, subtended by half the sum of the arcs they cut off.*

Let  $AB$  and  $CD$  be two chords, intersecting at  $E$  within the given  $\odot ADBC$ .

Then shall the  $\angle AEC$  be equal to the angle at the centre, subtended by half the sum of the arcs  $AC$ ,  $BD$ .

Join  $AD$ .

Then the ext.  $\angle AEC =$  the sum of the int. opp.  $\angle^s$   $EDA$ ,  $EAD$  ;


that is, the sum of the  $\angle^s$   $CDA$ ,  $BAD$ .

But the  $\angle^s$   $CDA$ ,  $BAD$  are the angles at the  $\odot^c$  subtended by the arcs  $AC$ ,  $BD$  ;

$\therefore$  their sum = half the sum of the angles at the centre subtended by the same arcs ;

or, the  $\angle AEC =$  the angle at the centre subtended by half the sum of the arcs  $AC$ ,  $BD$ .

Q. E. D.



2. *If two chords when produced intersect outside a circle, they form an angle equal to that at the centre subtended by half the difference of the arcs they cut off.*

3. The sum of the arcs cut off by two chords of a circle at right angles to one another is equal to the semi-circumference.

4. AB, AC are any two chords of a circle; and P, Q are the middle points of the minor arcs cut off by them: if PQ is joined, cutting AB and AC at X, Y, shew that  $AX = AY$ .

5. *If one side of a quadrilateral inscribed in a circle is produced, the exterior angle is equal to the opposite interior angle.*

6. If two circles intersect, and any straight lines are drawn, one through each point of section, terminated by the circumferences; shew that the chords which join their extremities towards the same parts are parallel.

7. ABCD is a quadrilateral inscribed in a circle; and the opposite sides AB, DC are produced to meet at P, and CB, DA to meet at Q: if the circles circumscribed about the triangles PBC, QAB intersect at R, shew that the points P, R, Q are collinear.

8. If a circle is described on one of the sides of a right-angled triangle, then the tangent drawn to it at the point where it cuts the hypotenuse bisects the other side.

9. Given three points not in the same straight line: shew how to find any number of points on the circle which passes through them, without finding the centre.

10. Through any one of three given points not in the same straight line, draw a tangent to the circle which passes through them, without finding the centre.

11. Of two circles which intersect at A and B, the circumference of one passes through the centre of the other: from A any straight line is drawn to cut the first at C, the second at D; shew that  $CB = CD$ .

12. Two tangents AP, AQ are drawn to a circle, and B is the middle point of the arc PQ, convex to A. Shew that PB bisects the angle APQ.

13. Two circles intersect at A and B; and at A tangents are drawn, one to each circle, to meet the circumferences at C and D; if CB, BD are joined, shew that the triangles ABC, DBA are equiangular to one another.

14. Two segments of circles are described on the same chord and on the same side of it; the extremities of the common chord are joined to any point on the arc of the exterior segment: shew that the arc intercepted on the interior segment is constant.

15. If a series of triangles are drawn standing on a fixed base, and having a given vertical angle, show that the bisectors of the vertical angles all pass through a fixed point.

16.  $ABC$  is a triangle inscribed in a circle, and  $E$  the middle point of the arc subtended by  $BC$  on the side remote from  $A$ : if through  $E$  a diameter  $ED$  is drawn, shew that the angle  $DEA$  is half the difference of the angles at  $B$  and  $C$ . [See Ex. 7, p. 109.]

17. If two circles touch each other internally at a point  $A$ , any chord of the exterior circle which touches the interior is divided at its point of contact into segments which subtend equal angles at  $A$ .

18. If two circles touch one another internally, and a straight line is drawn to cut them, the segments of it intercepted between the circumferences subtend equal angles at the point of contact.

#### THE ORTHOCENTRE OF A TRIANGLE.

19. *The perpendiculars drawn from the vertices of a triangle to the opposite sides are concurrent.*

In the  $\triangle ABC$ , let  $AD$ ,  $BE$  be the perp<sup>s</sup> drawn from  $A$  and  $B$  to the opposite sides; and let them intersect at  $O$ . Join  $CO$ ; and produce it to meet  $AB$  at  $F$ .

*It is required to shew that  $CF$  is perp. to  $AB$ .*

Join  $DE$ .

Then, because the  $\angle^s$   $OEC$ ,  $ODC$  are rt. angles,


*Hyp.*  
 $\therefore$  the points  $O$ ,  $E$ ,  $C$ ,  $D$  are concyclic:  
 $\therefore$  the  $\angle DEC =$  the  $\angle DOC$ , in the same segment;  
 $=$  the vert. opp.  $\angle FOA$ .

Again, because the  $\angle^s$   $AEB$ ,  $ADB$  are rt. angles,

*Hyp.*  
 $\therefore$  the points  $A$ ,  $E$ ,  $D$ ,  $B$  are concyclic:  
 $\therefore$  the  $\angle DEB =$  the  $\angle DAB$ , in the same segment.

$\therefore$  the sum of the  $\angle^s$   $FOA$ ,  $FAO =$  the sum of the  $\angle^s$   $DEC$ ,  $DEB$   
 $=$  a rt. angle: *Hyp.*  
 $\therefore$  the remaining  $\angle AFO =$  a rt. angle: I. 32.  
 that is,  $CF$  is perp. to  $AB$ .

Hence the three perp<sup>s</sup>  $AD$ ,  $BE$ ,  $CF$  meet at the point  $O$ . Q.E.D.



[For an Alternative Proof see p. 114.]

## DEFINITIONS.


- (i) The intersection of the perpendiculars drawn from the vertices of a triangle to the opposite sides is called its **orthocentre**.
- (ii) The triangle formed by joining the feet of the perpendiculars is called the **pedal** or **orthocentric triangle**.

20. In an acute-angled triangle the perpendiculars drawn from the vertices to the opposite sides bisect the angles of the pedal triangle through which they pass.

In the acute-angled  $\triangle ABC$ , let  $AD$ ,  $BE$ ,  $CF$  be the perp<sup>s</sup> drawn from the vertices to the opposite sides, meeting at the orthocentre  $O$ ; and let  $DEF$  be the pedal triangle.

Then shall  $AD$ ,  $BE$ ,  $CF$  bisect respectively the  $\angle$ 's  $FDE$ ,  $DEF$ ,  $EFD$ .

For, as in the last theorem, it may be shewn that the points  $O$ ,  $D$ ,  $C$ ,  $E$  are concyclic;



$\therefore$  the  $\angle ODE =$  the  $\angle OCE$ , in the same segment.

Similarly the points  $O$ ,  $D$ ,  $B$ ,  $F$  are concyclic;

$\therefore$  the  $\angle ODF =$  the  $\angle OBF$ , in the same segment.

But the  $\angle OCE =$  the  $\angle OBF$ , each being the comp<sup>t</sup> of the  $\angle BAC$ .

$\therefore$  the  $\angle ODE =$  the  $\angle ODF$ .

Similarly it may be shewn that the  $\angle$ 's  $DEF$ ,  $EFD$  are bisected by  $BE$  and  $CF$ . Q. E. D.

**COROLLARY.** (i) Every two sides of the pedal triangle are equally inclined to that side of the original triangle in which they meet.

For the  $\angle EDC =$  the comp<sup>t</sup> of the  $\angle ODE$   
 $=$  the comp<sup>t</sup> of the  $\angle OCE$   
 $=$  the  $\angle BAC$ .

Similarly it may be shewn that the  $\angle FDB =$  the  $\angle BAC$ ,

$\therefore$  the  $\angle EDC =$  the  $\angle FDB =$  the  $\angle A$ .

In like manner it may be proved that

the  $\angle DEC =$  the  $\angle FEA =$  the  $\angle B$ ,  
 and the  $\angle DFB =$  the  $\angle EFA =$  the  $\angle C$ .

**COROLLARY.** (ii) The triangles  $DEC$ ,  $AEF$ ,  $DBF$  are equiangular to one another and to the triangle  $ABC$ .

**NOTE.** If the angle  $BAC$  is obtuse, then the perpendiculars  $BE$ ,  $CF$  bisect externally the corresponding angles of the pedal triangle.

21. *In any triangle, if the perpendiculars drawn from the vertices on the opposite sides are produced to meet the circumscribed circle, then each side bisects that portion of the line perpendicular to it which lies between the orthocentre and the circumference.*

Let  $ABC$  be a triangle in which the perpendiculars  $AD$ ,  $BE$  are drawn, intersecting at  $O$  the orthocentre, and let  $AD$  be produced to meet the  $\circ^{\infty}$  of the circumscribing circle at  $G$ .

Then shall  $DO = DG$ .

Join  $BG$ .

Then in the two  $\triangle^s$   $OEA$ ,  $ODB$ ,  
the  $\angle OEA =$  the  $\angle ODB$ , being rt. angles ;  
and the  $\angle EOA =$  the vert. opp.  $\angle DOB$  ;

$\therefore$  the remaining  $\angle EAO =$  the remaining  $\angle DBO$ . I. 32.

But the  $\angle CAG =$  the  $\angle CBG$ , in the same segment ;

$\therefore$  the  $\angle DBO =$  the  $\angle DBG$ .

Then in the  $\triangle^s$   $DBO$ ,  $DBG$ ,


Because  $\left\{ \begin{array}{l} \text{the } \angle DBO = \text{the } \angle DBG, \\ \text{the } \angle BDO = \text{the } \angle BDG, \\ \text{and } BD \text{ is common ;} \end{array} \right.$

$\therefore DO = DG$ .

*Proved.*

I. 26.

Q. E. D.



22. *In an acute-angled triangle the three sides are the external bisectors of the angles of the pedal triangle : and in an obtuse-angled triangle the sides containing the obtuse angle are the internal bisectors of the corresponding angles of the pedal triangle.*

23. *If  $O$  is the orthocentre of the triangle  $ABC$ , shew that the angles  $BOC$ ,  $BAC$  are supplementary.*

24. *If  $O$  is the orthocentre of the triangle  $ABC$ , then any one of the four points  $O$ ,  $A$ ,  $B$ ,  $C$  is the orthocentre of the triangle whose vertices are the other three.*

25. *The three circles which pass through two vertices of a triangle and its orthocentre are each equal to the circle circumscribed about the triangle.*

26.  $D$ ,  $E$  are taken on the circumference of a semicircle described on a given straight line  $AB$ : the chords  $AD$ ,  $BE$  and  $AE$ ,  $BD$  intersect (produced if necessary) at  $F$  and  $G$ : shew that  $FG$  is perpendicular to  $AB$ .

27.  $ABCD$  is a parallelogram;  $AE$  and  $CE$  are drawn at right angles to  $AB$ , and  $CB$  respectively: shew that  $ED$ , if produced, will be perpendicular to  $AC$ .

28.  $ABC$  is a triangle,  $O$  is its orthocentre, and  $AK$  a diameter of the circumscribed circle: shew that  $BOCK$  is a parallelogram.

29. The orthocentre of a triangle is joined to the middle point of the base, and the joining line is produced to meet the circumscribed circle: prove that it will meet it at the same point as the diameter which passes through the vertex.

30. The perpendicular from the vertex of a triangle on the base, and the straight line joining the orthocentre to the middle point of the base, are produced to meet the circumscribed circle at  $P$  and  $Q$ : shew that  $PQ$  is parallel to the base.

31. *The distance of each vertex of a triangle from the orthocentre is double of the perpendicular drawn from the centre of the circumscribed circle on the opposite side.*

32. Three circles are described each passing through the orthocentre of a triangle and two of its vertices: shew that the triangle formed by joining their centres is equal in all respects to the original triangle.

33.  $ABC$  is a triangle inscribed in a circle, and the bisectors of its angles which intersect at  $O$  are produced to meet the circumference in  $PQR$ : shew that  $O$  is the orthocentre of the triangle  $PQR$ .

34. Construct a triangle, having given a vertex, the orthocentre, and the centre of the circumscribed circle.

#### LocI.

35. *Given the base and vertical angle of a triangle, find the locus of its orthocentre.*

Let  $BC$  be the given base, and  $X$  the given angle; and let  $BAC$  be any triangle on the base  $BC$ , having its vertical  $\angle A$  equal to the  $\angle X$ .

Draw the perp<sup>s</sup>  $BE$ ,  $CF$ , intersecting at the orthocentre  $O$ .

*It is required to find the locus of  $O$ .*

Since the  $\angle^s$   $OFA$ ,  $OEA$  are rt. angles,

$\therefore$  the points  $O$ ,  $F$ ,  $A$ ,  $E$  are concyclic;

$\therefore$  the  $\angle$   $FOE$  is the supplement of the  $\angle A$ :


$\therefore$  the vert. opp.  $\angle$   $BOC$  is the supplement of the  $\angle A$ .

But the  $\angle A$  is constant, being always equal to the  $\angle X$ ;

$\therefore$  its supplement is constant;

that is, the  $\triangle BOC$  has a fixed base, and constant vertical angle; hence the locus of its vertex  $O$  is the arc of a segment of which  $BC$  is the chord.

[See Corollary p. 201.]




III. 22.



36. Given the base and vertical angle of a triangle, find the locus of the intersection of the bisectors of its angles.

Let  $BAC$  be any triangle on the given base  $BC$ , having its vertical angle equal to the given  $\angle X$ ; and let  $AI, BI, CI$  be the bisectors of its angles. [See Ex. 2, p. 111.] It is required to find the locus of the point  $I$ .



Denote the angles of the  $\triangle ABC$  by  $A, B, C$ ; and let the  $\angle BIC$  be denoted by  $l$ .

Then from the  $\triangle BIC$ ,

$$(i) \quad l + \frac{1}{2}B + \frac{1}{2}C = \text{two rt. angles}, \quad \text{I. 32.}$$

and from the  $\triangle ABC$ ,

$$A + B + C = \text{two rt. angles}; \quad \text{I. 32.}$$

$$(ii) \quad \text{so that } \frac{1}{2}A + \frac{1}{2}B + \frac{1}{2}C = \text{one rt. angle},$$

$\therefore$ , taking the differences of the equals in (i) and (ii),

$$l - \frac{1}{2}A = \text{one rt. angle};$$

or,

$$l = \text{one rt. angle} + \frac{1}{2}A.$$


But  $A$  is constant, being always equal to the  $\angle X$ ;

$\therefore l$  is constant:

$\therefore$  the locus of  $l$  is the arc of a segment on the fixed chord  $BC$ .

37. Given the base and vertical angle of a triangle, find the locus of the centroid, that is, the intersection of the medians.

Let  $BAC$  be any triangle on the given base  $BC$ , having its vertical angle equal to the given angle  $S$ ; let the medians  $AX, BY, CZ$  intersect at the centroid  $G$ . [See Ex. 4, p. 113.]



It is required to find the locus of the point  $G$ .

Through  $G$  draw  $GP, GQ$  par<sup>l</sup> to  $AB$  and  $AC$  respectively.

Then  $ZG$  is a third part of  $ZC$ ;

Ex. 4, p. 113,

and since  $GP$  is par<sup>l</sup> to  $ZB$ ,

$\therefore BP$  is a third part of  $BC$ . Ex. 19, p. 107.

Similarly  $QC$  is a third part of  $BC$ ;

$\therefore P$  and  $Q$  are fixed points.

Now since  $PG, GQ$  are par<sup>l</sup> respectively to  $BA, AC$ , Constr.

$\therefore$  the  $\angle PGQ =$  the  $\angle BAC$ , I. 29.

$=$  the  $\angle S$ ,

that is, the  $\angle PGQ$  is constant;

$\therefore$  the locus of  $G$  is the arc of a segment on the fixed chord  $PQ$ .

NOTE. In this problem the points  $A$  and  $G$  move on the arcs of similar segments.

38. Given the base and the vertical angle of a triangle; find the locus of the intersection of the bisectors of the exterior base angles.

39. Through the extremities of a given straight line  $AB$  any two parallel straight lines  $AP$ ,  $BQ$  are drawn; find the locus of the intersection of the bisectors of the angles  $PAB$ ,  $QBA$ .

40. Find the locus of the middle points of chords of a circle drawn through a fixed point.

Distinguish between the cases when the given point is within, on, or without the circumference.

41. Find the locus of the points of contact of tangents drawn from a fixed point to a system of concentric circles.

42. Find the locus of the intersection of straight lines which pass through two fixed points on a circle and intercept on its circumference an arc of constant length.

43.  $A$  and  $B$  are two fixed points on the circumference of a circle, and  $PQ$  is any diameter: find the locus of the intersection of  $PA$  and  $QB$ .

44.  $BAC$  is any triangle described on the fixed base  $BC$  and having a constant vertical angle; and  $BA$  is produced to  $P$ , so that  $BP$  is equal to the sum of the sides containing the vertical angle: find the locus of  $P$ .

45.  $AB$  is a fixed chord of a circle, and  $AC$  is a moveable chord passing through  $A$ : if the parallelogram  $CB$  is completed, find the locus of the intersection of its diagonals.

46. A straight rod  $PQ$  slides between two rulers placed at right angles to one another, and from its extremities  $PX$ ,  $QX$  are drawn perpendicular to the rulers: find the locus of  $X$ .

47. Two circles whose centres are  $C$  and  $D$ , intersect at  $A$  and  $B$ : through  $A$ , any straight line  $PAQ$  is drawn terminated by the circumferences; and  $PC$ ,  $QD$  intersect at  $X$ : find the locus of  $X$ , and shew that it passes through  $B$ . [Ex. 9, p. 234.]

48. Two circles intersect at  $A$  and  $B$ , and through  $P$ , any point on the circumference of one of them, two straight lines  $PA$ ,  $PB$  are drawn, and produced if necessary, to cut the other circle at  $X$  and  $Y$ : find the locus of the intersection of  $AY$  and  $BX$ .

49. Two circles intersect at  $A$  and  $B$ ;  $HAK$  is a fixed straight line drawn through  $A$  and terminated by the circumferences, and  $PAQ$  is any other straight line similarly drawn: find the locus of the intersection of  $HP$  and  $QK$ .

50. Two segments of circles are on the same chord  $AB$  and on the same side of it; and  $P$  and  $Q$  are any points one on each arc: find the locus of the intersection of the bisectors of the angles  $PAQ$ ,  $PBQ$ .

51. Two circles intersect at  $A$  and  $B$ ; and through  $A$  any straight line  $PAQ$  is drawn terminated by the circumferences: find the locus of the middle point of  $PQ$ .

#### MISCELLANEOUS EXAMPLES ON ANGLES IN A CIRCLE.

52.  $ABC$  is a triangle, and circles are drawn through  $B$ ,  $C$ , cutting the sides in  $P$ ,  $Q$ ,  $P'$ ,  $Q'$ , ...: shew that  $PQ$ ,  $P'Q'$  ... are parallel to one another and to the tangent drawn at  $A$  to the circle circumscribed about the triangle.

53. Two circles intersect at  $B$  and  $C$ , and from any point  $A$ , on the circumference of one of them,  $AB$ ,  $AC$  are drawn, and produced if necessary, to meet the other at  $D$  and  $E$ : shew that  $DE$  is parallel to the tangent at  $A$ .

54. A secant  $PAB$  and a tangent  $PT$  are drawn to a circle from an external point  $P$ ; and the bisector of the angle  $ATB$  meets  $AB$  at  $C$ : shew that  $PC$  is equal to  $PT$ .

55. From a point  $A$  on the circumference of a circle two chords  $AB$ ,  $AC$  are drawn, and also the diameter  $AF$ : if  $AB$ ,  $AC$  are produced to meet the tangent at  $F$  in  $D$  and  $E$ , shew that the triangles  $ABC$ ,  $AED$  are equiangular to one another.

56.  $O$  is any point within a triangle  $ABC$ , and  $OD$ ,  $OE$ ,  $OF$  are drawn perpendicular to  $BC$ ,  $CA$ ,  $AB$  respectively: shew that the angle  $BOC$  is equal to the sum of the angles  $BAC$ ,  $EDF$ .

57. If two tangents are drawn to a circle from an external point, shew that they contain an angle equal to the difference of the angles in the segments cut off by the chord of contact.

58. Two circles intersect, and through a point of section a straight line is drawn bisecting the angle between the diameters through that point: shew that this straight line cuts off similar segments from the two circles.

59. Two equal circles intersect at  $A$  and  $B$ ; and from centre  $A$ , with any radius less than  $AB$  a third circle is described cutting the given circles on the same side of  $AB$  at  $C$  and  $D$ : shew that the points  $B$ ,  $C$ ,  $D$  are collinear.

60.  $ABC$  and  $A'B'C'$  are two triangles inscribed in a circle, so that  $AB$ ,  $AC$  are respectively parallel to  $A'B'$ ,  $A'C'$ : shew that  $BC'$  is parallel to  $B'C$ .

61. Two circles intersect at  $A$  and  $B$ , and through  $A$  two straight lines  $HAK$ ,  $PAQ$  are drawn terminated by the circumferences: if  $HP$  and  $KQ$  intersect at  $X$ , shew that the points  $H$ ,  $B$ ,  $K$ ,  $X$  are concyclic.

62. Describe a circle touching a given straight line at a given point, so that tangents drawn to it from two fixed points in the given line may be parallel. [See Ex. 10, p. 197.]

63.  $C$  is the centre of a circle, and  $CA$ ,  $CB$  two fixed radii: if from any point  $P$  on the arc  $AB$  perpendiculars  $PX$ ,  $PY$  are drawn to  $CA$  and  $CB$ , shew that the distance  $XY$  is constant.

64.  $AB$  is a chord of a circle, and  $P$  any point in its circumference;  $PM$  is drawn perpendicular to  $AB$ , and  $AN$  is drawn perpendicular to the tangent at  $P$ : shew that  $MN$  is parallel to  $PB$ .

65.  $P$  is any point on the circumference of a circle of which  $AB$  is a fixed diameter, and  $PN$  is drawn perpendicular to  $AB$ ; on  $AN$  and  $BN$  as diameters circles are described, which are cut by  $AP$ ,  $BP$  at  $X$  and  $Y$ : shew that  $XY$  is a common tangent to these circles.

66. Upon the same chord and on the same side of it three segments of circles are described containing respectively a given angle, its supplement and a right angle: shew that the intercept made by the two former segments upon any straight line drawn through an extremity of the given chord is bisected by the latter segment.

67. Two straight lines of indefinite length touch a given circle, and any chord is drawn so as to be bisected by the chord of contact: if the former chord is produced, shew that the intercepts between the circumference and the tangents are equal.

68. Two circles intersect one another: through one of the points of section draw a straight line of given length terminated by the circumferences.

69. On the three sides of any triangle equilateral triangles are described remote from the given triangle: shew that the circles described about them intersect at a point.

70. On  $BC$ ,  $CA$ ,  $AB$  the sides of a triangle  $ABC$ , any points  $P$ ,  $Q$ ,  $R$  are taken; shew that the circles described about the triangles  $AQR$ ,  $BRP$ ,  $CPQ$  meet in a point.

71. Find a point within a triangle at which the sides subtend equal angles.

72. Describe an equilateral triangle so that its sides may pass through three given points.

73. Describe a triangle equal in all respects to a given triangle, and having its sides passing through three given points.


## SIMSON'S LINE.

74. *If from any point on the circumference of the circle circumscribed about a triangle, perpendiculars are drawn to the three sides, the feet of these perpendiculars are collinear.*

Let  $P$  be any point on the  $\bigcirc^\infty$  of the circle circumscribed about the  $\triangle ABC$ ; and let  $PD$ ,  $PE$ ,  $PF$  be the perp<sup>s</sup> drawn from  $P$  to the three sides.

*It is required to prove that the points  $D$ ,  $E$ ,  $F$  are collinear.*

Join  $FD$  and  $DE$  :  
then  $FD$  and  $DE$  shall be in the same st. line.



Join  $PB$ ,  $PC$ .

Because the  $\angle^s$   $PDB$ ,  $PFB$  are rt. angles, *Hyp.*

$\therefore$  the points  $P$ ,  $D$ ,  $B$ ,  $F$  are concyclic :

$\therefore$  the  $\angle$   $PDF =$  the  $\angle$   $PBF$ , in the same segment. III. 21.

But since  $BACP$  is a quad<sup>l</sup> inscribed in a circle, having one of its sides  $AB$  produced to  $F$ ,

$\therefore$  the ext.  $\angle$   $PBF =$  the opp. int.  $\angle$   $ACP$ . *Ex. 3, p. 202.*

$\therefore$  the  $\angle$   $PDF =$  the  $\angle$   $ACP$ .

To each add the  $\angle$   $PDE$  :

then the  $\angle^s$   $PDF$ ,  $PDE =$  the  $\angle^s$   $ECP$ ,  $PDE$ .

But since the  $\angle^s$   $PDC$ ,  $PEC$  are rt. angles,

$\therefore$  the points  $P$ ,  $D$ ,  $E$ ,  $C$  are concyclic ;

$\therefore$  the  $\angle^s$   $ECP$ ,  $PDE$  together = two rt. angles :

$\therefore$  the  $\angle^s$   $PDF$ ,  $PDE$  together = two rt. angles ;

$\therefore$   $FD$  and  $DE$  are in the same st. line ;

I. 14.

that is, the points  $D$ ,  $E$ ,  $F$  are collinear.

Q.E.D.

[The line  $FDE$  is called the *Pedal* or *Simson's Line* of the triangle  $ABC$  for the point  $P$ ; though the tradition attributing the theorem to Robert Simson has been recently shaken by the researches of Dr. J. S. Mackay.]

75.  $ABC$  is a triangle inscribed in a circle; and from any point  $P$  on the circumference  $PD$ ,  $PF$  are drawn perpendicular to  $BC$  and  $AB$ : if  $FD$ , or  $FD$  produced, cuts  $AC$  at  $E$ , shew that  $PE$  is perpendicular to  $AC$ .

76. Find the locus of a point which moves so that if perpendiculars are drawn from it to the sides of a given triangle, their feet are collinear.

77.  $ABC$  and  $AB'C'$  are two triangles having a common vertical angle, and the circles circumscribed about them meet again at  $P$ ; shew that the feet of perpendiculars drawn from  $P$  to the four lines  $AB$ ,  $AC$ ,  $BC$ ,  $B'C'$  are collinear.


78. A triangle is inscribed in a circle, and any point P on the circumference is joined to the orthocentre of the triangle: shew that this joining line is bisected by the pedal of the point P.

IV. ON THE CIRCLE IN CONNECTION WITH RECTANGLES.

[See Propositions 35, 36, 37.]

1. If from any external point P two tangents are drawn to a given circle whose centre is O, and if OP meets the chord of contact at Q; then the rectangle OP, OQ is equal to the square on the radius.

Let PH, PK be tangents, drawn from the external point P to the  $\odot$  HAK, whose centre is O; and let OP meet HK the chord of contact at Q, and the  $\odot^{\infty}$  at A. Then shall the rect. OP, OQ = the sq. on OA.



On HP as diameter describe a circle: this circle must pass through Q, since the  $\angle$  HQP is a rt. angle. III. 31.

Join OH.

Then since PH is a tangent to the  $\odot$  HAK,  $\therefore$  the  $\angle$  OHP is a rt. angle.

And since HP is a diameter of the  $\odot$  HQP,

$\therefore$  OH touches the  $\odot$  HQP at H. III. 16.

$\therefore$  the rect. OP, OQ = the sq. on OH, III. 36.  
= the sq. on OA. Q.E.D.

2. ABC is a triangle, and AD, BE, CF the perpendiculars drawn from the vertices to the opposite sides, meeting in the orthocentre O: shew that the rect. AO, OD = the rect. BO, OE = the rect. CO, OF.

3. ABC is a triangle, and AD, BE the perpendiculars drawn from A and B on the opposite sides: shew that the rectangle CA, CE is equal to the rectangle CB, CD.

4. ABC is a triangle right-angled at C, and from D, any point in the hypotenuse AB, a straight line DE is drawn perpendicular to AB and meeting BC at E: shew that the square on DE is equal to the difference of the rectangles AD, DB and CE, EB.

5. From an external point P two tangents are drawn to a given circle whose centre is O, and OP meets the chord of contact at Q: shew that any circle which passes through the points P, Q will cut the given circle orthogonally. [See Def. p. 240.]

6. *A series of circles pass through two given points, and from a fixed point in the common chord produced tangents are drawn to all the circles: shew that the points of contact lie on a circle which cuts all the given circles orthogonally.*

7. *All circles which pass through a fixed point, and cut a given circle orthogonally, pass also through a second fixed point.*

8. Find the locus of the centres of all circles which pass through a given point and cut a given circle orthogonally.

9. Describe a circle to pass through two given points and cut a given circle orthogonally.

10. A, B, C, D are four points taken in order on a given straight line: find a point O between B and C such that the rectangle OA, OB may be equal to the rectangle OC, OD.

11. *AB is a fixed diameter of a circle, and CD a fixed straight line of indefinite length cutting AB or AB produced at right angles; any straight line is drawn through A to cut CD at P and the circle at Q: shew that the rectangle AP, AQ is constant.*

12. AB is a fixed diameter of a circle, and CD a fixed chord at right angles to AB; any straight line is drawn through A to cut CD at P and the circle at Q: shew that the rectangle AP, AQ is equal to the square on AC.

13. *A is a fixed point, and CD a fixed straight line of indefinite length; AP is any straight line drawn through A to meet CD at P; and in AP a point Q is taken such that the rectangle AP, AQ is constant: find the locus of Q.*

14. Two circles intersect orthogonally, and tangents are drawn from any point on the circumference of one to touch the other: prove that the first circle passes through the middle point of the chord of contact of the tangents. [Ex. I, p. 251.]

15. A semicircle is described on AB as diameter, and any two chords AC, BD are drawn intersecting at P: shew that

$$AB^2 = AC \cdot AP + BD \cdot BP.$$

16. Two circles intersect at B and C, and the two direct common tangents AE and DF are drawn: if the common chord is produced to meet the tangents at G and H, shew that  $GH^2 = AE^2 + BC^2$ .

17. If from a point P, without a circle, PM is drawn perpendicular to a diameter AB, and also a secant PCD, shew that

$$PM^2 = PC \cdot PD + AM \cdot MB.$$

18. Three circles intersect at D, and their other points of intersection are A, B, C; AD cuts the circle BDC at E, and EB, EC cut the circles ADB, ADC respectively at F and G: show that the points F, A, G are collinear, and F, B, C, G concyclic.

19. A semicircle is described on a given diameter BC, and from B and C any two chords BE, CF are drawn intersecting within the semicircle at O; BF and CE are produced to meet at A: shew that the sum of the squares on AB, AC is equal to twice the square on the tangent from A together with the square on BC.

20. X and Y are two fixed points in the diameter of a circle equidistant from the centre C: through X any chord PXQ is drawn, and its extremities are joined to Y; shew that the sum of the squares on the sides of the triangle PYQ is constant. [See p. 161, Ex. 24.]

## PROBLEMS ON TANGENCY.

21. To describe a circle to pass through two given points and to touch a given straight line.

Let A and B be the given points, and CD the given st. line.

It is required to describe a circle to pass through A and B and to touch CD.

Join BA, and produce it to meet CD at P.

Describe a square equal to the rect. PA, PB;


and from PD (or PC) cut off PQ equal to a side of this square. II. 14.

Through A, B, and Q describe a circle. Ex. 4, p. 171.

Then since the rect. PA, PB = the sq. on PQ,

$\therefore$  the  $\odot$  ABQ touches CD at Q.

III. 37.  
Q.E.F.



NOTES. (i) Since PQ may be taken on either side of P, it is clear that there are in general two solutions of the problem.

(ii) When AB is parallel to the given line CD, the above method is not applicable. In this case a simple construction follows from III. 1, Cor. and III. 16, and it will be found that only one solution exists.



22. To describe a circle to pass through two given points and to touch a given circle.

Let  $A$  and  $B$  be the given points, and  $\odot CRP$  the given circle.

It is required to describe a circle to pass through  $A$  and  $B$ , and to touch the  $\odot CRP$ .

Through  $A$  and  $B$  describe any circle to cut the given circle at  $P$  and  $Q$ .

Join  $AB$ ,  $PQ$ , and produce them to meet at  $D$ .

From  $D$  draw  $DC$  to touch the given circle, and let  $C$  be the point of contact.

Then the circle described through  $A, B, C$  will touch the given circle.


For, from the  $\odot ABQP$ , the rect.  $DA, DB =$  the rect.  $DP, DQ$ ;  
and from the  $\odot PQC$ , the rect.  $DP, DQ =$  the sq. on  $DC$ ; III. 36.

$\therefore$  the rect.  $DA, DB =$  the sq. on  $DC$ ;

$\therefore DC$  touches the  $\odot ABC$  at  $C$ ; III. 37.

But  $DC$  touches the  $\odot PQC$  at  $C$ ; Constr.

$\therefore$  the  $\odot ABC$  touches the given circle, and it passes through the given points  $A$  and  $B$ . Q. E. F.



NOTE. (i) Since two tangents may be drawn from  $D$  to the given circle, it follows that there will be two solutions of the problem.

(ii) The general construction fails when the straight line bisecting  $AB$  at right angles passes through the centre of the given circle: the problem then becomes symmetrical, and the solution is obvious.

23. To describe a circle to pass through a given point and to touch two given straight lines.

Let  $P$  be the given point, and  $AB, AC$  the given straight lines.


It is required to describe a circle to pass through  $P$  and to touch  $AB, AC$ .

Now the centre of every circle which touches  $AB$  and  $AC$  must lie on the bisector of the  $\angle BAC$ .

Ex. 7, p. 197.

Hence draw  $AE$  bisecting the  $\angle BAC$ .

From  $P$  draw  $PK$  perp. to  $AE$ , and produce it to  $P'$ , making  $KP'$  equal to  $PK$ .



Then every circle which has its centre in AE, and passes through P, must also pass through P'. Ex. 1, p. 233.

Hence the problem is now reduced to drawing a circle through P and P' to touch either AC or AB. Ex. 21, p. 253.

Produce P'P to meet AC at S.

Describe a square equal to the rect. SP, SP'; and cut off SR equal to a side of the square. II. 14.

Describe a circle through the points P', P, R.

Then since the rect. SP, SP' = the sq. on SR, Constr. III. 37,

$\therefore$  this circle touches AC at R;

and since its centre is in AE, the bisector of the  $\angle$  BAC, it may be shewn also to touch AB. Q.E.F.

NOTES. (i) Since SR may be taken on either side of S, it follows that there will be two solutions of the problem.


(ii) If the given straight lines are parallel, the centre lies on the parallel straight line mid-way between them, and the construction proceeds as before.

24. To describe a circle to touch two given straight lines and a given circle.

Let AB, AC be the two given st. lines, and D the centre of the given circle.

It is required to describe a circle to touch AB, AC and the circle whose centre is D.

Draw EF, GH par<sup>l</sup> to AB and AC respectively, on the sides remote from D, and at distances from them equal to the radius of the given circle.



Describe the  $\odot$  MND to touch EF and GH at M and N, and to pass through D. Ex. 23, p. 254.

Let O be the centre of this circle.

Join OM, ON, OD meeting AB, AC, and the given circle at P, Q, and R.

Then a circle described with centre O and radius OP will touch AB, AC and the given circle.

For since O is the centre of the  $\odot$  MND,

$\therefore$  OM = ON = OD. Constr.

But PM = QN = RD;

$\therefore$  OP = OQ = OR.

$\therefore$  a circle described with centre O, and radius OP, will pass through Q and R.

And since the  $\angle^s$  at M and N are rt. angles, III. 18.

$\therefore$  the  $\angle^s$  at P and Q are rt. angles; I. 29.

$\therefore$  the  $\odot$  PQR touches AB and AC.

And since R, the point in which the circles meet, is on the line of centres OD,

$\therefore$  the  $\odot$  PQR touches the given circle. Q.E.F.


NOTE. There will be two solutions of this problem, since two circles may be drawn to touch EF, GH and to pass through D.

25. To describe a circle to pass through a given point and touch a given straight line and a given circle.

Let P be the given point, AB the given st. line, and DHE the given circle, of which C is the centre.

It is required to describe a circle to pass through P, and to touch AB and the  $\odot$  DHE.

Through C draw DCEF perp. to AB, cutting the circle at the points D and E, of which E is between C and AB.



Join DP;

and by describing a circle through F, E, and P, find a point K in DP (or DP produced) such that the rect. DE, DF = the rect. DK, DP.

Describe a circle to pass through P, K, and touch AB: Ex. 21, p. 253. This circle shall also touch the given  $\odot$  DHE.

For let G be the point at which this circle touches AB. Join DG, cutting the given circle DHE at H.

Join HE.

Then the  $\angle$  DHE is a rt. angle, being in a semicircle, III. 31.  
also the angle at F is a rt. angle; Constr.

$\therefore$  the points E, F, G, H are concyclic:

$\therefore$  the rect. DE, DF = the rect. DH, DG: III. 36.  
but the rect. DE, DF = the rect. DK, DP: Constr.

$\therefore$  the rect. DH, DG = the rect. DK, DP:

$\therefore$  the point H is on the  $\odot$  PKG.

Let O be the centre of the  $\odot$  PHG.

Join OG, OH, CH.

Then OG and DF are par<sup>l</sup>, since they are both perp. to AB; and DG meets them.

$\therefore$  the  $\angle$  OGD = the  $\angle$  GDC. I. 29.

But since OG = OH, and CD = CH,

$\therefore$  the  $\angle$  OGH = the  $\angle$  OHG; and the  $\angle$  CDH = the  $\angle$  CHD:

$\therefore$  the  $\angle$  OHG = the  $\angle$  CHD;

$\therefore$  OH and CH are in one st. line.

$\therefore$  the  $\odot$  PHG touches the given  $\odot$  DHE. Q.E.F.

NOTES. (i) Since two circles may be drawn to pass through P, K and to touch AB, it follows that there will be two solutions of the present problem.

(ii) Two more solutions may be obtained by joining PE, and proceeding as before.

The student should examine the nature of the contact between the circles in each case.

26. Describe a circle to pass through a given point, to touch a given straight line, and to have its centre on another given straight line.

27. Describe a circle to pass through a given point, to touch a given circle, and to have its centre on a given straight line.

28. Describe a circle to pass through two given points, and to intercept an arc of given length on a given circle.

29. Describe a circle to touch a given circle and a given straight line at a given point.

30. Describe a circle to touch two given circles and a given straight line.

## V. ON MAXIMA AND MINIMA.

We gather from the Theory of Loci that the position of an angle, line or figure is capable under suitable conditions of gradual change; and it is usually found that change of *position* involves a corresponding and gradual change of *magnitude*.

Under these circumstances we may be required to note if any situations exist at which the magnitude in question, after increasing, begins to decrease; or after decreasing, to increase: in such situations the magnitude is said to have reached a **Maximum** or a **Minimum** value; for in the former case it is greater, and in the latter case less than in adjacent situations on either side. In the geometry of the circle and straight line we only meet with such cases of continuous change as admit of *one* transition from an increasing to a decreasing state—or vice versâ—so that in all the problems with which we have to deal (where a single circle is involved) there can be only one Maximum and one Minimum—the Maximum being the greatest, and the Minimum being the least value that the variable magnitude is capable of taking.

Thus a variable geometrical magnitude reaches its maximum or minimum value at a *turning point*, towards which the magnitude may mount or descend from either side: it is natural therefore to expect a maximum or minimum value to occur when, in the course of its change, the magnitude assumes a *symmetrical* form or position; and this is usually found to be the case.

This general connection between a symmetrical form or position and a maximum or minimum value is not exact enough to constitute a *proof* in any particular problem; but by means of it a situation is suggested, which on further examination may be shewn to give the maximum or minimum value sought for.

For example, suppose it is required

*to determine the greatest straight line that may be drawn perpendicular to the chord of a segment of a circle and intercepted between the chord and the arc:*

we immediately anticipate that the greatest perpendicular is that which occupies a *symmetrical* position in the figure, namely the perpendicular which passes through the middle point of the chord; and on further examination this may be proved to be the case by means of I. 19, and I. 34.

Again we are able to find at what point a geometrical magnitude, varying under certain conditions, assumes its Maximum or Minimum value, if we can discover a construction for drawing the magnitude so that it may have an *assigned* value: for we may then examine between what limits the assigned value must lie in order that the construction may be possible; and the higher or lower limit will give the Maximum or Minimum sought for.

It was pointed out in the chapter on the Intersection of Loci, [see page 125] that if under certain conditions existing among the data, *two* solutions of a problem are possible, and under other conditions, *no* solution exists, there will always be some intermediate condition under which *one* and *only one* distinct solution is possible.


Under these circumstances this single or limiting solution will always be found to correspond to the maximum or minimum value of the magnitude to be constructed.

1. For example, suppose it is required

*to divide a given straight line so that the rectangle contained by the two segments may be a maximum.*

We may first attempt to divide the given straight line so that the rectangle contained by its segments may have a *given* area—that is, be equal to the square on a given straight line.

Let  $AB$  be the given straight line, and  $K$  the side of the given square.



It is required to divide the st. line  $AB$  at a point  $M$ , so that the rect.  $AM, MB$  may be equal to the sq. on  $K$ .

Adopting a construction suggested by II. 14,

describe a semicircle on  $AB$ ; and at any point  $X$  in  $AB$ , or  $AB$  produced, draw  $XY$  perp. to  $AB$ , and equal to  $K$ .

Through  $Y$  draw  $YZ$  par<sup>l</sup> to  $AB$ , to meet the arc of the semicircle at  $P$ .

Then if the perp.  $PM$  is drawn to  $AB$ , it may be shewn after the manner of II. 14, or by III. 35 that

$$\begin{aligned} \text{the rect. } AM, MB &= \text{the sq. on } PM \\ &= \text{the sq. on } K. \end{aligned}$$

So that the rectangle  $AM, MB$  increases as  $K$  increases.

Now if  $K$  is less than the radius  $CD$ , then  $YZ$  will meet the arc of the semicircle in two points  $P, P'$ ; and it follows that  $AB$  may be divided at *two* points, so that the rectangle contained by its segments may be equal to the square on  $K$ . If  $K$  increases, the st. line  $YZ$  will recede from  $AB$ , and the points of intersection  $P, P'$  will continually approach one another; until, when  $K$  is equal to the radius  $CD$ , the st. line  $YZ$  (now in the position  $Y'Z'$ ) will meet the arc in *two coincident points*, that is, will touch the semicircle at  $D$ ; and there will be only *one* solution of the problem.

If  $K$  is greater than  $CD$ , the straight line  $YZ$  will not meet the semicircle, and the problem is impossible.

Hence the greatest length that  $K$  may have, in order that the construction may be possible, is the radius  $CD$ .

$\therefore$  the rect.  $AM, MB$  is a maximum, when it is equal to the square on  $CD$ ;

that is, when  $PM$  coincides with  $CD$ , and consequently when  $M$  is the middle point of  $AB$ .

**NOTE.** The special feature to be noticed in this problem is that the maximum is found at the transitional point between *two* solutions and *no* solution; that is, when the two solutions coincide and become identical.

The following example illustrates the same point.

2. *To find at what point in a given straight line the angle subtended by the line joining two given points, which are on the same side of the given straight line, is a maximum.*

Let  $CD$  be the given st. line, and  $A, B$  the given points on the same side of  $CD$ .

*It is required to find at what point in  $CD$  the angle subtended by the st. line  $AB$  is a maximum.*

First determine at what point in  $CD$ , the st. line  $AB$  subtends a given angle.

This is done as follows:—

On  $AB$  describe a segment of a circle containing an angle equal to the given angle. III. 33.

If the arc of this segment intersects  $CD$ , two points in  $CD$  are found at which  $AB$  subtends the given angle: but if the arc does not meet  $CD$ , no solution is given.

In accordance with the principles explained above, we expect that a maximum angle is determined at the limiting position; that is, when the arc touches  $CD$ , or meets it at two coincident points.

[See page 231.]

This we may prove to be the case.

Describe a circle to pass through  $A$  and  $B$ , and to touch the st. line  $CD$ .

[Ex. 21, p. 253.]

Let  $P$  be the point of contact.

Then shall the  $\angle APB$  be greater than any other angle subtended by  $AB$  at a point in  $CD$  on the same side of  $AB$  as  $P$ .

For take  $Q$ , any other point in  $CD$ , on the same side of  $AB$  as  $P$ ;

and join  $AQ, QB$ .

Since  $Q$  is a point in the tangent other than the point of contact, it must be without the circle;

$\therefore$  either  $BQ$  or  $AQ$  must meet the arc of the segment  $APB$ .


Let  $BQ$  meet the arc at  $K$ : join  $AK$ .

Then the  $\angle APB = \text{the } \angle AKB$ , in the same segment:  
but the ext.  $\angle AKB$  is greater than the int. opp.  $\angle AQB$ .

$\therefore$  the  $\angle APB$  is greater than  $\angle AQB$ .

Similarly the  $\angle APB$  may be shewn to be greater than any other angle subtended by  $AB$  at a point in  $CD$  on the same side of  $AB$ :  
that is, the  $\angle APB$  is the greatest of all such angles. Q.E.D.

NOTE. Two circles may be described to pass through  $A$  and  $B$ , and to touch  $CD$ , the points of contact being on opposite sides of  $AB$ ;



hence two points in CD may be found such that the angle subtended by AB at each of them is greater than the angle subtended at any other point in CD on the same side of AB.

We add two more examples of considerable importance.


3. In a straight line of indefinite length find a point such that the sum of its distances from two given points, on the same side of the given line, shall be a minimum.

Let CD be the given st. line of indefinite length, and A, B the given points on the same side of CD.

It is required to find a point P in CD, such that the sum of AP, PB is a minimum.

Draw AF perp. to CD; and produce AF to E, making FE equal to AF.

Join EB, cutting CD at P.  
Join AP, PB.



Then of all lines drawn from A and B to a point in CD, the sum of AP, PB shall be the least.

For, let Q be any other point in CD.  
Join AQ, BQ, EQ.

Now in the  $\triangle$ 's AFP, EFP,  
Because  $\left\{ \begin{array}{l} AF = EF, \\ \text{and FP is common;} \\ \text{and the } \angle AFP = \text{the } \angle EFP, \end{array} \right.$  being rt. angles. Constr.  
 $\therefore AP = EP.$  I. 4.

Similarly it may be shewn that  
 $AQ = EQ.$

Now in the  $\triangle$  EQB, the two sides EQ, QB are together greater than EB;

hence, AQ, QB are together greater than EB,  
that is, greater than AP, PB.

Similarly the sum of the st. lines drawn from A and B to any other point in CD may be shewn to be greater than AP, PB.

$\therefore$  the sum of AP, PB is a minimum.

Q. E. D.

NOTE. It follows from the above proof that

the  $\angle APF = \text{the } \angle EPF$  I. 4.  
= the  $\angle BPD.$  I. 15.

Thus the sum of AP, PB is a minimum, when these lines are equally inclined to CD.



4. Given two intersecting straight lines AB, AC, and a point P between them; shew that of all straight lines which pass through P and are terminated by AB, AC, that which is bisected at P cuts off the triangle of minimum area.

Let EF be the st. line, terminated by AB, AC, which is bisected at P.

Then the  $\triangle$  FAE shall be of minimum area.

For let HK be any other st. line passing through P.


Through E draw EM par<sup>l</sup> to AC.

Then in the  $\triangle$ <sup>s</sup> HPF, MPE,

Because {	the $\angle$ HPF = the $\angle$ MPE,	I. 15.
	and the $\angle$ HFP = the $\angle$ MEP,	I. 29.
	and FP = EP;	<i>Hyp.</i>
	$\therefore$ the $\triangle$ HPF = the $\triangle$ MPE.	I. 26, Cor.

But the  $\triangle$  MPE is less than the  $\triangle$  KPE;  
 $\therefore$  the  $\triangle$  HPF is less than the  $\triangle$  KPE;  
 to each add the fig. AHPE;  
 then the  $\triangle$  FAE is less than the  $\triangle$  HAK.

Similarly it may be shewn that the  $\triangle$  FAE is less than any other triangle formed by drawing a st. line through P:  
 that is, the  $\triangle$  FAE is a minimum.



#### EXAMPLES.

1. Two sides of a triangle are given in length; how must they be placed in order that the area of the triangle may be a maximum?
2. Of all triangles of given base and area, the isosceles is that which has the least perimeter.
3. Given the base and vertical angle of a triangle; construct it so that its area may be a maximum.
4. Find a point in a given straight line such that the tangents drawn from it to a given circle contain the greatest angle possible.
5. A straight rod slips between two straight rulers placed at right angles to one another; in what position is the triangle intercepted between the rulers and rod a maximum?

6. Divide a given straight line into two parts, so that the sum of the squares on the segments

(i) may be equal to a given square ;

(ii) may be a minimum.

7. Through a point of intersection of two circles draw a straight line terminated by the circumferences,

(i) so that it may be of given length ;

(ii) so that it may be a maximum.

8. Two tangents to a circle cut one another at right angles ; find the point on the intercepted arc such that the sum of the perpendiculars drawn from it to the tangents may be a minimum.

9. Straight lines are drawn from two given points to meet one another on the convex circumference of a given circle : prove that their sum is a minimum when they make equal angles with the tangent at the point of intersection.

10. Of all triangles of given vertical angle and altitude, that which is isosceles has the least area.

11. Two straight lines CA, CB of indefinite length are drawn from the centre of a circle to meet the circumference at A and B ; then of all tangents that may be drawn to the circle at points on the arc AB, that whose intercept is bisected at the point of contact cuts off the triangle of minimum area.

12. Given two intersecting tangents to a circle, draw a tangent to the convex arc so that the triangle formed by it and the given tangents may be of maximum area.

13. Of all triangles of given base and area, that which is isosceles has the greatest vertical angle.

14. Find a point on the circumference of a circle at which the straight line joining two given points (of which both are within, or both without the circle) subtends the greatest angle.

15. A bridge consists of three arches, whose spans are 49 ft., 32 ft. and 49 ft. respectively : shew that the point on either bank of the river at which the middle arch subtends the greatest angle is 63 feet distant from the bridge.

16. From a given point P without a circle whose centre is C, draw a straight line to cut the circumference at A and B, so that the triangle ACB may be of maximum area.

17. Shew that the greatest rectangle which can be inscribed in a circle is a square.

18. A and B are two fixed points without a circle : find a point P on the circumference, such that the sum of the squares on AP, PB may be a minimum. [See p. 161, Ex. 24.]

19. A segment of a circle is described on the chord  $AB$ : find a point  $C$  on its arc so that the sum of  $AC$ ,  $BC$  may be a maximum.

20. *Of all triangles that can be inscribed in a circle that which has the greatest perimeter is equilateral.*

21. *Of all triangles that can be inscribed in a given circle that which has the greatest area is equilateral.*

22. *Of all triangles that can be inscribed in a given triangle that which has the least perimeter is the triangle formed by joining the feet of the perpendiculars drawn from the vertices on opposite sides.*

23. Of all rectangles of given area, the square has the least perimeter.

24. Describe the triangle of maximum area, having its angles equal to those of a given triangle, and its sides passing through three given points.

#### VI. HARDER MISCELLANEOUS EXAMPLES.

1.  $AB$  is a diameter of a given circle; and  $AC$ ,  $BD$ , two chords on the same side of  $AB$ , intersect at  $E$ : shew that the circle which passes through  $D$ ,  $E$ ,  $C$  cuts the given circle orthogonally.

2. Two circles whose centres are  $C$  and  $D$  intersect at  $A$  and  $B$ ; and a straight line  $PAQ$  is drawn through  $A$  and terminated by the circumferences: prove that

$$(i) \text{ the angle } PBQ = \text{the angle } CAD$$

$$(ii) \text{ the angle } BPC = \text{the angle } BQD.$$

3. Two chords  $AB$ ,  $CD$  of a circle whose centre is  $O$  intersect at right angles at  $P$ : shew that

$$(i) PA^2 + PB^2 + PC^2 + PD^2 = 4 \text{ (radius)}^2.$$

$$(ii) AB^2 + CD^2 + 4OP^2 = 8 \text{ (radius)}^2.$$

4. Two parallel tangents to a circle intercept on any third tangent a portion which is so divided at its point of contact that the rectangle contained by its two parts is equal to the square on the radius.

5. Two equal circles move between two straight lines placed at right angles, so that each straight line is touched by one circle, and the two circles touch one another: find the locus of the point of contact.

6.  $AB$  is a given diameter of a circle, and  $CD$  is any parallel chord: if any point  $X$  in  $AB$  is joined to the extremities of  $CD$ , shew that

$$XC^2 + XD^2 = XA^2 + XB^2.$$

7.  $PQ$  is a fixed chord in a circle, and  $PX$ ,  $QY$  any two parallel chords through  $P$  and  $Q$ ; shew that  $XY$  touches a fixed concentric circle.

8. Two equal circles intersect at  $A$  and  $B$ ; and from  $C$ , any point on the circumference of one of them, a perpendicular is drawn to  $AB$ , meeting the other circle at  $O$  and  $O'$ ; shew that either  $O$  or  $O'$  is the orthocentre of the triangle  $ABC$ . Distinguish between the two cases.

9. Three equal circles pass through the same point  $A$ , and their other points of intersection are  $B$ ,  $C$ ,  $D$ : shew that of the four points  $A$ ,  $B$ ,  $C$ ,  $D$ , each is the orthocentre of the triangle formed by joining the other three.

10. From a given point without a circle draw a straight line to the concave circumference so as to be bisected by the convex circumference. When is this problem impossible?

11. Draw a straight line cutting two concentric circles so that the chord intercepted by the circumference of the greater circle may be double of the chord intercepted by the less.

12.  $ABC$  is a triangle inscribed in a circle, and  $A'$ ,  $B'$ ,  $C'$  are the middle points of the arcs subtended by the sides (remote from the opposite vertices): find the relation between the angles of the two triangles  $ABC$ ,  $A'B'C'$ ; and prove that the pedal triangle of  $A'B'C'$  is equiangular to the triangle  $ABC$ .

13. The opposite sides of a quadrilateral inscribed in a circle are produced to meet: shew that the bisectors of the two angles so formed are perpendicular to one another.

14. If a quadrilateral can have one circle inscribed in it, and another circumscribed about it; shew that the straight lines joining the opposite points of contact of the inscribed circle are perpendicular to one another.

15. Given the base of a triangle and the sum of the remaining sides; find the locus of the foot of the perpendicular from one extremity of the base on the bisector of the exterior vertical angle.

16. Two circles touch each other at  $C$ , and straight lines are drawn through  $C$  at right angles to one another, meeting the circles at  $P$ ,  $P'$  and  $Q$ ,  $Q'$  respectively: if the straight line which joins the centres is terminated by the circumferences at  $A$  and  $A'$ , shew that

$$P'P^2 + Q'Q^2 = A'A^2.$$

17. Two circles cut one another orthogonally at  $A$  and  $B$ ;  $P$  is any point on the arc of one circle intercepted by the other, and  $PA$ ,  $PB$  are produced to meet the circumference of the second circle at  $C$  and  $D$ : shew that  $CD$  is a diameter.

18.  $ABC$  is a triangle, and from any point  $P$  perpendiculars  $PD$ ,  $PE$ ,  $PF$  are drawn to the sides: if  $S_1$ ,  $S_2$ ,  $S_3$  are the centres of the circles circumscribed about the triangles  $EPF$ ,  $FPD$ ,  $DPE$ , shew that the triangle  $S_1S_2S_3$  is equiangular to the triangle  $ABC$ , and that the sides of the one are respectively half of the sides of the other.

19. Two tangents  $PA$ ,  $PB$  are drawn from an external point  $P$  to a given circle, and  $C$  is the middle point of the chord of contact  $AB$ ; if  $XY$  is any chord through  $P$ , shew that  $AB$  bisects the angle  $XCY$ .

20. Given the sum of two straight lines and the rectangle contained by them (equal to a given square): find the lines.

21. Given the sum of the squares on two straight lines and the rectangle contained by them: find the lines.

22. Given the sum of two straight lines and the sum of the squares on them: find the lines.

23. Given the difference between two straight lines, and the rectangle contained by them: find the lines.

24. Given the sum or difference of two straight lines and the difference of their squares: find the lines.

25.  $ABC$  is a triangle, and the internal and external bisectors of the angle  $A$  meet  $BC$ , and  $BC$  produced, at  $P$  and  $P'$ : if  $O$  is the middle point of  $PP'$ , shew that  $OA$  is a tangent to the circle circumscribed about the triangle  $ABC$ .

26.  $ABC$  is a triangle, and from  $P$ , any point on the circumference of the circle circumscribed about it, perpendiculars are drawn to the sides  $BC$ ,  $CA$ ,  $AB$  meeting the circle again in  $A'$ ,  $B'$ ,  $C'$ ; prove that

(i) the triangle  $A'B'C'$  is identically equal to the triangle  $ABC$ .

(ii)  $AA'$ ,  $BB'$ ,  $CC'$  are parallel.

27. Two equal circles intersect at fixed points  $A$  and  $B$ , and from any point in  $AB$  a perpendicular is drawn to meet the circumferences on the same side of  $AB$  at  $P$  and  $Q$ : shew that  $PQ$  is of constant length.

28. The straight lines which join the vertices of a triangle to the centre of its circumscribed circle, are perpendicular respectively to the sides of the pedal triangle.

29.  $P$  is any point on the circumference of a circle circumscribed about a triangle  $ABC$ ; and perpendiculars  $PD$ ,  $PE$  are drawn from  $P$  to the sides  $BC$ ,  $CA$ . Find the locus of the centre of the circle circumscribed about the triangle  $PDE$ .

30.  $P$  is any point on the circumference of a circle circumscribed about a triangle  $ABC$ : shew that the angle between Simson's Line for the point  $P$  and the side  $BC$  is equal to the angle between  $AP$  and the diameter of the circumscribed circle through  $A$ .

31. Shew that the circles circumscribed about the four triangles formed by two pairs of intersecting straight lines meet in a point.

32. Shew that the orthocentres of the four triangles formed by two pairs of intersecting straight lines are collinear.

#### ON THE CONSTRUCTION OF TRIANGLES.

33. Given the vertical angle, one of the sides containing it, and the length of the perpendicular from the vertex on the base: construct the triangle.

34. Given the feet of the perpendiculars drawn from the vertices on the opposite sides: construct the triangle.

35. Given the base, the altitude, and the radius of the circumscribed circle: construct the triangle.

36. Given the base, the vertical angle, and the sum of the squares on the sides containing the vertical angle: construct the triangle.

37. Given the base, the altitude and the sum of the squares on the sides containing the vertical angle: construct the triangle.

38. Given the base, the vertical angle, and the difference of the squares on the sides containing the vertical angle: construct the triangle.

39. Given the vertical angle, and the lengths of the two medians drawn from the extremities of the base: construct the triangle.

40. Given the base, the vertical angle, and the difference of the angles at the base: construct the triangle.

41. Given the base, and the position of the bisector of the vertical angle: construct the triangle.

42. Given the base, the vertical angle, and the length of the bisector of the vertical angle: construct the triangle.

43. Given the perpendicular from the vertex on the base, the bisector of the vertical angle, and the median which bisects the base: construct the triangle.

44. Given the bisector of the vertical angle, the median bisecting the base, and the difference of the angles at the base: construct the triangle.

## BOOK IV.

Book IV. consists entirely of problems, dealing with various rectilinear figures in relation to the circles which pass through their angular points, or are touched by their sides.


### DEFINITIONS.

1. A **Polygon** is a rectilinear figure bounded by more than four sides.


A Polygon of	<i>five</i> sides	is called a	<b>Pentagon,</b>
"	<i>six</i> sides	"	<b>Hexagon,</b>
"	<i>seven</i> sides	"	<b>Heptagon,</b>
"	<i>eight</i> sides	"	<b>Octagon,</b>
"	<i>ten</i> sides	"	<b>Decagon,</b>
"	<i>twelve</i> sides	"	<b>Dodecagon,</b>
"	<i>fifteen</i> sides	"	<b>Quindecagon.</b>

2. A Polygon is **Regular** when all its sides are equal, and all its angles are equal.

3. A rectilinear figure is said to be **inscribed** in a circle, when all its angular points are on the circumference of the circle; and a circle is said to be **circumscribed about** a rectilinear figure, when the circumference of the circle passes through all the angular points of the figure.




4. A circle is said to be **inscribed in** a rectilinear figure, when the circumference of the circle is touched by each side of the figure; and a rectilinear figure is said to be **circumscribed about** a circle, when each side of the figure is a tangent to the circle.



5. A straight line is said to be **placed in** a circle, when its extremities are on the circumference of the circle.

PROPOSITION 1. PROBLEM.

*In a given circle to place a chord equal to a given straight line, which is not greater than the diameter of the circle.*



Let ABC be the given circle, and D the given straight line not greater than the diameter of the circle.

*It is required to place in the  $\odot$  ABC a chord equal to D.*

**Construction.** Draw CB, a diameter of the  $\odot$  ABC.

Then if  $CB = D$ , the thing required is done.

But if not, CB must be greater than D. *Hyp.*

From CB cut off CE equal to D: I. 3.

and with centre C, and radius CE, describe the  $\odot$  AEF, cutting the given circle at A.

Join CA.

*Then CA shall be the chord required.*

**Proof.** For  $CA = CE$ , being radii of the  $\odot$  AEF;

and  $CE = D$ :

$\therefore CA = D$ .

*Constr.*

Q.E.F.

EXERCISES.

1. In a given circle place a chord of given length so as to pass through a given point (i) without, (ii) within the circle.


When is this problem impossible?

2. In a given circle place a chord of given length so that it may be parallel to a given straight line.



## PROPOSITION 2. PROBLEM.

*In a given circle to inscribe a triangle equiangular to a given triangle.*



Let  $ABC$  be the given circle, and  $DEF$  the given triangle.  
*It is required to inscribe in the  $\odot ABC$  a triangle equiangular to the  $\triangle DEF$ .*

**Construction.** At any point  $A$ , on the  $\circ^{\infty}$  of the  $\odot ABC$ , draw the tangent  $GAH$ . III. 17.

At  $A$  make the  $\angle GAB$  equal to the  $\angle DFE$ ; I. 23.

and make the  $\angle HAC$  equal to the  $\angle DEF$ . I. 23.

Join  $BC$ .

*Then  $ABC$  shall be the triangle required.*

**Proof.** Because  $GH$  is a tangent to the  $\odot ABC$ ,  
 and from  $A$  its point of contact the chord  $AB$  is drawn,  
 $\therefore$  the  $\angle GAB =$  the  $\angle ACB$  in the alt. segment: III. 32.  
 but the  $\angle GAB =$  the  $\angle DFE$ ; *Constr.*  
 $\therefore$  the  $\angle ACB =$  the  $\angle DFE$ .

Similarly the  $\angle HAC =$  the  $\angle ABC$ , in the alt. segment:  
 $\therefore$  the  $\angle ABC =$  the  $\angle DEF$ . *Constr.*


Hence the third  $\angle BAC =$  the third  $\angle EDF$ ,  
 for the three angles in each triangle are together equal to  
 two rt. angles. I. 32.

$\therefore$  the  $\triangle ABC$  is equiangular to the  $\triangle DEF$ , and it is  
 inscribed in the  $\odot ABC$ .

Q. E. F.

PROPOSITION 3. PROBLEM.

About a given circle to circumscribe a triangle equiangular to a given triangle.



Let  $ABC$  be the given circle, and  $DEF$  the given triangle. It is required to circumscribe about the  $\odot ABC$  a triangle equiangular to the  $\triangle DEF$ .

**Construction.** Produce  $EF$  both ways to  $G$  and  $H$ .

Find  $K$  the centre of the  $\odot ABC$ , III. 1.  
and draw any radius  $KB$ .

At  $K$  make the  $\angle BKA$  equal to the  $\angle DEG$ ; I. 23.  
and make the  $\angle BKC$  equal to the  $\angle DFH$ .

Through  $A, B, C$  draw  $LM, MN, NL$  perp. to  $KA, KB, KC$ .

Then  $LMN$  shall be the triangle required.

**Proof.** Because  $LM, MN, NL$  are drawn perp. to radii at their extremities.

$\therefore LM, MN, NL$  are tangents to the circle. III. 16.

And because the four angles of the quadrilateral  $AKBM$  together = four rt. angles; I. 32. Cor.

and of these, the  $\angle^s KAM, KBM$  are rt. angles; *Constr.*

$\therefore$  the  $\angle^s AKB, AMB$  together = two rt. angles.

But the  $\angle^s DEG, DEF$  together = two rt. angles; I. 13.

$\therefore$  the  $\angle^s AKB, AMB =$  the  $\angle^s DEG, DEF$ ;

and of these, the  $\angle AKB =$  the  $\angle DEG$ ; *Constr.*

$\therefore$  the  $\angle AMB =$  the  $\angle DEF$ .


Similarly it may be shewn that the  $\angle LNM =$  the  $\angle DFE$ .

$\therefore$  the third  $\angle MLN =$  the third  $\angle EDF$ . I. 32.

$\therefore$  the  $\triangle LMN$  is equiangular to the  $\triangle DEF$ , and it is circumscribed about the  $\odot ABC$ . Q.E.F.

## PROPOSITION 4. PROBLEM.

To inscribe a circle in a given triangle



Let  $ABC$  be the given triangle.

It is required to inscribe a circle in the  $\triangle ABC$ .

**Construction.** Bisect the  $\angle^s$   $ABC$ ,  $ACB$  by the st. lines  $BI$ ,  $CI$ , which intersect at  $I$ . I. 9.

From  $I$  draw  $IE$ ,  $IF$ ,  $IG$  perp. to  $AB$ ,  $BC$ ,  $CA$ . I. 12.

**Proof.** Then in the  $\triangle^s$   $EIB$ ,  $FIB$ ,  
 Because  $\left\{ \begin{array}{l} \text{the } \angle EBI = \text{the } \angle FBI; \\ \text{and the } \angle BEI = \text{the } \angle BFI, \text{ being rt. angles;} \\ \text{and } BI \text{ is common;} \end{array} \right.$  *Constr.*  
 $\therefore IE = IF$ . I. 26.

Similarly it may be shewn that  $IF = IG$ .

$\therefore IE$ ,  $IF$ ,  $IG$  are all equal.

With centre  $I$ , and radius  $IE$ , describe a circle.

*This circle must pass through the points  $E$ ,  $F$ ,  $G$ ;  
 and it will be inscribed in the  $\triangle ABC$ .*

For since  $IE$ ,  $IF$ ,  $IG$ , being equal, are radii of the  $\odot EFG$ ;  
 and since the  $\angle^s$  at  $E$ ,  $F$ ,  $G$  are rt. angles; *Constr.*

$\therefore$  the  $\odot EFG$  is touched at these points by  $AB$ ,  $BC$ ,  $CA$ :  
 III. 16.

$\therefore$  the  $\odot EFG$  is inscribed in the  $\triangle ABC$ .

Q. E. F.

**NOTE.** From page 111 it is seen that if  $AI$  is joined, then  $AI$  bisects the angle  $BAC$ : hence it follows that

*The bisectors of the angles of a triangle are concurrent, the point of intersection being the centre of the inscribed circle.*

The centre of the circle inscribed in a triangle is usually called its **in-centre**.

### DEFINITION.

A circle which touches one side of a triangle and the other two sides produced is said to be an **escribed circle** of the triangle.

*To draw an escribed circle of a given triangle.*

Let  $ABC$  be the given triangle, of which the two sides  $AB, AC$  are produced to  $E$  and  $F$ .

*It is required to describe a circle touching  $BC$ , and  $AB, AC$  produced.*

Bisect the  $\angle^s CBE, BCF$  by the st. lines  $BI_1, CI_1$ , which intersect at  $I_1$ . I. 9.

From  $I_1$  draw  $I_1G, I_1H, I_1K$  perp. to  $AE, BC, AF$ . I. 12.

Then in the  $\triangle I_1BG, I_1BH$ ,  
 Because  $\left\{ \begin{array}{l} \text{the } \angle I_1BG = \text{the } \angle I_1BH, \text{ Constr.} \\ \text{and the } \angle I_1GB = \text{the } \angle I_1HB, \\ \text{being rt. angles;} \\ \text{also } I_1B \text{ is common;} \end{array} \right.$   
 $\therefore I_1G = I_1H$ .

Similarly it may be shewn that  $I_1H = I_1K$ ;

$\therefore I_1G, I_1H, I_1K$  are all equal.

With centre  $I_1$  and radius  $I_1G$ , describe a circle.

*This circle must pass through the points  $G, H, K$ ;  
 and it will be an escribed circle of the  $\triangle ABC$ .*


For since  $I_1H, I_1G, I_1K$ , being equal, are radii of the  $\odot HGK$ ,  
 and since the angles at  $H, G, K$  are rt. angles,  
 $\therefore$  the  $\odot HGK$  is touched at these points by  $BC$ , and by  $AB, AC$  produced:

$\therefore$  the  $\odot HGK$  is an escribed circle of the  $\triangle ABC$ . Q.E.F.

It is clear that every triangle has *three* escribed circles.


**NOTE.** From page 112 it is seen that if  $AI_1$  is joined, then  $AI_1$  bisects the angle  $BAC$ : hence it follows that

*The bisectors of two exterior angles of a triangle and the bisector of the third angle are concurrent, the point of intersection being the centre of an escribed circle.*



## PROPOSITION 5. PROBLEM.

To circumscribe a circle about a given triangle.



Let  $ABC$  be the given triangle.

It is required to circumscribe a circle about the  $\triangle ABC$ .

**Construction.** Draw  $DS$  bisecting  $AB$  at rt. angles; I. 11.  
and draw  $ES$  bisecting  $AC$  at rt. angles.

Then since  $AB, AC$  are neither par<sup>l</sup>, nor in the same st. line,  
 $\therefore DS$  and  $ES$  must meet at some point  $S$ .

Join  $SA$ ;

and if  $S$  be not in  $BC$ , join  $SB, SC$ .

**Proof.** Then in the  $\triangle ADS, BDS$ ,

$AD = BD$ ,

Because  $\left\{ \begin{array}{l} \text{and } DS \text{ is common to both;} \\ \text{and the } \angle ADS = \text{the } \angle BDS, \text{ being rt. angles;} \end{array} \right.$

$\therefore SA = SB$ .

I. 4.

Similarly it may be shewn that  $SC = SA$ .

$\therefore SA, SB, SC$  are all equal.

With centre  $S$ , and radius  $SA$ , describe a circle:  
this circle must pass through the points  $A, B, C$ , and is  
therefore circumscribed about the  $\triangle ABC$ . Q.E.F.

It follows that

(i) when the centre of the circumscribed circle falls *within* the triangle, each of its angles must be acute, for each angle is then in a segment greater than a semicircle:

(ii) when the centre falls *on one of the sides* of the triangle, the angle opposite to this side must be a right angle, for it is the angle in a semicircle:

(iii) when the centre falls *without* the triangle, the angle opposite to the side beyond which the centre falls, must be obtuse, for it is the angle in a segment less than a semicircle.

*Therefore, conversely, if the given triangle be acute-angled, the centre of the circumscribed circle falls within it: if it be a right-angled triangle, the centre falls on the hypotenuse: if it be an obtuse-angled triangle, the centre falls without the triangle.*

**NOTE.** From page 111 it is seen that if  $S$  is joined to the middle point of  $BC$ , then the joining line is perpendicular to  $BC$ .

*Hence the perpendiculars drawn to the sides of a triangle from their middle points are concurrent, the point of intersection being the centre of the circle circumscribed about the triangle.*

The centre of the circle circumscribed about a triangle is usually called its **circum-centre**.

#### EXERCISES.

##### ON THE INSCRIBED, CIRCUMSCRIBED, AND ESCRIBED CIRCLES OF A TRIANGLE.

1. An equilateral triangle is inscribed in a circle, and tangents are drawn at its vertices, prove that

- (i) the resulting figure is an equilateral triangle:
- (ii) its area is four times that of the given triangle.

2. Describe a circle to touch two parallel straight lines and a third straight line which meets them. Shew that two such circles can be drawn, and that they are equal.

3. *Triangles which have equal bases and equal vertical angles have equal circumscribed circles.*

4.  $I$  is the centre of the circle inscribed in the triangle  $ABC$ , and  $I_1$  is the centre of the circle which touches  $BC$  and  $AB, AC$  produced: shew that  $A, I, I_1$  are collinear.

5. *If the inscribed and circumscribed circles of a triangle are concentric, shew that the triangle is equilateral; and that the diameter of the circumscribed circle is double that of the inscribed circle.*

6.  $ABC$  is a triangle, and  $I, S$  are the centres of the inscribed and circumscribed circles; if  $A, I, S$  are collinear, shew that  $AB = AC$

7. The sum of the diameters of the inscribed and circumscribed circles of a right-angled triangle is equal to the sum of the sides containing the right angle.

8. If the circle inscribed in a triangle  $ABC$  touches the sides at  $D, E, F$ , shew that the triangle  $DEF$  is acute-angled; and express its angles in terms of the angles at  $A, B, C$ .

9. If  $I$  is the centre of the circle inscribed in the triangle  $ABC$ , and  $I_1$  the centre of the escribed circle which touches  $BC$ ; shew that  $I, B, I_1, C$  are concyclic.

10. In any triangle the difference of two sides is equal to the difference of the segments into which the third side is divided at the point of contact of the inscribed circle.

11. In the triangle  $ABC$  the bisector of the angle  $BAC$  meets the base at  $D$ , and from  $I$  the centre of the inscribed circle a perpendicular  $IE$  is drawn to  $BC$ : shew that the angle  $BID$  is equal to the angle  $CIE$ .

12. In the triangle  $ABC$ ,  $I$  and  $S$  are the centres of the inscribed and circumscribed circles: shew that  $IS$  subtends at  $A$  an angle equal to half the difference of the angles at the base of the triangle.

13. In a triangle  $ABC$ ,  $I$  and  $S$  are the centres of the inscribed and circumscribed circles, and  $AD$  is drawn perpendicular to  $BC$ : shew that  $AI$  is the bisector of the angle  $DAS$ .

14. Shew that the area of a triangle is equal to the rectangle contained by its semi-perimeter and the radius of the inscribed circle.

15. The diagonals of a quadrilateral  $ABCD$  intersect at  $O$ : shew that the centres of the circles circumscribed about the four triangles  $AOB, BOC, COD, DOA$  are at the angular points of a parallelogram.

16. In any triangle  $ABC$ , if  $I$  is the centre of the inscribed circle, and if  $AI$  is produced to meet the circumscribed circle at  $O$ ; shew that  $O$  is the centre of the circle circumscribed about the triangle  $BIC$ .

17. Given the base, altitude, and the radius of the circumscribed circle; construct the triangle.


18. Describe a circle to intercept equal chords of given length on three given straight lines.

19. In an equilateral triangle the radii of the circumscribed and escribed circles are respectively double and treble of the radius of the inscribed circle.

20. Three circles whose centres are  $A, B, C$  touch one another externally two by two at  $D, E, F$ : shew that the inscribed circle of the triangle  $ABC$  is the circumscribed circle of the triangle  $DEF$ .

PROPOSITION 6. PROBLEM.

To inscribe a square in a given circle.



Let ABCD be the given circle.

It is required to inscribe a square in the  $\odot$  ABCD.

**Construction.** Find E the centre of the circle : III. 1.  
and draw two diameters AC, BD perp. to one another. I. 11.  
Join AB, BC, CD, DA.

Then the fig. ABCD shall be the square required.

**Proof.** For in the  $\triangle$  BEA, DEA,  
Because  $\left\{ \begin{array}{l} BE = DE, \quad \text{I. Def. 15.} \\ \text{and EA is common;} \\ \text{and the } \angle BEA = \text{the } \angle DEA, \text{ being rt. angles;} \\ \therefore BA = DA. \quad \text{I. 4.} \end{array} \right.$

Similarly it may be shewn that  $CD = DA$ , and that  $BC = CD$ .  
 $\therefore$  the fig. ABCD is equilateral.

And since BD is a diameter of the  $\odot$  ABCD,  
 $\therefore$  BAD is a semicircle;  
 $\therefore$  the  $\angle$  BAD is a rt. angle. III. 31.

Similarly the other angles of the fig. ABCD are rt. angles.  
 $\therefore$  the fig. ABCD is a square;  
and it is inscribed in the given circle.


Q.E.F.

[For Exercises see page 281.]



## PROPOSITION 7. PROBLEM.

To circumscribe a square about a given circle.



Let  $\odot ABCD$  be the given circle.

It is required to circumscribe a square about the  $\odot ABCD$ .

**Construction.** Find  $E$  the centre of the  $\odot ABCD$ : III. 1. and draw two diameters  $AC, BD$  perp. to one another. I. 11. Through  $A, B, C, D$  draw  $FG, GH, HK, KF$  perp. to  $EA, EB, EC, ED$ .

Then the fig.  $GK$  shall be the square required.

**Proof.** Because  $FG, GH, HK, KF$  are drawn perp. to radii at their extremities,

$\therefore FG, GH, HK, KF$  are tangents to the circle. III. 16.

And because the  $\angle^s AEB, EBG$  are both rt. angles, *Constr.*

$\therefore GH$  is par<sup>l</sup> to  $AC$ . I. 28.

Similarly  $FK$  is par<sup>l</sup> to  $AC$ :

and in like manner  $GF, BD, HK$  are par<sup>l</sup>.

Hence the figs.  $GK, GC, AK, GD, BK, GE$  are par<sup>ms</sup>.

$\therefore GF$  and  $HK$  each =  $BD$ ;

also  $GH$  and  $FK$  each =  $AC$ ;

but  $AC = BD$ ;

$\therefore GF, FK, KH, HG$  are all equal:

that is, the fig.  $GK$  is equilateral.

And since the fig.  $GE$  is a par<sup>m</sup>,

$\therefore$  the  $\angle BGA =$  the  $\angle BEA$ ;

I. 34.

but the  $\angle BEA$  is a rt. angle;

*Constr.*


$\therefore$  the  $\angle$  at  $G$  is a rt. angle.

Similarly the  $\angle^s$  at  $F, K, H$  are rt. angles.

$\therefore$  the fig.  $GK$  is a square, and it has been circumscribed about the  $\odot ABCD$  Q.E.F.

PROPOSITION 8. PROBLEM.

To inscribe a circle in a given square.



Let ABCD be the given square.

It is required to inscribe a circle in the square ABCD.

**Construction.** Bisect the sides AB, AD at F and E. I. 10.  
 Through E draw EH par<sup>l</sup> to AB or DC: I. 31.  
 and through F draw FK par<sup>l</sup> to AD or BC, meeting EH at G.

**Proof.** Now AB = AD, being the sides of a square;  
 and their halves are equal Ax. 7.  
 $\therefore AF = AE.$   
 But the fig. AG is a par<sup>m</sup>; Constr.  
 $\therefore AF = GE,$  and  $AE = GF;$   
 $\therefore GE = GF.$

Similarly it may be shewn that  $GE = GK,$  and  $GK = GH:$   
 $\therefore GF, GE, GK, GH$  are all equal.


With centre G, and radius GE, describe a circle.  
 This circle must pass through the points F, E, K, H;  
 and it will be touched by BA, AD, DC, CB; III. 16.  
 for GF, GE, GK, GH, being equal, are radii;  
 and the angles at F, E, K, H are rt. angles. I. 29.  
 Hence the  $\odot FEKH$  is inscribed in the sq. ABCD.

Q.E.F.

[For Exercises see p. 281.]

## PROPOSITION 9. PROBLEM.

*To circumscribe a circle about a given square.*



Let ABCD be the given square.

*It is required to circumscribe a circle about the square ABCD.*

**Construction.** Join AC, BD, intersecting at E.

**Proof.**

Then in the  $\triangle^s$  BAC, DAC,

Because  $\left\{ \begin{array}{l} BA = DA, \\ \text{and AC is common;} \\ \text{and BC = DC;} \end{array} \right.$  I. Def. 30.

$\therefore$  the  $\angle$  BAC = the  $\angle$  DAC; I. 8.

that is, the diagonal AC bisects the  $\angle$  BAD.

Similarly the remaining angles of the square are bisected by the diagonals AC or BD.

Hence each of the  $\angle^s$  EAD, EDA is half a rt. angle;

$\therefore$  the  $\angle$  EAD = the  $\angle$  EDA:

$\therefore$  EA = ED. I. 6.

Similarly it may be shewn that ED = EC, and EC = EB.

$\therefore$  EA, EB, EC, ED are all equal.

With centre E, and radius EA, describe a circle:  
this circle must pass through the points A, B, C, D, and is  
therefore circumscribed about the sq. ABCD. Q.E.F.

**DEFINITION.** A rectilineal figure about which a circle may be described is said to be **Cyclic**.

## EXERCISES ON PROPOSITIONS 6-9.

1. *If a circle can be inscribed in a quadrilateral, shew that the sum of one pair of opposite sides is equal to the sum of the other pair.*

2. *If the sum of one pair of opposite sides of a quadrilateral is equal to the sum of the other pair, shew that a circle may be inscribed in the figure.*

[Bisect two adjacent angles of the figure, and so describe a circle to touch three of its sides. Then prove indirectly by means of the last exercise that this circle must also touch the fourth side.]

3. *Prove that a rhombus and a square are the only parallelograms in which a circle can be inscribed.*

4. *All cyclic parallelograms are rectangular.*

5. *The greatest rectangle which can be inscribed in a given circle is a square.*

6. Circumscribe a rhombus about a given circle.

7. All squares circumscribed about a given circle are equal.

8. The area of a square circumscribed about a circle is double of the area of the inscribed square.

9.  $ABCD$  is a square inscribed in a circle, and  $P$  is any point on the arc  $AD$ : shew that the side  $AD$  subtends at  $P$  an angle three times as great as that subtended at  $P$  by any one of the other sides.

10. Inscribe a square in a given square  $ABCD$ , so that one of its angular points shall be at a given point  $X$  in  $AB$ .

11. In a given square inscribe the square of minimum area.


12. Describe (i) a circle, (ii) a square about a given rectangle.

13. Inscribe (i) a circle, (ii) a square in a given quadrant.

14.  $ABCD$  is a square inscribed in a circle, and  $P$  is any point on the circumference; shew that the sum of the squares on  $PA$ ,  $PB$ ,  $PC$ ,  $PD$  is double the square on the diameter. [See Ex. 24, p. 161.]

## PROPOSITION 10. PROBLEM.

To describe an isosceles triangle having each of the angles at the base double of the third angle.



**Construction.** Take any straight line  $AB$ .  
Divide  $AB$  at  $C$ , so that the rect.  $BA, BC =$  the sq. on  $AC$ . II. 11.

With centre  $A$ , and radius  $AB$ , describe the  $\odot BDE$ ;  
and in it place the chord  $BD$  equal to  $AC$ . IV. 1.  
Join  $DA$ .

*Then  $ABD$  shall be the triangle required.*

Join  $CD$ ;  
and about the  $\triangle ACD$  circumscribe a circle. IV. 5.

**Proof.** Now the rect.  $BA, BC =$  the sq. on  $AC$  *Constr.*  
 $=$  the sq. on  $BD$ . *Constr.*

Hence  $BD$  is a tangent to the  $\odot ACD$ : III. 37.  
and from the point of contact  $D$  a chord  $DC$  is drawn;  
 $\therefore$  the  $\angle BDC =$  the  $\angle CAD$  in the alt. segment. III. 32.

To each of these equals add the  $\angle CDA$ :  
then the whole  $\angle BDA =$  the sum of the  $\angle^s CAD, CDA$ .

But the ext.  $\angle DCB =$  the sum of the  $\angle^s CAD, CDA$ ; I. 32.  
 $\therefore$  the  $\angle DCB =$  the  $\angle BDA$ .

And since  $AB = AD$ , being radii of the  $\odot BDE$ ,  
 $\therefore$  the  $\angle DBA =$  the  $\angle BDA$ ; I. 5.  
 $\therefore$  the  $\angle DBC =$  the  $\angle DCB$ ;

$\therefore DC = DB;$  I. 6.  
 that is,  $DC = CA;$  *Constr.*  
 $\therefore$  the  $\angle CAD =$  the  $\angle CDA;$  I. 5.  
 $\therefore$  the sum of the  $\angle^s CAD, CDA =$  twice the angle at A.  
**But** the  $\angle ADB =$  the sum of the  $\angle^s CAD, CDA;$  *Proved.*  
 $\therefore$  each of the  $\angle^s ABD, ADB =$  twice the angle at A.  
 Q.E.F.

## EXERCISES ON PROPOSITION 10.

1. In an isosceles triangle in which each of the angles at the base is double of the vertical angle, shew that the vertical angle is one-fifth of two right angles.

2. Divide a right angle into five equal parts.

3. Describe an isosceles triangle whose vertical angle shall be three times either angle at the base. Point out a triangle of this kind in the figure of Proposition 10.

4. In the figure of Proposition 10, if the two circles intersect at F, shew that  $BD = DF$ .

5. In the figure of Proposition 10, shew that the circle ACD is equal to the circle circumscribed about the triangle ABD.

6. In the figure of Proposition 10, if the two circles intersect at F, shew that

(i) BD, DF are sides of a regular decagon inscribed in the circle EBD.


(ii) AC, CD, DF are sides of a regular pentagon inscribed in the circle ACD.

7. In the figure of Proposition 10, shew that the centre of the circle circumscribed about the triangle DBC is the middle point of the arc CD.

8. In the figure of Proposition 10, if I is the centre of the circle inscribed in the triangle ABD, and I', S' the centres of the inscribed and circumscribed circles of the triangle DBC, shew that  $S'I = S'I'$ .

## PROPOSITION 11. PROBLEM.

To inscribe a regular pentagon in a given circle.



Let  $\odot ABC$  be a given circle.

It is required to inscribe a regular pentagon in the  $\odot ABC$ .

**Construction.** Describe an isosceles  $\triangle FGH$ , having each of the angles at  $G$  and  $H$  double of the angle at  $F$ . IV. 10.

In the  $\odot ABC$  inscribe the  $\triangle ACD$  equiangular to the  $\triangle FGH$ , IV. 2.

so that each of the  $\angle^s ACD, ADC$  is double of the  $\angle CAD$ .

Bisect the  $\angle^s ACD, ADC$  by  $CE$  and  $DB$ , which meet the  $\odot^{\circ}$  at  $E$  and  $B$ . I. 9.

Join  $AB, BC, AE, ED$ .

Then  $ABCDE$  shall be the required regular pentagon.

**Proof.** Since each of the  $\angle^s ACD, ADC =$  twice the  $\angle CAD$ ;

and since the  $\angle^s ACD, ADC$  are bisected by  $CE, DB$ ,

$\therefore$  the five  $\angle^s ADB, BDC, CAD, DCE, ECA$  are all equal.

$\therefore$  the five arcs  $AB, BC, CD, DE, EA$  are all equal. III. 26.

$\therefore$  the five chords  $AB, BC, CD, DE, EA$  are all equal. III. 29.

$\therefore$  the pentagon  $ABCDE$  is equilateral.

Again the arc  $AB =$  the arc  $DE$ ; *Proved.*

to each of these equals add the arc  $BCD$ ;

$\therefore$  the arc  $ABCD =$  the arc  $BCDE$ ;

hence the angles at the  $\odot^{\circ}$  which stand upon these equal arcs are equal; III. 27.

that is, the  $\angle AED =$  the  $\angle BAE$ .


In like manner the remaining angles of the pentagon may be shewn to be equal;

$\therefore$  the pentagon  $ABCDE$  is equiangular.

Hence the pentagon, being both equilateral and equiangular, is regular; and it is inscribed in the  $\odot ABC$ . Q.E.F.

PROPOSITION 12. PROBLEM.

To circumscribe a regular pentagon about a given circle.



Let ABCD be the given circle.

It is required to circumscribe a regular pentagon about the  $\odot$  ABCD.

**Construction.**

Inscribe a regular pentagon in the  $\odot$  ABCD, IV. 11.  
and let A, B, C, D, E be its angular points.

At the points A, B, C, D, E draw GH, HK, KL, LM, MG,  
tangents to the circle. III. 17.

Then shall GHKLM be the required regular pentagon.

Find F the centre of the  $\odot$  ABCD ; III. 1.  
and join FB, FK, FC, FL, FD.

**Proof.**

In the  $\triangle$  BFK, CFK,

Because  $\left\{ \begin{array}{l} \text{BF} = \text{CF, being radii of the circle,} \\ \text{and FK is common;} \\ \text{and KB} = \text{KC, being tangents to the circle from} \\ \text{the same point K;} \end{array} \right.$  III. 17, Cor.

$\therefore$  the  $\angle$  BFK = the  $\angle$  CFK, I. 8.

also the  $\angle$  BKF = the  $\angle$  CKF. I. 8, Cor.

Hence the  $\angle$  BFC = twice the  $\angle$  CFK,

and the  $\angle$  BKC = twice the  $\angle$  CKF.

Similarly it may be shewn

that the  $\angle$  CFD = twice the  $\angle$  CFL,

and that the  $\angle$  CLD = twice the  $\angle$  CLF.


But since the arc BC = the arc CD, IV. 11.

$\therefore$  the  $\angle$  BFC = the  $\angle$  CFD ; III. 27.

and the halves of these angles are equal,

that is, the  $\angle$  CFK = the  $\angle$  CFL.





Then in the  $\triangle$ 's CFK, CFL,  
 Because { the  $\angle$  CFK = the  $\angle$  CFL, *Proved—*  
 and the  $\angle$  FCK = the  $\angle$  FCL, being rt. angles, III. 18—  
 and FC is common ;  
 $\therefore$  CK = CL, I. 26—  
 and the  $\angle$  FKC = the  $\angle$  FLC.

Hence KL is double of KC; similarly HK is double of KB.

And since KC = KB, III. 17, *Cor.*

$\therefore$  KL = HK.

In the same way it may be shewn that every two consecutive sides are equal ;

$\therefore$  the pentagon GHKLM is equilateral.

Again, it has been proved that the  $\angle$  FKC = the  $\angle$  FLC,  
 and that the  $\angle$ 's HKL, KLM are respectively double of these angles :

$\therefore$  the  $\angle$  HKL = the  $\angle$  KLM.

In the same way it may be shewn that every two consecutive angles of the figure are equal ;

$\therefore$  the pentagon GHKLM is equiangular.


$\therefore$  the pentagon is regular, and it is circumscribed about the  $\odot$  ABCD. Q.E.F.

**COROLLARY.** *Similarly it may be proved that if tangents are drawn at the vertices of any regular polygon inscribed in a circle, they will form another regular polygon of the same species circumscribed about the circle.*

[For Exercises see p. 293.]

PROPOSITION 13. PROBLEM.

To inscribe a circle in a given regular pentagon.



Let  $ABCDE$  be the given regular pentagon.

It is required to inscribe a circle within the figure  $ABCDE$ .

**Construction.** Bisect two consecutive  $\angle$ 's  $BCD$ ,  $CDE$  by  $CF$  and  $DF$  which intersect at  $F$ . I. 9.

Join  $FB$ ;  
and draw  $FH$ ,  $FK$  perp. to  $BC$ ,  $CD$ . I. 12.


**Proof.** In the  $\triangle$ 's  $BCF$ ,  $DCF$ ,  
Because  $\left\{ \begin{array}{l} BC = DC, \\ \text{and } CF \text{ is common to both;} \\ \text{and the } \angle BCF = \text{the } \angle DCF; \end{array} \right.$  Hyp.  
 $\therefore$  the  $\angle CBF = \text{the } \angle CDF$ . Constr. I. 4.

But the  $\angle CDF$  is half an angle of the regular pentagon:  
 $\therefore$  also the  $\angle CBF$  is half an angle of the regular pentagon:  
that is,  $FB$  bisects the  $\angle ABC$ .

So it may be shewn that if  $FA$ ,  $FE$  were joined, these lines would bisect the  $\angle$ 's at  $A$  and  $E$ .

Again, in the  $\triangle$ 's  $FCH$ ,  $FCK$ ,  
Because  $\left\{ \begin{array}{l} \text{the } \angle FCH = \text{the } \angle FCK, \\ \text{and the } \angle FHC = \text{the } \angle FKC, \text{ being rt. angles;} \\ \text{also } FC \text{ is common;} \end{array} \right.$  Constr.  
 $\therefore FH = FK$ . I. 26.

Similarly if  $FG$ ,  $FM$ ,  $FL$  be drawn perp. to  $BA$ ,  $AE$ ,  $ED$ , it may be shewn that the five perpendiculars drawn from  $F$  to the sides of the pentagon are all equal.



With centre  $F$ , and radius  $FH$ , describe a circle ;  
 this circle must pass through the points  $H, K, L, M, G$  ;  
 and it will be touched at these points by the sides of the  
 pentagon, for the  $\angle$ 's at  $H, K, L, M, G$  are  $\text{rt. } \angle$ 's. *Constr.*  
 $\therefore$  the  $\odot$   $HKLMG$  is inscribed in the given pentagon. *Q.E.F.*

*COROLLARY.* *The bisectors of the angles of a regular pentagon meet at a point.*

*NOTE.* In the same way it may be shewn that the bisectors of the angles of any regular polygon meet at a point. [See Ex. 1, p. 294.]

[For Exercises on Regular Polygons see p. 293.]

#### MISCELLANEOUS EXERCISES.

1. Two tangents  $AB, AC$  are drawn from an external point  $A$  to a given circle : describe a circle to touch  $AB, AC$  and the convex arc intercepted by them on the given circle.


2.  $ABC$  is an isosceles triangle, and from the vertex  $A$  a straight line is drawn to meet the base at  $D$  and the circumference of the circumscribed circle at  $E$  : shew that  $AB$  is a tangent to the circle circumscribed about the triangle  $BDE$ .

3. An equilateral triangle is inscribed in a given circle : shew that twice the square on one of its sides is equal to three times the area of the square inscribed in the same circle.

4.  $ABC$  is an isosceles triangle in which each of the angles at  $B$  and  $C$  is double of the angle at  $A$  : shew that the square on  $AB$  is equal to the rectangle  $AB, BC$  with the square on  $BC$ .

PROPOSITION 14. PROBLEM.

To circumscribe a circle about a given regular pentagon.



Let ABCDE be the given regular pentagon.

It is required to circumscribe a circle about the figure ABCDE.

**Construction.** Bisect the  $\angle$  BCD, CDE by CF, DF, intersecting at F. I. 9.

Join FB, FA, FE.

**Proof.** In the  $\triangle$  BCF, DCF,  
 Because  $\left\{ \begin{array}{l} BC = DC, \\ \text{and CF is common to both;} \\ \text{and the } \angle BCF = \text{the } \angle DCF; \end{array} \right.$  Hyp.  
 $\therefore$  the  $\angle CBF = \text{the } \angle CDF.$  Constr. I. 4.

But the  $\angle CDF$  is half an angle of the regular pentagon :  
 $\therefore$  also the  $\angle CBF$  is half an angle of the regular pentagon :  
 that is, FB bisects the  $\angle ABC$ .

So it may be shewn that FA, FE bisect the  $\angle$  at A and E.

Now the  $\angle$  FCD, FDC are each half an angle of the given regular pentagon ;

$\therefore$  the  $\angle FCD = \text{the } \angle FDC,$  IV. Def. 2.  
 $\therefore FC = FD.$  I. 6.


Similarly it may be shewn that FA, FB, FC, FD, FE are all equal.

With centre F, and radius FA, describe a circle :  
 this circle must pass through the points A, B, C, D, E,  
 and therefore is circumscribed about the pentagon. Q.E.F.

**NOTE.** In the same way a circle may be circumscribed about any regular polygon.

## PROPOSITION 15. PROBLEM.

To inscribe a regular hexagon in a given circle.



Let  $ABDF$  be the given circle.

It is required to inscribe a regular hexagon in the  $\odot ABDF$ .

**Construction.** Find  $G$  the centre of the  $\odot ABDF$ ; III. 1. and draw a diameter  $AGD$ .

With centre  $D$ , and radius  $DG$ , describe the  $\odot EGCH$ .

Join  $CG$ ,  $EG$ , and produce them to cut the  $\odot$  of the given circle at  $F$  and  $B$ .

Join  $AB$ ,  $BC$ ,  $CD$ ,  $DE$ ,  $EF$ ,  $FA$ .

Then  $ABCDEF$  shall be the required regular hexagon.

**Proof.** Now  $GE = GD$ , being radii of the  $\odot ACE$  ;  
and  $DG = DE$ , being radii of the  $\odot EHC$  :

$\therefore GE$ ,  $ED$ ,  $DG$  are all equal, and the  $\triangle EGD$  is equilateral.

Hence the  $\angle EGD =$  one-third of two rt. angles. I. 32.

Similarly the  $\angle DGC =$  one-third of two rt. angles.

But the  $\angle EGD$ ,  $DGC$ ,  $CGB$  together = two rt. angles ; I. 13.

$\therefore$  the remaining  $\angle CGB =$  one-third of two rt. angles.

$\therefore$  the three  $\angle EGD$ ,  $DGC$ ,  $CGB$  are equal to one another.

And to these angles the vert. opp.  $\angle BGA$ ,  $AGF$ ,  $FGE$  are respectively equal :

$\therefore$  the  $\angle EGD$ ,  $DGC$ ,  $CGB$ ,  $BGA$ ,  $AGF$ ,  $FGE$  are all equal ;

$\therefore$  the arcs  $ED$ ,  $DC$ ,  $CB$ ,  $BA$ ,  $AF$ ,  $FE$  are all equal : III. 26.

$\therefore$  the chords  $ED$ ,  $DC$ ,  $CB$ ,  $BA$ ,  $AF$ ,  $FE$  are all equal : III. 29.

$\therefore$  the hexagon is equilateral.

Again the arc  $FA =$  the arc  $DE$  : *Proved.*

to each of these equals add the arc  $ABCD$  ;

then the arc  $FABCD =$  the arc  $ABCDE$  :

hence the angles at the  $\odot$  which stand on these equal arcs are equal.

that is, the  $\angle FED =$  the  $\angle AFE$ . III, 27.

In like manner the remaining angles of the hexagon may be shewn to be equal.

$\therefore$  the hexagon is equiangular;

$\therefore$  the hexagon ABCDEF is regular, and it is inscribed in the

$\odot$  ABDF.

Q.E.F.

**COROLLARY.** *The side of a regular hexagon inscribed in a circle is equal to the radius of the circle.*

#### SUMMARY OF THE PROPOSITIONS OF BOOK IV.

The following summary will assist the student in remembering the sequence of the Propositions of Book IV.

(i) Of the sixteen Propositions of this Book, Props. 1, 10, 15, 16 deal with isolated constructions.

(ii) The remaining twelve Propositions may be divided into *three* groups of *four* each, as follows:

(a) Group 1. Props. 2, 3, 4, 5 deal with *triangles* and *circles*.

(b) Group 2. Props. 6, 7, 8, 9 deal with *squares* and *circles*.


(c) Group 3. Props. 11, 12, 13, 14 deal with *pentagons* and *circles*.

(iii) In each group the problem of *inscription* precedes the corresponding problem of *circumscription*.

Further, each group deals with the inscription and circumscription of *rectilineal* figures first and of *circles* afterwards.

## PROPOSITION 16. PROBLEM.

To inscribe a regular quindecagon in a given circle.



Let ABCD be the given circle.

It is required to inscribe a regular quindecagon in the  $\odot$  ABCD.

**Construction.**

In the  $\odot$  ABCD inscribe an equilateral triangle, IV. 2.  
and let AC be one of its sides.

In the same circle inscribe a regular pentagon, IV. 11.  
and let AB be one of its sides.

**Proof.**

Now of such equal parts as the whole  $\odot^{\infty}$  contains fifteen,  
the arc AC, which is one-third of the  $\odot^{\infty}$ , contains five,  
and the arc AB, which is one-fifth of the  $\odot^{\infty}$ , contains three;  
 $\therefore$  their difference, the arc BC, contains two.

Bisect the arc BC at E: III. 30.  
then each of the arcs BE, EC is one-fifteenth of the  $\odot^{\infty}$ .

$\therefore$  if BE, EC be joined, and st. lines equal to them be  
placed successively round the circle, a regular quindecagon  
will be inscribed in it. Q.E.F.

EXERCISES ON PROPOSITIONS 11—16.

1. Express in terms of a right angle the magnitude of an angle of the following *regular* polygons :

- (i) a pentagon, (ii) a hexagon, (iii) an octagon,  
(iv) a decagon, (v) a quindecagon.

2. Any angle of a regular pentagon is trisected by the straight lines which join it to the opposite vertices.

3. In a polygon of  $n$  sides the straight lines which join any angular point to the vertices not adjacent to it, divide the angle into  $n - 2$  equal parts.

4. Shew how to construct on a given straight line  
(i) a regular pentagon, (ii) a regular hexagon, (iii) a regular octagon.

5. An equilateral triangle and a regular hexagon are inscribed in a given circle ; shew that

- (i) the area of the triangle is half that of the hexagon ;  
(ii) the square on the side of the triangle is three times the square on the side of the hexagon.

6. ABCDE is a regular pentagon, and AC, BE intersect at H : shew that

- (i)  $AB = CH = EH$ .  
(ii) AB is a tangent to the circle circumscribed about the triangle BHC.  
(iii) AC and BE cut one another in medial section.

7. The straight lines which join alternate vertices of a regular pentagon intersect so as to form another regular pentagon.

8. The straight lines which join alternate vertices of a regular polygon of  $n$  sides, intersect so as to form another regular polygon of  $n$  sides.

If  $n = 6$ , shew that the area of the resulting hexagon is one-third of the given hexagon.

9. By means of iv. 16, inscribe in a circle a triangle whose angles are as the numbers 2, 5, 8.

10. Shew that the area of a regular hexagon inscribed in a circle is three-fourths of that of the corresponding circumscribed hexagon.




## NOTE ON REGULAR POLYGONS.

The following propositions, proved by Euclid for a regular pentagon, hold good for all regular polygons.

1. *The bisectors of the angles of any regular polygon are concurrent.*

Let D, E, A, B, C be consecutive angular points of a regular polygon of any number of sides.

Bisect the  $\angle^s$  EAB, ABC by AO, BO, which intersect at O.



Join EO.

*It is required to prove that EO bisects the  $\angle$  DEA.*

For in the  $\triangle^s$  EAO, BAO,  
 Because  $\left\{ \begin{array}{l} EA = BA, \text{ being sides of a regular polygon;} \\ \text{and } AO \text{ is common;} \\ \text{and the } \angle \text{ EAO} = \text{the } \angle \text{ BAO;} \end{array} \right. \quad \begin{array}{l} \text{Constr.} \\ \text{I. 4.} \end{array}$   
 $\therefore$  the  $\angle$  OEA = the  $\angle$  OBA.

But the  $\angle$  OBA is half the  $\angle$  ABC; Constr.  
 also the  $\angle$  ABC = the  $\angle$  DEA, since the polygon is regular;  
 $\therefore$  the  $\angle$  OEA is half the  $\angle$  DEA:  
 that is, EO bisects the  $\angle$  DEA.

Similarly if O be joined to the remaining angular points of the polygon, it may be proved that each joining line bisects the angle to whose vertex it is drawn.

That is to say, the bisectors of the angles of the polygon meet at the point O. Q. E. D.

COROLLARIES. Since the  $\angle$  EAB = the  $\angle$  ABC; Hyp.  
 and since the  $\angle^s$  OAB, OBA are respectively half of the  $\angle^s$  EAB, ABC;  
 $\therefore$  the  $\angle$  OAB = the  $\angle$  OBA;

$\therefore$  OA = OB. I. 6.  
 OE = OA.

Similarly

*Hence the bisectors of the angles of a regular polygon are all equal.  
 Therefore a circle described with centre O, and radius OA, will be circumscribed about the polygon.*

Also it may be shewn, as in Proposition 13, that perpendiculars drawn from O to the sides of the polygon are all equal.

*Therefore a circle described with centre O, and any one of these perpendiculars as radius, will be inscribed in the polygon.*


2. *If a polygon inscribed in a circle is equilateral, it is also equiangular.*

Let AB, BC, CD be consecutive sides of an equilateral polygon inscribed in the  $\odot$  ADK.

Then shall this polygon be equiangular.

Because the chord AB = the chord DC. Hyp.  
 $\therefore$  the minor arc AB = the minor arc DC. III. 28.

To each of these equals add the arc AKD :  
 then the arc BAKD = the arc AKDC ;  
 $\therefore$  the angles at the  $\odot^{\infty}$ , which stand on these equal arcs, are equal ;  
 that is, the  $\angle BCD =$  the  $\angle ABC$ . III. 27.




Similarly the remaining angles of the polygon may be shewn to be equal :

the polygon is equiangular. Q.E.D.

3. *If a polygon inscribed in a circle is equiangular, it is also equilateral, provided that the number of its sides is odd.*

[Observe that Theorems 2 and 3 are only true of polygons inscribed in a circle.



The above figures are sufficient to shew that otherwise a polygon may be equilateral without being equiangular, Fig. 1 ; or equiangular without being equilateral, Fig. 2.]

NOTE. The following extensions of Euclid's constructions for Regular Polygons should be noticed.

By continual bisection of arcs, we are enabled to divide the circumference of a circle,

- by means of Proposition 6, into 4, 8, 16, ...,  $2 \cdot 2^n$ , ... equal parts ;
- by means of Proposition 15, into 3, 6, 12, ...,  $3 \cdot 2^n$ , ... equal parts ;
- by means of Proposition 11, into 5, 10, 20, ...,  $5 \cdot 2^n$ , ... equal parts ;
- by means of Proposition 16, into 15, 30, 60, ...,  $15 \cdot 2^n$ , ... equal parts.

Hence we can inscribe in a circle a regular polygon the number of whose sides is included in any one of the formulæ  $2 \cdot 2^n$ ,  $3 \cdot 2^n$ ,  $5 \cdot 2^n$ ,  $15 \cdot 2^n$ ,  $n$  being any positive integer. It has also been shewn (by Gauss, 1800) that a regular polygon of  $2^n + 1$  sides may be inscribed in a circle, provided  $2^n + 1$  is a prime number.

## QUESTIONS FOR REVISION ON BOOK IV.

1. With what difference of meaning is the word *inscribed* used in the following cases?

- (i) a triangle *inscribed* in a circle ;
- (ii) a circle *inscribed* in a triangle.

2. What is meant by a *cyclic figure*? Shew that *all triangles* are cyclic.

What is the condition that a quadrilateral may be *cyclic*?

Shew that cyclic parallelograms must be rectangular.

3. Shew that the only *regular* figures which may be fitted together so as to form a plane surface are (i) *equilateral triangles*, (ii) *squares*, (iii) *regular hexagons*.

4. Employ the first Corollary of I. 32 to shew that in any regular polygon of  $n$  sides each interior angle contains  $\frac{2(n-2)}{n}$  right angles?

5. *The bisectors of the angles of a regular polygon are concurrent.* State the *method of proof* employed in this and similar theorems.

6. Shew that

- (i) all squares inscribed in a given circle are equal ; and
- (ii) all equilateral triangles circumscribed about a given circle are equal.

7. How many circles can be described to touch each of three given straight lines of unlimited length?

- (i) when no two of the lines are parallel ;
- (ii) when two only are parallel ;
- (iii) when all three are parallel.

8. Prove that the greatest triangle which can be inscribed in a circle on a diameter as base, is one-fourth of the circumscribed square.

9. The radius of a given circle is 10 inches : find the length of a side of

- (i) the circumscribed square ; [20 inches.]
- (ii) the inscribed square ;  $\sqrt{2}$  inches.]
- (iii) the inscribed equilateral triangle ; [10 $\sqrt{3}$  inches.]
- (iv) the circumscribed equilateral triangle ; [20 $\sqrt{3}$  inches.]
- (v) the inscribed regular hexagon. [10 inches.]

Shew also that the areas of these figures are respectively 400, 200, 75 $\sqrt{3}$ , 300 $\sqrt{3}$ , and 150 $\sqrt{3}$  square inches.


## THEOREMS AND EXAMPLES ON BOOK IV.

### I. ON THE TRIANGLE AND ITS CIRCLES.


1.  $D, E, F$  are the points of contact of the inscribed circle of the triangle  $ABC$ , and  $D_1, E_1, F_1$  the points of contact of the escribed circle, which touches  $BC$  and the other sides produced:  $a, b, c$  denote the length of the sides  $BC, CA, AB$ ;  $s$  the semi-perimeter of the triangle, and  $r, r_1$  the radii of the inscribed and escribed circles.

*Prove the following equalities:*

- (i)  $AE = AF = s - a,$   
 $BD = BF = s - b,$   
 $CD = CE = s - c,$
- (ii)  $AE_1 = AF_1 = s.$
- (iii)  $CD_1 = CE_1 = s - b,$   
 $BD_1 = BF_1 = s - c.$
- (iv)  $CD = BD_1$  and  $BD = CD_1.$
- (v)  $EE_1 = FF_1 = a.$
- (vi) The area of the  $\triangle ABC$   
 $= rs = r_1(s - a).$



2. In the triangle  $ABC$ ,  $I$  is the centre of the inscribed circle, and  $I_1, I_2, I_3$  the centres of the escribed circles touching respectively the sides  $BC, CA, AB$  and the other sides produced.



Prove the following properties :

- (i) The points  $A, I, I_1$  are collinear: so are  $B, I, I_2$ ; and  $C, I, I_3$ .
- (ii) The points  $I_2, A, I_3$  are collinear; so are  $I_3, B, I_1$ ; and  $I_1, C, I_2$ .
- (iii) The triangles  $BI_1C, CI_2A, AI_3B$  are equiangular to one another.
- (iv) The triangle  $I_1I_2I_3$  is equiangular to the triangle formed by joining the points of contact of the inscribed circle.
- (v) Of the four points  $I, I_1, I_2, I_3$  each is the orthocentre of the triangle whose vertices are the other three.
- (vi) The four circles, each of which passes through three of the points  $I, I_1, I_2, I_3$ , are all equal.

3. With the notation of page 297, shew that in a triangle  $ABC$ , if the angle at  $C$  is a right angle,

$$r = s - c; \quad r_1 = s - b; \quad r_2 = s - a; \quad r_3 = s.$$

4. With the figure given on page 298, shew that if the circles whose centres are  $l, l_1, l_2, l_3$  touch  $BC$  at  $D, D_1, D_2, D_3$ , then

$$(i) \quad DD_2 = D_1D_3 = b.$$

$$(ii) \quad DD_3 = D_1D_2 = c.$$

$$(iii) \quad D_2D_3 = b + c.$$

$$(iv) \quad DD_1 = b - c.$$

5. Shew that the orthocentre and vertices of a triangle are the centres of the inscribed and escribed circles of the pedal triangle.

[See Ex. 20, p. 243.]

6. Given the base and vertical angle of a triangle, find the locus of the centre of the inscribed circle.

[See Ex. 36, p. 246.]

7. Given the base and vertical angle of a triangle, find the locus of the centre of the escribed circle which touches the base.

8. Given the base and vertical angle of a triangle, shew that the centre of the circumscribed circle is fixed.

9. Given the base  $BC$ , and the vertical angle  $A$  of a triangle, find the locus of the centre of the escribed circle which touches  $AC$ .

10. Given the base, the vertical angle, and the radius of the inscribed circle; construct the triangle.

11. Given the base, the vertical angle, and the radius of the escribed circle, (i) which touches the base, (ii) which touches one of the sides containing the vertical angle; construct the triangle.

12. Given the base, the vertical angle, and the point of contact with the base of the inscribed circle; construct the triangle.

13. Given the base, the vertical angle, and the point of contact with the base, or base produced, of an escribed circle; construct the triangle.

14. From an external point  $A$  two tangents  $AB, AC$  are drawn to a given circle; and the angle  $BAC$  is bisected by a straight line which meets the circumference in  $l$  and  $l_1$ ; shew that  $l$  is the centre of the circle inscribed in the triangle  $ABC$ , and  $l_1$  the centre of one of the escribed circles.

15.  $l$  is the centre of the circle inscribed in a triangle, and  $l_1, l_2, l_3$  the centres of the escribed circles; shew that  $ll_1, ll_2, ll_3$  are bisected by the circumference of the circumscribed circle.

16.  $ABC$  is a triangle, and  $l_2, l_3$  the centres of the escribed circles which touch  $AC$ , and  $AB$  respectively; shew that the points  $B, C, l_2, l_3$  lie upon a circle whose centre is on the circumference of the circle circumscribed about  $ABC$ .

17. With three given points as centres describe three circles touching one another two by two. How many solutions will there be?

18. Two tangents  $AB, AC$  are drawn to a given circle from an external point  $A$ ; and in  $AB, AC$  two points  $D$  and  $E$  are taken so that  $DE$  is equal to the sum of  $DB$  and  $EC$ : shew that  $DE$  touches the circle.

19. Given the perimeter of a triangle, and one angle in magnitude and position: shew that the opposite side always touches a fixed circle.

20. Given the centres of the three escribed circles; construct the triangle.

21. Given the centre of the inscribed circle, and the centres of two escribed circles; construct the triangle.

22. Given the vertical angle, perimeter, and the length of the bisector of the vertical angle; construct the triangle.

23. Given the vertical angle, perimeter, and altitude; construct the triangle.

24. Given the vertical angle, perimeter, and radius of the inscribed circle; construct the triangle.

25. Given the vertical angle, the radius of the inscribed circle, and the length of the perpendicular from the vertex to the base; construct the triangle.

26. Given the base, the difference of the sides containing the vertical angle, and the radius of the inscribed circle; construct the triangle. [See Ex. 10, p. 276.]

27. Given a vertex, the centre of the circumscribed circle, and the centre of the inscribed circle, construct the triangle.

28. In a triangle  $ABC$ ,  $I$  is the centre of the inscribed circle; shew that the centres of the circles circumscribed about the triangles  $BIC, CIA, AIB$  lie on the circumference of the circle circumscribed about the given triangle.

29. In a triangle  $ABC$ , the inscribed circle touches the base  $BC$  at  $D$ ; and  $r, r_1$  are the radii of the inscribed circle and of the escribed circle which touches  $BC$ : shew that  $r \cdot r_1 = BD \cdot DC$ .

30.  $ABC$  is a triangle,  $D, E, F$  the points of contact of its inscribed circle; and  $D'E'F'$  is the pedal triangle of the triangle  $DEF$ : shew that the sides of the triangle  $D'E'F'$  are parallel to those of  $ABC$ .

31. In a triangle  $ABC$  the inscribed circle touches  $BC$  at  $D$ . Shew that the circles inscribed in the triangles  $ABD, ACD$  touch one another.

## ON THE NINE-POINTS CIRCLE.

32. In any triangle the middle points of the sides, the feet of the perpendiculars drawn from the vertices to the opposite sides, and the middle points of the lines joining the orthocentre to the vertices are concyclic.

In the  $\triangle ABC$ , let  $X, Y, Z$  be the middle points of the sides  $BC, CA, AB$ ; let  $D, E, F$  be the feet of the perp<sup>s</sup> drawn to these sides from  $A, B, C$ ; let  $O$  be the orthocentre, and  $\alpha, \beta, \gamma$  the middle points of  $OA, OB, OC$ .

Then shall the nine points  $X, Y, Z, D, E, F, \alpha, \beta, \gamma$  be concyclic.

Join  $XY, XZ, X\alpha, Y\alpha, Z\alpha$ .

Now from the  $\triangle ABO$ , since  $AZ = ZB$ , and  $A\alpha = \alpha O$ , *Hyp.*  
 $\therefore Z\alpha$  is par<sup>l</sup> to  $BO$ . Ex. 2, p. 104.

And from the  $\triangle ABC$ , since  $BZ = ZA$ , and  $BX = XC$ , *Hyp.*  
 $\therefore ZX$  is par<sup>l</sup> to  $AC$ .

But  $BO$  produced makes a rt. angle with  $AC$ ; *Hyp.*  
 $\therefore$  the  $\angle XZ\alpha$  is a rt. angle.

Similarly, the  $\angle XY\alpha$  is a rt. angle. I. 29.  
 $\therefore$  the points  $X, Z, \alpha, Y$  are concyclic:

that is,  $\alpha$  lies on the  $\bigcirc^{ce}$  of the circle, which passes through  $X, Y, Z$ ; and  $X\alpha$  is a diameter of this circle.


Similarly it may be shewn that  $\beta$  and  $\gamma$  lie on the  $\bigcirc^{ce}$  of the circle which passes through  $X, Y, Z$ .

Again, since  $\alpha DX$  is a rt. angle, *Hyp.*  
 $\therefore$  the circle on  $X\alpha$  as diameter passes through  $D$ .

Similarly it may be shewn that  $E$  and  $F$  lie on the circumference of the same circle.

$\therefore$  the points  $X, Y, Z, D, E, F, \alpha, \beta, \gamma$  are concyclic. Q.E.D.

From this property the circle which passes through the middle points of the sides of a triangle is called the **Nine-Points Circle**; many of its properties may be derived from the fact of its being the circle circumscribed about the pedal triangle.






33. To prove that

- (i) the centre of the nine-points circle is the middle point of the straight line which joins the orthocentre to the circumscribed centre.
- (ii) the radius of the nine-points circle is half the radius of the circumscribed circle.
- (iii) the centroid is collinear with the circumscribed centre, the nine-points centre, and the orthocentre.

In the  $\triangle ABC$ , let  $X, Y, Z$  be the middle points of the sides;  $D, E, F$  the feet of the perp<sup>s</sup>;  $O$  the orthocentre;  $S$  and  $N$  the centres of the circumscribed and nine-points circles respectively.



(i) To prove that  $N$  is the middle point of  $SO$ .

It may be shewn that the perp. to  $XD$  from its middle point bisects  $SO$ ;

Ex. 14, p. 106.

Similarly the perp. to  $EY$  at its middle point bisects  $SO$  :

that is, these perp<sup>s</sup> intersect at the middle point of  $SO$  :

And since  $XD$  and  $EY$  are chords of the nine-points circle,  
 $\therefore$  the intersection of the lines which bisect  $XD$  and  $EY$  at rt. angles is its centre :

III. 1.

$\therefore$  the centre  $N$  is the middle point of  $SO$

(ii) To prove that the radius of the nine-points circle is half the radius of the circumscribed circle.

By the last Proposition,  $Xa$  is a diameter of the nine-points circle.

$\therefore$  the middle point of  $Xa$  is its centre :

but the middle point of  $SO$  is also the centre of the nine-points circle.

(Proved.)

Hence  $Xa$  and  $SO$  bisect one another at  $N$ .

Then from the  $\triangle^s SNX, ONa$ ,  
 Because  $\left\{ \begin{array}{l} SN = ON, \\ \text{and } NX = Na, \\ \text{and the } \angle SNX = \text{the } \angle ONa; \end{array} \right.$  I. 15.  
 $\therefore SX = Oa$  I. 4.  
 $= Aa$ .

And  $SX$  is also par<sup>l</sup> to  $Aa$ ,  
 $\therefore SA = Xa$ . I. 33.

But  $SA$  is a radius of the circumscribed circle ;  
 and  $Xa$  is a diameter of the nine-points circle ;

$\therefore$  the radius of the nine-points circle is half the radius of the circumscribed circle.

(iii) To prove that the centroid is collinear with points S, N, O.

Join AX and draw  $ag$  par<sup>l</sup> to SO.

Let AX meet SO at G.

Then from the  $\triangle AGO$ , since  $Aa = aO$ , and  $ag$  is par<sup>l</sup> to OG,

$$\therefore Ag = gG. \quad \text{Ex. 1, p. 104.}$$

And from the  $\triangle Xag$ , since  $aN = NX$ , and NG is par<sup>l</sup> to  $ag$ ,

$$\therefore gG = GX.$$

$$\therefore AG = \frac{2}{3} \text{ of } AX;$$

$\therefore$  G is the centroid of the triangle ABC. Ex. 4, p. 113.

That is, the centroid is collinear with the points S, N, O. Q.E.D.

34. Given the base and vertical angle of a triangle, find the locus of the centre of the nine-points circle.

35. The nine-points circle of any triangle ABC, whose orthocentre is O, is also the nine-points circle of each of the triangles AOB, BOC, COA.

36. If  $l, l_1, l_2, l_3$  are the centres of the inscribed and escribed circles of a triangle ABC, then the circle circumscribed about ABC is the nine-points circle of each of the four triangles formed by joining three of the points  $l, l_1, l_2, l_3$ .

37. All triangles which have the same orthocentre and the same circumscribed circle, have also the same nine-points circle.

38. Given the base and vertical angle of a triangle, shew that one angle and one side of the pedal triangle are constant.

39. Given the base and vertical angle of a triangle, find the locus of the centre of the circle which passes through the three escribed centres.

NOTE. For another important property of the Nine-points Circle see Miscellaneous Examples on Book VI., Ex. 60.

## II. MISCELLANEOUS EXAMPLES.

1. If four circles are described to touch every three sides of a quadrilateral, shew that their centres are concyclic.

2. If the straight lines which bisect the angles of a rectilineal figure are concurrent, a circle may be inscribed in the figure.

3. Within a given circle describe three equal circles touching one another and the given circle.

4. The perpendiculars drawn from the centres of the three escribed circles of a triangle to the sides which they touch, are concurrent.

5. Given an angle and the radii of the inscribed and circumscribed circles; construct the triangle.

6. Given the base, an angle at the base, and the distance between the centre of the inscribed circle and the centre of the escribed circle which touches the base; construct the triangle.

7. In a given circle inscribe a triangle such that two of its sides may pass through two given points, and the third side be of given length.

8. In any triangle  $ABC$ ,  $I, I_1, I_2, I_3$  are the centres of the inscribed and escribed circles, and  $S_1, S_2, S_3$  are the centres of the circles circumscribed about the triangles  $BIC, CIA, AIB$ : shew that the triangle  $S_1S_2S_3$  has its sides parallel to those of the triangle  $I_1I_2I_3$ , and is one-fourth of it in area: also that the triangles  $ABC$  and  $S_1S_2S_3$  have the same circumscribed circle.

9.  $O$  is the orthocentre of a triangle  $ABC$ : shew that

$$AO^2 + BC^2 = BO^2 + CA^2 = CO^2 + AB^2 = d^2,$$

where  $d$  is the diameter of the circumscribed circle.

10. If from any point within a regular polygon of  $n$  sides perpendiculars are drawn to the sides, the sum of the perpendiculars is equal to  $n$  times the radius of the inscribed circle.

11. The sum of the perpendiculars drawn from the vertices of a regular polygon of  $n$  sides on any straight line is equal to  $n$  times the perpendicular drawn from the centre of the inscribed circle.

12. The area of a cyclic quadrilateral is independent of the order in which the sides are placed in the circle.

13. Given the orthocentre, the centre of the nine-points circle, and the middle point of the base; construct the triangle.

14. Of all polygons of a given number of sides, which may be inscribed in a given circle, that which is regular has the maximum area and the maximum perimeter.

15. Of all polygons of a given number of sides circumscribed about a given circle, that which is regular has the minimum area and the minimum perimeter.

16. Given the vertical angle of a triangle in position and magnitude, and the sum of the sides containing it: find the locus of the centre of the circumscribed circle.

17.  $P$  is any point on the circumference of a circle circumscribed about an equilateral triangle  $ABC$ : shew that  $PA^2 + PB^2 + PC^2$  is constant.

\* \* *Book V. is now very rarely read. The subject-matter, so far as it is introductory to Book VI., is dealt with in a simpler manner at page 317, in the chapter called 'Elementary Principles of Proportion.'* The student is advised to proceed at once to that chapter, leaving *Book V.* in its stricter form to be studied at a later stage, if it is thought desirable.

## BOOK V.

**Book V.** treats of Ratio and Proportion, and the method adopted is such as to place these subjects on a basis independent of arithmetical principles.

The following notation will be employed throughout this section.

Capital letters,  $A, B, C, \dots$  will be used to denote the magnitudes themselves, *not any numerical or algebraical measures of them*, and small letters,  $m, n, p, \dots$  will be used to denote whole numbers. Also it will be assumed that multiplication, in the sense of repeated addition, can be applied to any magnitude, so that  $m \cdot A$  or  $mA$  will denote the magnitude  $A$  taken  $m$  times.

The symbol  $>$  will be used for the words *greater than*, and  $<$  for *less than*.

### DEFINITIONS.

*Definition 1.* One magnitude is said to be a **multiple** of another, when the first contains the second an *exact* number of times.

*Definition 2.* One magnitude is said to be a **submultiple** of another, when the first is contained an *exact* number of times in the second.

The following properties of multiples will be assumed as self-evident.

- (1)  $mA >, =, \text{ or } < nB$  according as  $A >, =, \text{ or } < B$ ; and conversely.
- (2)  $mA + mB + \dots = m(A + B + \dots)$ .
- (3) If  $A > B$ , then  $mA - nB = m(A - B)$ .
- (4)  $mA + nA + \dots = (m + n + \dots)A$ .
- (5) If  $m > n$ , then  $mA - nA = (m - n)A$ .
- (6)  $m \cdot nA = mn, A = nm \cdot A = n \cdot mA$ .

*Definition 3.* The **Ratio** of one magnitude to another of the same kind is the relation which the first bears to the second in respect of *quantuplicity*.

The ratio of  $A$  to  $B$  is denoted thus,  $A : B$ ; and  $A$  is called the **antecedent**,  $B$  the **consequent** of the ratio.

The term *quantuplicity* denotes the capacity of the first magnitude to contain the second with or without remainder.

If the magnitudes are commensurable, their quantuplicity may be expressed *numerically* by observing what multiples of the two magnitudes are equal to one another.

Thus if  $A = ma$ , and  $B = na$ , it follows that  $nA = mB$ . In this case  $A = \frac{m}{n}B$ , and the quantuplicity of  $A$  with respect to  $B$  is the arithmetical fraction  $\frac{m}{n}$ .

But if the magnitudes are incommensurable, no multiple of the first can be equal to any multiple of the second, and therefore the quantuplicity of one with respect to the other cannot exactly be expressed numerically: in this case it is determined by examining how the multiples of one magnitude are distributed among the multiples of the other.

Thus, let all the multiples of  $A$  be formed, the scale extending *ad infinitum*; also let all the multiples of  $B$  be formed and placed in their proper order of magnitude among the multiples of  $A$ . This forms the relative scale of the two magnitudes, and the quantuplicity of  $A$  with respect to  $B$  is estimated by examining how the multiples of  $A$  are distributed among those of  $B$  in their relative scale.

In other words, the ratio of  $A$  to  $B$  is known, if for all integral values of  $m$  we know the multiples  $nB$  and  $(n+1)B$  between which  $mA$  lies.

In the case of two given magnitudes  $A$  and  $B$ , the relative scale of multiples is definite, and is different from that of  $A$  to  $C$ , if  $C$  differs from  $B$  by any magnitude however small.

For let  $D$  be the difference between  $B$  and  $C$ ; then however small  $D$  may be, it will be possible to find a number  $m$  such that  $mD > A$ . In this case,  $mB$  and  $mC$  would differ by a magnitude greater than  $A$ , and therefore could not lie between the same two multiples of  $A$ ; so that after a certain point the relative scale of  $A$  and  $B$  would differ from that of  $A$  and  $C$ .

*Definition 4.* Magnitudes are said to have a ratio to one another, when the less can be multiplied so as to exceed the other.

*Definition 5.* The ratio of one magnitude to another is equal to that of a third magnitude to a fourth, when if any equimultiples whatever of the antecedents of the ratios are taken, and also any equimultiples whatever of the consequents, the multiple of one antecedent is greater than, equal to, or less than that of its consequent, according as the multiple of the other antecedent is greater than, equal to, or less than that of its consequent.

Thus the ratio  $A$  to  $B$  is equal to that of  $C$  to  $D$  when  $mC >$ ,  $=$ , or  $< nD$  according as  $mA >$ ,  $=$ , or  $< nB$ , whatever whole numbers  $m$  and  $n$  may be.

Again, let  $m$  be any whole number whatever, and  $n$  another whole number determined in such a way that either  $mA$  is equal to  $nB$ , or  $mA$  lies between  $nB$  and  $(n+1)B$ ; then the definition asserts that the ratio of  $A$  to  $B$  is equal to that of  $C$  to  $D$  if  $mC = nD$  when  $mA = nB$ ; or if  $mC$  lies between  $nD$  and  $(n+1)D$  when  $mA$  lies between  $nB$  and  $(n+1)B$ .

In other words, the ratio of  $A$  to  $B$  is equal to that of  $C$  to  $D$  when the multiples of  $A$  are distributed among those of  $B$  in the same manner as the multiples of  $C$  are distributed among those of  $D$ .

When the ratio of  $A$  to  $B$  is equal to that of  $C$  to  $D$  the four magnitudes are called **proportionals**. This is expressed by saying " $A$  is to  $B$  as  $C$  is to  $D$ ," and the **proportion** is written

$$A : B :: C : D, \text{ or } A : B = C : D.$$

$A$  and  $D$  are called the **extremes**,  $B$  and  $C$  the **means**; also  $D$  is said to be a **fourth proportional** to  $A$ ,  $B$ , and  $C$ .

*Definition 6.* Two terms in a proportion are said to be **homologous** when they are both antecedents, or both consequents of the ratios.

*Definition 7.* The ratio of one magnitude to another is greater than that of a third magnitude to a fourth, when it is possible to find equimultiples of the antecedents and equimultiples of the consequents such that while the multiple of the antecedent of the first ratio is greater than, or equal to, that of its consequent, the multiple of the antecedent of the second is not greater, or is less, than that of its consequent.

This definition asserts that if whole numbers  $m$  and  $n$  can be found such that while  $mA$  is greater than  $nB$ ,  $mC$  is not greater than  $nD$ , or while  $mA = nB$ ,  $mC$  is less than  $nD$ , then the ratio of  $A$  to  $B$  is greater than that of  $C$  to  $D$ .

If  $A$  is equal to  $B$ , the ratio of  $A$  to  $B$  is called a **ratio of equality**.

If  $A$  is greater than  $B$ , the ratio of  $A$  to  $B$  is called a **ratio of greater inequality**.

If  $A$  is less than  $B$ , the ratio of  $A$  to  $B$  is called a **ratio of less inequality**.

*Definition 8.* Two ratios are said to be **reciprocal** when the antecedent and consequent of one are the consequent and antecedent of the other respectively; thus  $B : A$  is the reciprocal of  $A : B$ .

*Definition 9.* Three magnitudes of the same kind are said to be **proportionals**, when the ratio of the first to the second is equal to that of the second to the third.

Thus  $A, B, C$  are proportionals if

$$A : B :: B : C.$$

$B$  is called a **mean proportional** to  $A$  and  $C$ , and  $C$  is called a **third proportional** to  $A$  and  $B$ .

*Definition 10.* Three or more magnitudes are said to be in **continued proportion** when the ratio of the first to the second is equal to that of the second to the third, and the ratio of the second to the third is equal to that of the third to the fourth, and so on.

*Definition 11.* When there are any number of magnitudes of the same kind, the first is said to have to the last the **ratio compounded** of the ratios of the first to the second, of the second to the third, and so on up to the ratio of the last but one to the last magnitude.

For example, if  $A, B, C, D, E$  be magnitudes of the same kind,  $A : E$  is the ratio compounded of the ratios  $A : B, B : C, C : D$ , and  $D : E$ .

This is sometimes expressed by the following notation :

$$A : E = \left\{ \begin{array}{l} A : B \\ B : C \\ C : D \\ D : E. \end{array} \right.$$

*Definition 12.* If there are any number of ratios, and a set of magnitudes is taken such that the ratio of the first to the second is equal to the first ratio, and the ratio of the second to the third is equal to the second ratio, and so on, then the first of the set of magnitudes is said to have to the last the **ratio compounded** of the given ratios.

Thus, if  $A : B, C : D, E : F$  be given ratios, and if  $P, Q, R, S$  be magnitudes taken so that

$$\begin{array}{l} P : Q :: A : B, \\ Q : R :: C : D, \\ R : S :: E : F; \end{array}$$

then

$$P : S = \left\{ \begin{array}{l} A : B \\ C : D \\ E : F. \end{array} \right.$$

*Definition 13.* When three magnitudes are proportionals, the first is said to have to the third the **duplicate ratio** of that which it has to the second.

Thus if  $A : B :: B : C,$

then  $A$  is said to have to  $C$  the duplicate ratio of that which it has to  $B$ .

Since  $A : C = \begin{cases} A : B \\ B : C, \end{cases}$

it is clear that the ratio compounded of two equal ratios is the duplicate ratio of either of them.

*Definition 14.* When four magnitudes are in *continued proportion*, the first is said to have to the fourth the **triplicate ratio** of that which it has to the second.

It may be shewn as above that the ratio compounded of three equal ratios is the triplicate ratio of any one of them.

## PROPOSITIONS.

*Obs.* Of the propositions of Book V., which, it may be noticed are all theorems, we here give only the more important.

### PROPOSITION 1.

*Ratios which are equal to the same ratio are equal to one another.*

Let  $A : B :: P : Q,$  and also  $C : D :: P : Q;$  then shall  $A : B :: C : D.$

For it is evident that two scales or arrangements of multiples which agree in every respect with a third scale, will agree with one another.

### PROPOSITION 2.

*If two ratios are equal, the antecedent of the second is greater than, equal to, or less than its consequent according as the antecedent of the first is greater than, equal to, or less than its consequent.*

Let  $A : B :: C : D,$

then  $C >, =, \text{ or } < D,$

according as  $A >, =, \text{ or } < B.$

This follows at once from Def. 5, by taking  $m$  and  $n$  each equal to unity.



## PROPOSITION 3.

**Invertendo or Inversely.** *If two ratios are equal, their reciprocals are equal.*

Let  $A : B :: C : D$ ,  
then shall  $B : A :: D : C$ .

For, by hypothesis, the multiples of  $A$  are distributed among those of  $B$  in the same manner as the multiples of  $C$  are among those of  $D$ .

therefore also, the multiples of  $B$  are distributed among those of  $A$  in the same manner as the multiples of  $D$  are among those of  $C$ .

That is,  $B : A :: D : C$ .

**NOTE.** This proposition is sometimes enunciated thus:

*If four magnitudes are proportionals, they are also proportionals when taken inversely.*

## PROPOSITION 4.

*Equal magnitudes have the same ratio to the same magnitude; and the same magnitude has the same ratio to equal magnitudes.*

Let  $A, B, C$  be three magnitudes of the same kind, and let  $A$  be equal to  $B$ ;

then shall  $A : C :: B : C$   
and  $C : A :: C : B$ .

Since  $A = B$ , their multiples are identical and therefore are distributed in the same way among the multiples of  $C$ .

$\therefore A : C :: B : C$ , Def. 5.  
 $\therefore$  also, *invertendo*,  $C : A :: C : B$ . v. 3.

## PROPOSITION 5.

*Of two unequal magnitudes, the greater has a greater ratio to a third magnitude than the less has; and the same magnitude has a greater ratio to the less of two magnitudes than it has to the greater.*

First, let  $A$  be  $> B$ ;  
then shall  $A : C$  be  $> B : C$ .

Since  $A > B$ , it will be possible to find  $m$  such that  $mA$  exceeds  $mB$  by a magnitude greater than  $C$ ;

hence if  $mA$  lies between  $nC$  and  $(n+1)C$ ,  $mB < nC$ ;

and if  $mA = nC$ , then  $mB < nC$ ;

$\therefore A : C > B : C$ . Def. 7.

Secondly, let  $B$  be  $< A$  ;  
 then shall  $C : B$  be  $> C : A$ .  
 For taking  $m$  and  $n$  as before,  
 $nC > mB$ , while  $nC$  is not  $> mA$  ;  
 $\therefore C : B > C : A$ .

Def. 7.

PROPOSITION 6.

*Magnitudes which have the same ratio to the same magnitude are equal to one another ; and those to which the same magnitude has the same ratio are equal to one another.*

First, let  $A : C :: B : C$  ;  
 then shall  $A = B$ .  
 For if  $A > B$ , then  $A : C > B : C$ ,  
 and if  $B > A$ , then  $B : C > A : C$ ,  
 which contradict the hypothesis ;  
 $\therefore A = B$ .

v. 5.

Secondly, let  $C : A :: C : B$  ;  
 then shall  $A = B$ .  
 Because  $C : A :: C : B$ ,  
 $\therefore$  *invertendo*,  $A : C :: B : C$ ,  
 $\therefore A = B$ ,

v. 3.

by the first part of the proof.

PROPOSITION 7.

*That magnitude which has a greater ratio than another has to the same magnitude is the greater of the two ; and that magnitude to which the same has a greater ratio than it has to another magnitude is the less of the two.*

First, let  $A : C$  be  $> B : C$  ;  
 then shall  $A$  be  $> B$ .  
 For if  $A = B$ , then  $A : C :: B : C$ ,  
 which is contrary to the hypothesis.  
 And if  $A < B$ , then  $A : C < B : C$  ;  
 which is contrary to the hypothesis ;  
 $\therefore A > B$ .

v. 4.

v. 5.

Secondly, let  $C : A$  be  $> C : B$ ;  
 then shall  $A$  be  $< B$ .  
 For if  $A = B$ , then  $C : A :: C : B$ , v. 4.  
 which is contrary to the hypothesis.  
 And if  $A > B$ , then  $C : A < C : B$ ; v. 5.  
 which is contrary to the hypothesis ;  
 $\therefore A < B$ .

## PROPOSITION 8.

*Magnitudes have the same ratio to one another which their equimultiples have.*

Let  $A, B$  be two magnitudes ;  
 then shall  $A : B :: mA : mB$ .  
 If  $p, q$  be any two whole numbers,  
 then  $m \cdot pA >, =, \text{ or } < m \cdot qB$   
 according as  $pA >, =, \text{ or } < qB$ .  
 But  $m \cdot pA = p \cdot mA$ , and  $m \cdot qB = q \cdot mB$ ;  
 $\therefore p \cdot mA >, =, \text{ or } < q \cdot mB$   
 according as  $pA >, =, \text{ or } < qB$ ;  
 $\therefore A : B :: mA : mB$ . Def. 5.  
 COR. Let  $A : B :: C : D$ .  
 Then since  $A : B :: mA : mB$ ,  
 and  $C : D :: nC : nD$ ;  
 $\therefore mA : mB :: nC : nD$ . v. 1.

## PROPOSITION 9.

*If two ratios are equal, and any equimultiples of the antecedents and also of the consequents are taken, the multiple of the first antecedent has to that of its consequent the same ratio as the multiple of the other antecedent has to that of its consequent.*

Let  $A : B :: C : D$ ;  
 then shall  $mA : nB :: mC : nD$ .  
 Let  $p, q$  be any two whole numbers ;  
 then because  $A : B :: C : D$ ,  
 $pm \cdot C >, =, \text{ or } < qn \cdot D$   
 according as  $pm \cdot A >, =, \text{ or } < qn \cdot B$ , Def. 5.  
 that is,  $p \cdot mC >, =, \text{ or } < q \cdot nD$ ,  
 according as  $p \cdot mA >, =, \text{ or } < q \cdot nB$ ;  
 $\therefore mA : nB :: mC : nD$ . Def. 5.

PROPOSITION 10.

*If four magnitudes of the same kind are proportionals, the first is greater than, equal to, or less than the third, according as the second is greater than, equal to, or less than the fourth.*

Let  $A, B, C, D$  be four magnitudes of the same kind such that

$$A : B :: C : D ;$$

then  $A >, =, \text{ or } < C$

according as  $B >, =, \text{ or } < D$ .

If  $B > D$ , then  $A : B < A : D$ ; v. 5

but  $A : B :: C : D$ ;

$$\therefore C : D < A : D ;$$

$$\therefore A : D > C : D ;$$

$$\therefore A > C. \quad \text{v. 7.}$$

Similarly it may be shewn that

if  $B < D$ , then  $A < C$ ,

and if  $B = D$ , then  $A = C$ .

PROPOSITION 11.

**Alternando or Alternately.** *If four magnitudes of the same kind are proportionals, they are also proportionals when taken alternately.*

Let  $A, B, C, D$  be four magnitudes of the same kind such that

$$A : B :: C : D ;$$

then shall

$$A : C :: B : D.$$

Because  $A : B :: mA : mB$ , v. 8.

and  $C : D :: nC : nD$ ;

$$\therefore mA : mB :: nC : nD. \quad \text{v. 1.}$$

$$\therefore mA >, =, \text{ or } < nC$$

according as  $mB >, =, \text{ or } < nD$ . v. 10.

And  $m$  and  $n$  are any whole numbers ;

$$\therefore A : C :: B : D. \quad \text{Def. 5.}$$

## PROPOSITION 12.

**Addendo.** *If any number of magnitudes of the same kind are proportionals, as one of the antecedents is to its consequent, so is the sum of the antecedents to the sum of the consequents.*

Let  $A, B, C, D, E, F, \dots$  be magnitudes of the same kind such that

$$A : B :: C : D :: E : F :: \dots ;$$

then shall  $A : B :: A + C + E + \dots : B + D + F + \dots$ .

Because  $A : B :: C : D :: E : F :: \dots$ ,

$\therefore$  according as  $mA >, =, \text{ or } < nB$ ,  
so is  $mC >, =, \text{ or } < nD$ ,  
and  $mE >, =, \text{ or } < nF$ ,

$\therefore$  so is  $mA + mC + mE + \dots >, =, \text{ or } < nB + nD + nF + \dots$   
or  $m(A + C + E + \dots) >, =, \text{ or } < n(B + D + F + \dots)$ ;

and  $m$  and  $n$  are any whole numbers;

$\therefore A : B :: A + C + E + \dots : B + D + F + \dots$  Def. 5.

## PROPOSITION 13.

**Componendo.** *If four magnitudes are proportionals, the sum of the first and second is to the second as the sum of the third and fourth is to the fourth.*

Let  $A : B :: C : D$ ;

then shall

$$A + B : B :: C + D : D.$$

If  $m$  be any whole number, it is possible to find another number  $n$  such that  $mA = nB$ , or lies between  $nB$  and  $(n+1)B$ ,

$\therefore mA + mB = mB + nB$ , or lies between  $mB + nB$  and  $mB + (n+1)B$ .

But  $mA + mB = m(A + B)$ , and  $mB + nB = (m+n)B$ ;

$\therefore m(A + B) = (m+n)B$ , or lies between  $(m+n)B$  and  $(m+n+1)B$ .

Also because  $A : B :: C : D$ ,

$\therefore mC = nD$ , or lies between  $nD$  and  $(n+1)D$ ; Def. 5.

$\therefore m(C + D) = (m+n)D$  or lies between  $(m+n)D$  and  $(m+n+1)D$ ;

that is, the multiples of  $C + D$  are distributed among those of  $D$  in the same way as the multiples of  $A + B$  among those of  $B$ ;

$\therefore A + B : B :: C + D : D$ .

**Dividendo.** In the same way it may be proved that

$$A - B : B :: C - D : D,$$

or  $B - A : B :: D - C : D$ ,  
according as  $A$  is  $>$  or  $<$   $B$ .

PROPOSITION 14.

**Ex Aequali.** *If there are two sets of magnitudes, such that the first is to the second of the first set as the first to the second of the other set, and the second to the third of the first set as the second to the third of the other, and so on to the last magnitude : then the first is to the last of the first set as the first to the last of the other.*

*First*, let there be three magnitudes  $A, B, C$  of one set, and three,  $P, Q, R$ , of another set,

$$\text{and let } A : B :: P : Q,$$

$$\text{and } B : C :: Q : R;$$

$$\text{then shall } A : C :: P : R.$$

$$\text{Because } A : B :: P : Q,$$

$$\therefore mA : mB :: mP : mQ; \quad \text{v. 8, Cor.}$$

$$\text{and because } B : C :: Q : R,$$

$$\therefore mB : nC :: mQ : nR, \quad \text{v. 9.}$$

$$\therefore, \text{ invertendo,} \quad nC : mB :: nR : mQ. \quad \text{v. 3.}$$

Now, if

$$mA > nC,$$

$$\text{then } mA : mB > nC : mB; \quad \text{v. 5.}$$

$$\therefore mP : mQ > nR : mQ,$$

$$\text{and } \therefore mP > nR. \quad \text{v. 7.}$$

Similarly  $mP =$  or  $< nR$  according as  $mA =$  or  $< nC$ .

$$\therefore A : C :: P : R. \quad \text{Def. 5.}$$

*Secondly*, let there be any number of magnitudes,  $A, B, C, \dots L, M$ , of one set, and the same number  $P, Q, R, \dots Y, Z$ , of another set, such that

$$A : B :: P : Q,$$

$$B : C :: Q : R,$$

$$\dots :: \dots$$

$$L : M :: Y : Z;$$

$$\text{then shall } A : M :: P : Z.$$

$$\text{For } A : C :: P : R,$$

$$\text{and } C : D :: R : S;$$

$$\therefore \text{ by the first case } A : D :: P : S,$$

$$\text{and so on, until finally } A : M = P : Z.$$

*Proved.*

*Hyp.*

**COROLLARY.**

$$\text{If } A : B :: P : Q,$$

$$\text{and } B : C :: R : P:$$

$$\text{then } A : C :: R : Q.$$

## PROPOSITION 15.

If  $A : B :: X : Y$ ,  
 and  $C : B :: Z : Y$ ;  
 then shall  $A + C : B :: X + Z : Y$ .

	For since $C : B :: Z : Y$ ,	<i>Hyp.</i>
$\therefore$ , <i>invertendo</i> ,	$B : C :: Y : Z$ .	v. 3.
	Also $A : B :: X : Y$ ,	
$\therefore$ , <i>ex æquali</i> ,	$A : C :: X : Z$ ,	v. 14.
$\therefore$ , <i>componendo</i> ,	$A + C : C :: X + Z : Z$ .	v. 13.
	Again, $C : B :: Z : Y$ ,	<i>Hyp.</i>
$\therefore$ , <i>ex æquali</i> ,	$A + C : B :: X + Z : Y$ .	v. 14.

## PROPOSITION 16.

*If two ratios are equal, their duplicate ratios are equal.*

Let  $A : B :: C : D$ ;

then shall the duplicate ratio of  $A$  to  $B$  be equal to that of  $C$  to  $D$ .

Let  $X$  be a third proportional to  $A$  and  $B$ , and  $Y$  a third proportional to  $C$  and  $D$ ,

so that  $A : B :: B : X$ , and  $C : D :: D : Y$ ;

then because  $A : B :: C : D$ ,

$\therefore B : X :: D : Y$ ;

$\therefore$ , *ex æquali*,

$A : X :: C : Y$ .

But  $A : X$  and  $C : Y$  are respectively the duplicate ratios of

$A : B$  and  $C : D$ ,

*Def. 13.*

$\therefore$  the duplicate ratio of  $A : B =$  that of  $C : D$ .

NOTE. The converse of this theorem may be readily proved; namely,

*If the duplicates of two ratios are equal, the ratios themselves are equal.*

## ELEMENTARY PRINCIPLES OF PROPORTION.

### INTRODUCTION TO BOOK VI.

1. The first four books of Euclid deal with the absolute equality or inequality of geometrical magnitudes. In Book VI. such magnitudes are compared by considering their *ratio* or *relative greatness*.

2. The meaning of the words *ratio* and *proportion* in their simplest arithmetical sense may be given as follows:

(i) *The ratio of one number to another is the multiple or fraction which the first is of the second.*

(ii) *Four numbers are in proportion when the ratio of the first to the second is equal to the ratio of the third to the fourth*

3. These definitions are however not strictly applicable to the purposes of Pure Geometry, for the following reasons:

(i) Pure Geometry deals only with magnitudes *as represented by diagrams*, without measuring them in terms of a common unit: in other words, it makes no use of *number* for the purpose of comparing magnitudes.

(ii) It commonly happens that Geometrical magnitudes of the same kind are *incommensurable*, that is, they are such that it is impossible to express them *exactly* in terms of some common unit. Nevertheless it is always possible to express the arithmetical ratio of two such magnitudes *within any required degree of accuracy*. [See Note, p. 131: also Hall and Knight's *Elementary Algebra*, Art. 289.]



4. Accordingly, the object of Euclid's Fifth Book is to establish the Theory of Proportion on a basis independent of *number*. But as Book V. is now very rarely read, we propose here merely to illustrate *algebraically* such principles of proportion as are required before proceeding to Book VI. The strict treatment of the subject given in Book V. may be studied at a later stage, if it is thought desirable.

*Obs.* In what follows the symbol  $>$  will be used for the words *greater than*, and  $<$  for *less than*.

5. The following definitions are selected from Book V.

*Definition 1.* One magnitude is said to be a **multiple** of another, when the first contains the second an *exact* number of times.

Thus  $ma$  is a multiple of  $a$ , if  $m$  is any whole number.

*Definition 2.* One magnitude is said to be a **submultiple** of another, when the first is contained in the second an *exact* number of times.

Thus  $\frac{a}{m}$  is a submultiple of  $a$ , if  $m$  is any whole number.

*Definition 3.* The **ratio** of one magnitude to another of *the same kind* is the relation which the first bears to the second in regard to quantity; this is measured by the fraction which the first is of the second.

Thus if two such magnitudes contain  $a$  and  $b$  units respectively, the ratio of the first to the second is expressed by the fraction  $\frac{a}{b}$ .

The ratio of  $a$  to  $b$  is generally denoted thus,  $a : b$ ; and  $a$  is called the **antecedent** and  $b$  the **consequent** of the ratio.

The two magnitudes compared in a ratio must be *of the same kind*; for example, both must be lines, or both angles, or both areas. It is clearly impossible to compare the *length* of a straight line with a magnitude of a different kind, such as the *area* of a triangle.

*Definition 5.* Four quantities are in **proportion**, when the ratio of the *first* to the *second* is equal to the ratio of the *third* to the *fourth*.

When the ratio of  $a$  to  $b$  is equal to that of  $x$  to  $y$ , the four magnitudes are called **proportionals**. This is expressed by saying " $a$  is to  $b$  as  $x$  is to  $y$ ," and the proportion is written

$$a : b :: x : y ;$$

$$\text{or } a : b = x : y.$$

Here  $a$  and  $y$  are called the **extremes**, and  $b$  and  $x$  the **means**.

(i) **Algebraical Test of Proportion.** The ratios  $a : b$  and  $x : y$  may be expressed algebraically by the fractions  $\frac{a}{b}$  and  $\frac{x}{y}$ ; thus the four magnitudes  $a, b, x, y$  are in proportion if

$$\frac{a}{b} = \frac{x}{y}$$

(ii) **Geometrical Test of Proportion.** The ratio of one magnitude to another is equal to that of a third magnitude to a fourth, when if any equimultiples whatever of the antecedents of the ratios are taken, and also any equimultiples whatever of the consequents, the multiple of one antecedent is greater than, equal to, or less than that of its consequent, according as the multiple of the other antecedent is greater than, equal to, or less than that of its consequent.

Thus the ratio of  $a$  to  $b$  is equal to that of  $x$  to  $y$ , that is to say,

$a, b, x, y$  are in proportion,

if  $mx >, =, \text{ or } < ny,$

according as  $ma >, =, \text{ or } < nb,$

whatever whole numbers  $m$  and  $n$  may be.

**NOTE.** The Algebraical and Geometrical Tests of Proportion, though differing widely in method, really determine the same property; for each may be deduced from the other. This is fully explained on the following page.

COMPARISON BETWEEN THE ALGEBRAICAL AND GEOMETRICAL  
TESTS OF PROPORTION.

(i) *If a, b, x, y satisfy the Algebraical test of proportion, to shew that they also satisfy the geometrical test.*

By hypothesis  $\frac{a}{b} = \frac{x}{y}$ ;

and, multiplying both sides by  $\frac{m}{n}$ , where  $m$  and  $n$  are any whole

numbers, we obtain  $\frac{ma}{nb} = \frac{mx}{ny}$ ;

thus these fractions are both improper, or both proper, or both equal to unity;

hence  $mx >$ ,  $=$ , or  $< ny$ , according as  $ma >$ ,  $=$ , or  $< nb$ , which is the Geometrical test of proportion.

(ii) *If a, b, x, y satisfy the Geometrical test of proportion, to shew that they also satisfy the Algebraical test.*

By hypothesis  $mx >$ ,  $=$ , or  $< ny$ , according as  $ma >$ ,  $=$ , or  $< nb$ , it is required to prove that

$$\frac{a}{b} = \frac{x}{y}$$

If  $\frac{a}{b}$  is not equal to  $\frac{x}{y}$ , one of them must be the greater.

Suppose  $\frac{a}{b} > \frac{x}{y}$ ; then it will be possible to find some fraction  $\frac{n}{m}$  which lies between them,  $n$  and  $m$  being positive integers.

Hence  $\frac{a}{b} > \frac{n}{m}$ , .....(1)

and  $\frac{x}{y} < \frac{n}{m}$ . .....(2)

From (1),  $ma > nb$ ;

from (2),  $mx < ny$ ;

and these contradict the hypothesis.

Therefore  $\frac{a}{b}$  and  $\frac{x}{y}$  are not unequal; that is  $\frac{a}{b} = \frac{x}{y}$ .

*Definition 6.* Two terms in a proportion are said to be **homologous**, when they are both antecedents or both consequents of the ratios.

Thus if  $a : b :: x : y$ ,

$a$  and  $x$  are homologous; also  $b$  and  $y$  are homologous,

*Definition 8.* Two ratios are said to be **reciprocal**, when the antecedent and consequent of one are respectively the consequent and antecedent of the other.

Thus  $b : a$  is the reciprocal of  $a : b$ .

*Definition 9.* Three magnitudes of the same kind are said to be **proportionals**, when the ratio of the *first* to the *second* is equal to that of the *second* to the *third*.

Thus  $a, b, c$  are proportionals if

$$a : b :: b : c.$$

Here  $b$  is called a **mean proportional** to  $a$  and  $c$ ; and  $c$  is called a **third proportional** to  $a$  and  $b$ .

When *four* magnitudes are in proportion, namely when

$$a : b :: c : d,$$

then  $d$  is called a **fourth proportional** to  $a, b$ , and  $c$ .

*Definition 10.* A series of magnitudes of the same kind are said to be in **continued proportion**, when the ratios of the *first* to the *second*, of the *second* to the *third*, of the *third* to the *fourth*, and so on, are all equal.

Thus  $a, b, c, d, e$  are in continued proportion, if

$$a : b = b : c = c : d = d : e;$$

that is, if  $\frac{a}{b} = \frac{b}{c} = \frac{c}{d} = \frac{d}{e}$ .

*Definition 11.* When there are any number of magnitudes of the same kind, the *first* is said to have to the *last* the **ratio compounded** of the ratios of the *first* to the *second*, of the *second* to the *third*, and so on up to the ratio of the *last but one* to the *last* magnitude.

Thus if  $a, b, c, d, e$  are magnitudes of the same kind, then  $a : e$  is the ratio compounded of the ratios

$$a : b, b : c, c : d, d : e.$$

**NOTE.** *Algebra* defines the **ratio compounded** of given ratios as that formed by *multiplying together* the fractions which represent the given ratios. In the above illustration it will be seen that on multiplying together the ratios  $\frac{a}{b}, \frac{b}{c}, \frac{c}{d}, \frac{d}{e}$  we obtain the ratio  $\frac{a}{e}$ .

*Definition 13.* When three magnitudes are proportionals, the *first* is said to have to the *third* the **duplicate ratio** of that which it has to the *second*.

Thus if  $a : b :: b : c$ ,  
then  $a : c$  is said to be the duplicate of the ratio  $a : b$ .

**NOTE.** In *Algebra* the duplicate of the ratio  $a : b$  is defined as the ratio of  $a^2$  to  $b^2$ .

It is easy to show that the two definitions are identical.

For if  $a : b :: b : c$ ,  
then  $\frac{a}{b} = \frac{b}{c}$ .  
Now  $\frac{a}{c} = \frac{a}{b} \cdot \frac{b}{c} = \frac{a}{b} \cdot \frac{a}{b} = \frac{a^2}{b^2}$ ;  
that is,  $a : c :: a^2 : b^2$ .

6. The following theorems from Book V. are here proved algebraically. Reference is made to them in Book VI. under certain technical names.

**THEOREM 1. By Equal Ratios.** *Ratios which are equal to the same ratio are equal to one another.*

That is, if  $a : b = x : y$ , and  $c : d = x : y$ ;  
then shall  $a : b = c : d$ .

For, by hypothesis,  $\frac{a}{b} = \frac{x}{y}$ , and  $\frac{c}{d} = \frac{x}{y}$ ;

hence  $\frac{a}{b} = \frac{c}{d}$

or  $a : b = c : d$ .

**THEOREM 3. Invertendo, or Inversely.** *If four magnitudes are proportionals, they are also proportionals taken inversely.*

That is, if  $a : b = x : y$ ,  
then shall  $b : a = y : x$ .

Since, by hypothesis,  $\frac{a}{b} = \frac{x}{y}$ , it follows that  $\frac{b}{a} = \frac{y}{x}$ ;

or  $b : a = y : x$ .

**THEOREM 11. Alternando, or Alternately.** *If four magnitudes of the same kind are proportionals, they are also proportionals when taken alternately.*

That is, if  $a : b = x : y$ ,  
then shall  $a : x = b : y$ .

For, by hypothesis,  $\frac{a}{b} = \frac{x}{y}$ .

Multiplying both sides by  $\frac{b}{x}$

we have  $\frac{a}{b} \cdot \frac{b}{x} = \frac{x}{y} \cdot \frac{b}{x}$ ;

that is,  $\frac{a}{x} = \frac{b}{y}$ ,

or  $a : x = b : y$ .

**NOTE.** In this theorem the *hypothesis* requires that  $a$  and  $b$  shall be of the same kind, also that  $x$  and  $y$  shall be of the same kind; while the *conclusion* requires that  $a$  and  $x$  shall be of the same kind, and also  $b$  and  $y$  of the same kind.

**THEOREM 12. Addendo.** *In a series of equal ratios (the magnitudes being all of the same kind), as any antecedent is to its consequent so is the sum of the antecedents to the sum of the consequents.*

That is, if  $a : x = b : y = c : z = \dots$ ;  
then shall  $a : x = a + b + c + \dots : x + y + z + \dots$ .

Let each of the equal ratios  $\frac{a}{x}, \frac{b}{y}, \frac{c}{z} \dots$  be equal to  $k$ .

Then  $a = kx, b = ky, c = kz, \dots$ ;  
 $\therefore$ , by addition,

$$a + b + c + \dots = k(x + y + z + \dots);$$

$$\therefore \frac{a + b + c + \dots}{x + y + z + \dots} = k = \frac{a}{x}$$

or  $a : x = a + b + c + \dots : x + y + z + \dots$ .

**THEOREM 13. Componendo.** *If four magnitudes are proportionals, the sum of the first and second is to the second as the sum of the third and fourth is to the fourth.*

That is, if  $a : b = x : y$ ;  
then shall  $a + b : b = x + y : y$ .

For, by hypothesis,  $\frac{a}{b} = \frac{x}{y}$ ;

$$\therefore \frac{a}{b} + 1 = \frac{x}{y} + 1, \text{ or } \frac{a+b}{b} = \frac{x+y}{y};$$

that is,  $a + b : b = x + y : y$ .

**Dividendo.** Similarly it may be shewn that  $a - b : b = x - y : y$ .

**THEOREM 14. Ex Aequali.** *If there are three magnitudes  $a, b, c$  of one set, and three magnitudes  $x, y, z$  of another set; and if these are so related that*

and  $a : b = x : y,$   
 $b : c = y : z,$   
then shall  $a : c = x : z$ .

For, by hypothesis,  $\frac{a}{b} = \frac{x}{y}$ , and  $\frac{b}{c} = \frac{y}{z}$ ;

$\therefore$ , by multiplication,  $\frac{a}{b} \cdot \frac{b}{c} = \frac{x}{y} \cdot \frac{y}{z}$ ;

that is,  $\frac{a}{c} = \frac{x}{z}$ ;

or  $a : c = x : z$ .

**THEOREM 15.** *If two proportions have the same consequents,*

that is, if  $a : b = x : y,$   
and  $c : b = z : y,$   
then shall  $a + c : b = x + z : y$ .

For, by hypothesis,  $\frac{a}{b} = \frac{x}{y}$ , and  $\frac{c}{b} = \frac{z}{y}$ ;

$\therefore$ , by addition,  $\frac{a+c}{b} = \frac{x+z}{y}$ ;

or  $a + c : b = x + z : y$ .

## BOOK VI.


### DEFINITIONS.

1. Two rectilinear figures are said to be **equiangular** to one another when the angles of the first, taken in order, are equal respectively to those of the second, taken in order.

2. Rectilinear figures are said to be **similar** when they are equiangular to one another, and also have the sides about the equal angles taken in order proportionals.

Thus the two quadrilaterals ABCD, EFGH are similar if the angles at A, B, C, D are respectively equal to those at E, F, G, H, and if the following proportions hold :

$$\begin{aligned} AB : BC &:: EF : FG, \\ BC : CD &:: FG : GH, \\ CD : DA &:: GH : HE, \\ DA : AB &:: HE : EF. \end{aligned}$$



In these proportions, sides which are *both antecedents* or *both consequents* of the ratios are said to be *homologous* or *corresponding*.  
[Def. 6, p. 320.]

Thus AB and EF are homologous sides ; so are BC and FG.

3. Two similar rectilinear figures are said to be **similarly situated** with respect to two of their sides when these sides are *homologous*.


4. Two figures are said to have their sides about one angle in each **reciprocally proportional** when a side of the *first* figure is to a side of the *second* as the remaining side of the *second* figure is to the remaining side of the *first*.

5. A straight line is said to be divided in **extreme and mean ratio** when the whole is to the greater segment as the greater segment is to the less.



## PROPOSITION 1. THEOREM. [EUCLID'S PROOF.]

*The areas of triangles of the same altitude are to one another as their bases.*



Let  $\triangle ABC$ ,  $\triangle ACD$  be two triangles of the same altitude, namely the perpendicular from  $A$  to  $BD$ .

*Then shall the  $\triangle ABC : \text{the } \triangle ACD :: BC : CD$ .*

Produce  $BD$  both ways ;  
and from  $CB$  produced cut off *any* number of parts  $BG$ ,  $GH$ , each equal to  $BC$  ;  
and from  $CD$  produced cut off *any* number of parts  $DK$ ,  $KL$ ,  $LM$ , each equal to  $CD$ .

Join  $AH$ ,  $AG$ ,  $AK$ ,  $AL$ ,  $AM$ .

Since the  $\triangle ABC$ ,  $\triangle ABG$ ,  $\triangle AGH$  are of the same altitude, and stand on the equal bases  $CB$ ,  $BG$ ,  $GH$ ,

$\therefore$  the  $\triangle ABC$ ,  $\triangle ABG$ ,  $\triangle AGH$  are equal in area ; I. 38.  
 $\therefore$  the  $\triangle AHC$  is the same multiple of the  $\triangle ABC$  that  $HC$  is of  $BC$ .

Similarly the  $\triangle ACM$  is the same multiple of the  $\triangle ACD$  that  $CM$  is of  $CD$ .

And if  $HC = CM$ ,  
the  $\triangle AHC = \text{the } \triangle ACM$  ; I. 38.

and if  $HC$  is greater than  $CM$ ,  
the  $\triangle AHC$  is greater than the  $\triangle ACM$  ; I. 38, *Cor.*

and if  $HC$  is less than  $CM$ ,  
the  $\triangle AHC$  is less than the  $\triangle ACM$ . I. 38, *Cor.*


Now since there are four magnitudes, namely, the  $\triangle ABC$ ,  $\triangle ACD$ , and the bases  $BC$ ,  $CD$  ; and of the antecedents, any equimultiples have been taken, namely, the  $\triangle AHC$

and the base HC; and of the consequents, any equimultiples have been taken, namely the  $\triangle ACM$  and the base CM; and since it has been shewn that the  $\triangle AHC$  is greater than, equal to, or less than the  $\triangle ACM$ , according as HC is greater than, equal to, or less than CM;

$\therefore$  the four original magnitudes are proportionals; v. *Def.* 5. that is,

the  $\triangle ABC$  : the  $\triangle ACD$  :: the base BC : the base CD.  $\cdot$ Q.E.D.

*COROLLARY.* The areas of parallelograms of the same altitude are to one another as their bases.



Let EC, CF be par<sup>m</sup>s of the same altitude.

Then shall the par<sup>m</sup> EC : the par<sup>m</sup> CF :: BC : CD.

Join BA, AD.

Then the  $\triangle ABC$  : the  $\triangle ACD$  :: BC : CD; *Proved.*  
but the par<sup>m</sup> EC is double of the  $\triangle ABC$ , I. 34.

and the par<sup>m</sup> CF is double of the  $\triangle ACD$ ;  
 $\therefore$  the par<sup>m</sup> EC : the par<sup>m</sup> CF :: BC : CD. v. 8.


**NOTE.**

This proof of Proposition 1 is founded on Euclid's Test of Proportion, and therefore holds good whether the bases BC, CD are commensurable or otherwise.

The numerical treatment given on the following page applies in strict theory only to the former case; but the beginner would do well to accept it, at any rate provisionally, and thus postpone to a later reading the acknowledged difficulty of Euclid's Theory of Proportion.

## PROPOSITION 1. [NUMERICAL ILLUSTRATION.]

*The areas of triangles of equal altitude are to one another as their bases.*



Let  $ABC$ ,  $DEF$  be two triangles between the same par<sup>ls</sup>, and therefore of equal altitude.

Then shall the  $\triangle ABC$  : the  $\triangle DEF$  = the base  $BC$  : the base  $EF$ .

Suppose  $BC$  contains 4 units of length, and  $EF$  3 units; and let  $BL$ ,  $LM$ ,  $MN$ ,  $NC$  each represent *one* unit, as also  $ER$ ,  $RS$ ,  $SF$ .

Then  $BC : EF = 4 : 3$ .

Join  $AL$ ,  $AM$ ,  $AN$ ; also  $DR$ ,  $DS$ .

Then the four  $\triangle^s$   $ABL$ ,  $ALM$ ,  $AMN$ ,  $ANC$  are all equal; for they stand on equal bases, and are of equal altitude.

$\therefore$  the  $\triangle ABC$  is *four* times the  $\triangle ABL$ .

Similarly, the  $\triangle DEF$  is *three* times the  $\triangle DER$ .

But the  $\triangle^s$   $ABL$  and  $DER$  are equal, for they are on equal bases  $BL$ ,  $ER$ , and of equal altitude;

hence the  $\triangle ABC$  : the  $\triangle DEF = 4 : 3$   
 $= BC : EF$ .

This reasoning holds good however many units of length the bases  $BC$ ,  $EF$  contain.

Thus if  $BC = m$  units, and  $EF = n$  units, then, whatever whole numbers  $m$  and  $n$  represent,

the  $\triangle ABC$  : the  $\triangle DEF = m : n$   
 $= BC : EF$ .

The corollary should then be proved as on page 327.

EXERCISES ON PROPOSITION 1.

1. Two triangles of equal altitude stand on bases of 6·3 inches and 5·4 inches respectively ; if the area of the first triangle is  $12\frac{1}{4}$  square inches, find the area of the other. [ $10\frac{1}{2}$  sq. in.]

2. The areas of two triangles of equal altitude have the ratio 24 : 17 ; if the base of the first is 4·2 centimetres, find the base of the second to the nearest millimetre. [3·0 c.m.]

3. Two triangles lying between the same parallels have bases of 16·20 metres and 20·70 metres ; find to the nearest square centimetre the area of the second triangle, if that of the first is 50·1204 sq. metres. [60·0427 sq. m.]

4. Assuming that *the area of a triangle* =  $\frac{1}{2}$  *base* × *altitude*, prove algebraically that

- (i) Triangles of equal altitudes are proportional to their bases ;
- (ii) Triangles on equal bases are proportional to their altitudes.

Also deduce the second of these propositions *geometrically* from the first.

5. Two triangular fields lie on opposite sides of a common base ; and their altitudes with respect to it are 4·20 chains and 3·71 chains. If the first field contains 18 acres, find the acreage of the whole quadrilateral. [33·9 acres.]

DEFINITION.

Two straight lines are cut *proportionally* when the segments of one line are in the same ratio as the corresponding segments of the other. [See definition, page 139.]

Fig. 1.





Fig. 2.




Thus AB and CD are cut proportionally at X and Y, if  
 $AX : XB :: CY : YD.$

And the same definition applies equally whether X and Y divide AB and CD internally as in Fig. 1 or externally as in Fig. 2.

## PROPOSITION 2. THEOREM.

If a straight line is drawn parallel to one side of a triangle, it cuts the other sides, or those sides produced, proportionally.

Conversely, if the sides, or the sides produced, are cut proportionally, the straight line which joins the points of section, is parallel to the remaining side of the triangle.



Let  $XY$  be drawn par<sup>l</sup> to  $BC$ , one of the sides of the  $\triangle ABC$ .

Then shall  $BX : XA :: CY : YA$

Join  $BY, CX$ .

Now the  $\triangle BXY, CXY$  are on the same base  $XY$  and between the same par<sup>l</sup>  $XY, BC$ ;

$\therefore$  the  $\triangle BXY =$  the  $\triangle CXY$ ; I. 37.  
and  $AXY$  is another triangle;

$\therefore$  the  $\triangle BXY : \text{the } \triangle AXY :: \text{the } \triangle CXY : \text{the } \triangle AXY$ . v. 4.

But the  $\triangle BXY : \text{the } \triangle AXY :: BX : XA$ , VI. 1.

and the  $\triangle CXY : \text{the } \triangle AXY :: CY : YA$ ;

$\therefore BX : XA :: CY : YA$ . v. 1.

*Conversely.* Let  $BX : XA :: CY : YA$ , and let  $XY$  be joined.

Then shall  $XY$  be par<sup>l</sup> to  $BC$ .

As before, join  $BY, CX$ .

By hypothesis,  $BX : XA :: CY : YA$ ;

but  $BX : XA :: \text{the } \triangle BXY : \text{the } \triangle AXY$ , VI. 1.

and  $CY : YA :: \text{the } \triangle CXY : \text{the } \triangle AXY$ ;

$\therefore$  the  $\triangle BXY : \text{the } \triangle AXY :: \text{the } \triangle CXY : \text{the } \triangle AXY$ . v. 1.

$\therefore$  the  $\triangle BXY =$  the  $\triangle CXY$ ; v. 6.

and these triangles are on the same base and on the same side of it;

$\therefore XY$  is par<sup>l</sup> to  $BC$ .

I. 39.

Q.E.D.

## EXERCISES.

1. Shew that every quadrilateral is divided by its diagonals into four triangles whose areas are proportionals.

2. *If any two straight lines are cut by three parallel straight lines, they are cut proportionally.*

3. From the point E in the common base of two triangles ACB, ADB, straight lines are drawn parallel to AC, AD, meeting BC, BD at F, G : shew that FG is parallel to CD.

4. In a triangle ABC the straight line DEF meets the sides BC, CA, AB at the points D, E, F respectively, and it makes equal angles with AB and AC : prove that

$$BD : CD :: BF : CE.$$

5. In a triangle ABC, AD is drawn perpendicular to BC, the bisector of the angle at B : shew that a straight line through D parallel to AC will bisect AD.

6. From B and C, the extremities of the base of a triangle ABC, straight lines BE, CF are drawn to the opposite sides so as to intersect on the median from A : shew that EF is parallel to BC.


7. From P, a given point in the side AB of a triangle ABC, draw a straight line to AC produced, so that it will be bisected by BC.

8. Find a point within a triangle such that, if straight lines be drawn from it to the three angular points, the triangle will be divided into three equal triangles.

## PROPOSITION 3. THEOREM.

*If the vertical angle of a triangle be bisected by a straight line which cuts the base, the segments of the base shall have to one another the same ratio as the remaining sides of the triangle.*

*Conversely, if the base be divided so that its segments have to one another the same ratio as the remaining sides of the triangle, the straight line drawn from the vertex to the point of section shall bisect the vertical angle.*



In the  $\triangle ABC$ , let the  $\angle BAC$  be bisected by  $AX$ , which meets the base at  $X$ .

Then shall  $BX : XC :: BA : AC$ .

Through  $C$  draw  $CE$  par<sup>l</sup> to  $XA$ , to meet  $BA$  produced at  $E$ . I. 31.

Then because  $XA$  and  $CE$  are par<sup>l</sup>,  
 $\therefore$  the  $\angle BAX =$  the int. opp.  $\angle AEC$ , I. 29.  
 and the  $\angle XAC =$  the alt.  $\angle ACE$ . I. 29.  
 But the  $\angle BAX =$  the  $\angle XAC$ ; *Hyp.*  
 $\therefore$  the  $\angle AEC =$  the  $\angle ACE$ ;  
 $\therefore AC = AE$ . I. 6.

Again, because  $XA$  is par<sup>l</sup> to  $CE$ , a side of the  $\triangle BCE$ ,  
 $\therefore BX : XC :: BA : AE$ ; VI. 2.  
 that is,  $BX : XC :: BA : AC$ .

*Conversely.* Let  $BX : XC :: BA : AC$  ; and let  $AX$  be joined.

*Then shall the  $\angle BAX =$  the  $\angle XAC$ .*

For, with the same construction as before,  
because  $XA$  is par<sup>l</sup> to  $CE$ , a side of the  $\triangle BCE$ ,

$$\therefore BX : XC :: BA : AE. \quad \text{VI. 2.}$$

But, *by hypothesis*,  $BX : XC :: BA : AC$  ;

$$\therefore BA : AE :: BA : AC ; \quad \text{V. 1.}$$

$$\therefore AE = AC ;$$

$$\therefore \text{the } \angle ACE = \text{the } \angle AEC. \quad \text{I. 5.}$$

But because  $XA$  is par<sup>l</sup> to  $CE$ ,

$$\therefore \text{the } \angle XAC = \text{the alt. } \angle ACE. \quad \text{I. 29.}$$

and the ext.  $\angle BAX =$  the int. opp.  $\angle AEC$  ; I. 29.

$$\therefore \text{the } \angle BAX = \text{the } \angle XAC.$$

Q.E.D.

#### EXERCISES.

1. The side  $BC$  of a triangle  $ABC$  is bisected at  $D$ , and the angles  $ADB, ADC$  are bisected by the straight lines  $DE, DF$ , meeting  $AB, AC$  at  $E, F$  respectively : shew that  $EF$  is parallel to  $BC$ .

2. Apply Proposition 3 to trisect a given finite straight line.

3. If the line bisecting the vertical angle of a triangle is divided into parts which are to one another as the base to the sum of the sides, the point of division is the centre of the inscribed circle.

4.  $ABCD$  is a quadrilateral : shew that if the bisectors of the angles  $A$  and  $C$  meet in the diagonal  $BD$ , the bisectors of the angles  $B$  and  $D$  will meet on  $AC$ .

5. Construct a triangle having given the base, the vertical angle, and the ratio of the remaining sides.

6. Employ Proposition 3 to shew that the bisectors of the angles of a triangle are concurrent.


7.  $AB$  is a diameter of a circle,  $CD$  is a chord at right angles to it, and  $E$  any point in  $CD$  ;  $AE$  and  $BE$  are drawn and produced to cut the circle in  $F$  and  $G$  : shew that the quadrilateral  $CFDG$  has any two of its adjacent sides in the same ratio as the remaining two.



## PROPOSITION A. THEOREM.

If one side of a triangle be produced, and the exterior angle so formed be bisected by a straight line which cuts the base produced, the segments between the point of section and the extremities of the base shall have to one another the same ratio as the remaining sides of the triangle.

Conversely, if the segments of the base produced have to one another the same ratio as the remaining sides of the triangle, the straight line drawn from the vertex to the point of section shall bisect the exterior vertical angle.



In the  $\triangle ABC$  let  $BA$  be produced to  $F$ , and let the exterior  $\angle CAF$  be bisected by  $AX$  which meets the base produced at  $X$ .

Then shall

$$BX : XC :: BA : AC.$$

Through  $C$  draw  $CE$  par<sup>l</sup> to  $XA$ ,  
and let  $CE$  meet  $BA$  at  $E$ .

I. 31.

Then because  $AX$  and  $CE$  are par<sup>l</sup>,  
 $\therefore$  the ext.  $\angle FAX =$  the int. opp.  $\angle AEC$ ,  
and the  $\angle XAC =$  the alt.  $\angle ACE$ .

I. 29.

But the  $\angle FAX =$  the  $\angle XAC$ ;

Hyp.

$\therefore$  the  $\angle AEC =$  the  $\angle ACE$ ;

$\therefore AC = AE$ .

I. 6.

Again, because  $XA$  is par<sup>l</sup> to  $CE$ , a side of the  $\triangle BCE$ ,

Constr.

$$\therefore BX : XC :: BA : AE ;$$

VI. 2.

that is,

$$BX : XC :: BA : AC.$$

*Conversely.* Let  $BX : XC :: BA : AC$ , and let  $AX$  be joined.

*Then shall the  $\angle FAX =$  the  $\angle XAC$ .*

For, with the same construction as before,  
because  $AX$  is par<sup>l</sup> to  $CE$ , a side of the  $\triangle BCE$ ,

$$\therefore BX : XC :: BA : AE. \quad \text{VI. 2.}$$

But, *by hypothesis*,  $BX : XC :: BA : AC$ ;

$$\therefore BA : AE :: BA : AC; \quad \text{V. 1.}$$

$$\therefore AE = AC;$$

$$\therefore \text{the } \angle ACE = \text{the } \angle AEC. \quad \text{I. 5.}$$

But because  $AX$  is par<sup>l</sup> to  $CE$ ,

$$\therefore \text{the } \angle XAC = \text{the alt. } \angle AEC,$$

and the ext.  $\angle FAX =$  the int. opp.  $\angle AEC$ ; I. 29.

$$\therefore \text{the } \angle FAX = \text{the } \angle XAC. \quad \text{Q.E.D.}$$

Propositions 3 and A may be both included in one enunciation as follows :

*If the interior or exterior vertical angle of a triangle be bisected by a straight line which also cuts the base, the base shall be divided internally or externally into segments which have the same ratio as the other sides of the triangle.*

*Conversely, if the base be divided internally or externally into segments which have the same ratio as the other sides of the triangle, the straight line drawn from the point of division to the vertex will bisect the interior or exterior vertical angle.*

#### EXERCISES.


1. In the circumference of a circle of which  $AB$  is a diameter, a point  $P$  is taken ; straight lines  $PC, PD$  are drawn equally inclined to  $AP$  and on opposite sides of it, meeting  $AB$  in  $C$  and  $D$  ;  
shew that  $AC : CB :: AD : DB$ .

2. From a point  $A$  straight lines are drawn making the angles  $BAC, CAD, DAE$ , each equal to half a right angle, and they are cut by a straight line  $BCDE$ , which makes  $BAE$  an isosceles triangle : shew that  $BC$  or  $DE$  is a mean proportional between  $BE$  and  $CD$ .

3. By means of Propositions 3 and A, prove that the straight lines bisecting one angle of a triangle internally, and the other two externally, are concurrent.

## PROPOSITION 4. THEOREM.

If two triangles be equiangular to one another, the sides about the equal angles shall be proportionals, those sides which are opposite to equal angles being homologous.



Let the  $\triangle ABC$  be equiangular to the  $\triangle DCE$ ,  
 having the  $\angle ABC$  equal to the  $\angle DCE$ ,  
 the  $\angle BCA$  equal to the  $\angle CED$ ,  
 and consequently the  $\angle CAB$  equal to the  $\angle EDC$ . I. 32.  
 Then shall the sides about these equal angles be proportionals,  
 namely

$$\begin{aligned} AB : BC &:: DC : CE, \\ BC : CA &:: CE : ED, \\ \text{and } AB : AC &:: DC : DE. \end{aligned}$$

Let the  $\triangle DCE$  be placed so that its side  $CE$  may be contiguous to  $BC$ , and in the same straight line with it.

Then because the  $\angle^s ABC, ACB$  are together less than two rt. angles, I. 17.

and the  $\angle ACB =$  the  $\angle DEC$ ; Hyp.

$\therefore$  the  $\angle^s ABC, DEC$  are together less than two rt. angles;

$\therefore BA$  and  $ED$  will meet if produced. Ax. 12.

Let them be produced and meet at  $F$ .

Then because the  $\angle ABC =$  the  $\angle DCE$ , Hyp.

$\therefore BF$  is par<sup>l</sup> to  $CD$ ; I. 28.

and because the  $\angle ACB =$  the  $\angle DEC$ , Hyp.

$\therefore AC$  is par<sup>l</sup> to  $FE$ ; I. 28.

$\therefore FACD$  is a par<sup>m</sup>;

$\therefore AF = CD$ , and  $AC = FD$ . I. 34.

Again, because CD is par<sup>l</sup> to BF, a side of the  $\triangle$  EBF,  
 $\therefore BC : CE :: FD : DE ;$  VI. 2.  
 but  $FD = AC ;$   
 $\therefore BC : CE :: AC : DE ;$   
 and, *alternately*,  $BC : CA :: CE : ED.$  V. 11

Again, because AC is par<sup>l</sup> to FE, a side of the  $\triangle$  FBE,  
 $\therefore BA : AF :: BC : CE ;$  VI. 2.  
 but  $AF = CD ;$   
 $\therefore BA : CD :: BC : CE ;$   
 and, *alternately*,  $AB : BC :: DC : CE.$  V. 11.  
 Also  $BC : CA :: CE : ED ;$  *Proved.*  
 $\therefore$ , *ex æquali*,  $AB : AC :: DC : DE.$  V. 14.

Q.E.D.

[For Alternative Proof see Page 342.]

## EXERCISES.


1. If one of the parallel sides of a trapezium is double the other, shew that the diagonals intersect one another at a point of trisection.
2. In the side AC of a triangle ABC any point D is taken : shew that if AD, DC, AB, BC are bisected in E, F, G, H respectively, then EG is equal to HF.
3. AB and CD are two parallel straight lines ; E is the middle point of CD ; AC and BE meet at F, and AE and BD meet at G : shew that FG is parallel to AB.
4. ABCDE is a regular pentagon, and AD and BE intersect in F : shew that  $AF : AE :: AE : AD.$
5. In the figure of I. 43 shew that EH and GF are parallel, and that FH and GE will meet on CA produced.
6. Chords AB and CD of a circle are produced towards B and D respectively to meet in the point E, and through E, the line EF is drawn parallel to AD to meet CB produced in F. Prove that EF is a mean proportional between FB and FC.

H.S.E.

Y

## PROPOSITION 5. THEOREM.

If the sides of two triangles, taken in order about each of their angles, be proportionals, the triangles shall be equiangular to one another, having those angles equal which are opposite to the homologous sides.



Let the  $\triangle^s$  ABC, DEF have their sides proportionals, so that

$$AB : BC :: DE : EF,$$

$$BC : CA :: EF : FD,$$

and consequently, *ex æquali*,

$$AB : AC :: DE : DF.$$

Then shall the  $\triangle^s$  ABC, DEF be equiangular to one another.

At E in FE make the  $\angle$  FEG equal to the  $\angle$  ABC; I. 23.

and at F in EF make the  $\angle$  EFG equal to the  $\angle$  BCA;

$\therefore$  the remaining  $\angle$  EGF = the remaining  $\angle$  BAC. I. 32.

Then the  $\triangle^s$  ABC, GEF are equiangular to one another;

$$\therefore AB : BC :: GE : EF. \quad \text{VI. 4.}$$

But, *by hypothesis*,  $AB : BC :: DE : EF$ ;

$$\therefore GE : EF :: DE : EF; \quad \text{V. 1.}$$

$$\therefore GE = DE.$$

Similarly  $GF = DF$ .

Then in the  $\triangle^s$  GEF, DEF,

Because  $\left\{ \begin{array}{l} GE = DE, \\ GF = DF, \\ \text{and } EF \text{ is common;} \end{array} \right.$

$\therefore$  the  $\angle$  GEF = the  $\angle$  DEF,  
and the  $\angle$  GFE = the  $\angle$  DFE,  
and the  $\angle$  EGF = the  $\angle$  EDF. I. 8.

But the  $\angle$  GEF = the  $\angle$  ABC ;  
 $\therefore$  the  $\angle$  DEF = the  $\angle$  ABC.

*Constr.*

Similarly, the  $\angle$  EFD = the  $\angle$  BCA ;  
 $\therefore$  the remaining  $\angle$  FDE = the remaining  $\angle$  CAB ; I. 32.  
 that is, the  $\triangle$  DEF is equiangular to the  $\triangle$  ABC.

Q. E. D.

#### NOTE ON SIMILAR FIGURES.

Similar figures may be described as those which have the *same shape*.


For this, *two conditions* are necessary [see VI., *Def.* 2];

- (i) *the figures must have their angles equal each to each ;*
- (ii) *their sides about the equal angles taken in order must be proportional.*

In the case of *triangles* we have learned that these conditions are not independent, for each follows from the other : thus


- (i) if the triangles are *equiangular*, Proposition 4 proves the *proportionality of their sides ;*
- (ii) if the triangles have their *sides proportional*, Proposition 5 proves their *equiangularity*.

This, however, is not necessarily the case with rectilinear figures of more than three sides. For example, the first diagram in the margin shews two figures which are equiangular to one another, but which clearly have not their sides proportional ; while the figures in the second diagram have their sides proportional, but are not equiangular to one another.



## PROPOSITION 6. THEOREM.

*If two triangles have one angle of the one equal to one angle of the other, and the sides about the equal angles proportionals, the triangles shall be similar.*



In the  $\triangle^s$  BAC, EDF, let the  $\angle$  BAC = the  $\angle$  EDF,  
and let  $BA : AC :: ED : DF$ .

*Then shall the  $\triangle^s$  BAC, EDF be similar.*

At D in FD make the  $\angle$  FDG equal to the  $\angle$  CAB : I. 23.  
at F in DF make the  $\angle$  DFG equal to the  $\angle$  ACB ;  
 $\therefore$  the remaining  $\angle$  DGF = the remaining  $\angle$  ABC. I. 32.

Then the  $\triangle^s$  BAC, GDF are equiangular to one another ;

$\therefore BA : AC :: GD : DF$ . VI. 4.

But, *by hypothesis*,  $BA : AC :: ED : DF$  ;

$\therefore GD : DF :: ED : DF$ ,

$\therefore GD = ED$ .

Then in the  $\triangle^s$  GDF, EDF,

$GD = ED$ ,

Because { and DF is common ; *Constr.*  
{ and the  $\angle$  GDF = the  $\angle$  EDF ;

$\therefore$  the  $\triangle^s$  GDF, EDF are equal in all respects ; I. 4.

so that the  $\triangle$  EDF is equiangular to the  $\triangle$  GDF ;

but the  $\triangle$  GDF is equiangular to the  $\triangle$  BAC ; *Constr.*

$\therefore$  the  $\triangle$  EDF is equiangular to the  $\triangle$  BAC ;

$\therefore$  their sides about the equal angles are proportionals ; VI. 4.  
that is, the  $\triangle^s$  BAC, EDF are similar.

Q.E.D.

## EXERCISES.

## ON PROPOSITIONS 1 TO 6.

1. Shew that the diagonals of a trapezium cut one another in the same ratio.

2. If three straight lines drawn from a point cut two parallel straight lines in A, B, C and P, Q, R respectively, prove that

$$AB : BC :: PQ : QR.$$

3. From a point O, a tangent OP is drawn to a given circle, and a secant OQR is drawn cutting it in Q and R; shew that

$$OQ : OP :: OP : OR.$$

4. *If two triangles are on equal bases and between the same parallels, any straight line parallel to their bases will cut off equal areas from the two triangles.*

5. *If two straight lines PQ, XY intersect in a point O, so that PO : OX :: YO : OQ, prove that P, X, Q, Y are concyclic.*

6. On the same base and on the same side of it two equal triangles ACB, ADB are described; AC and BD intersect in O, and through O lines parallel to DA and CB are drawn meeting the base in E and F. Shew that AE = BF.

7. BD, CD are perpendicular to the sides AB, AC of a triangle ABC, and CE is drawn perpendicular to AD, meeting AB in E: shew that the triangles ABC, ACE are similar.

8. AC and BD are drawn perpendicular to a given straight line CD from two given points A and B; AD and BC intersect in E, and EF is perpendicular to CD: shew that AF and BF make equal angles with CD.

9. ABCD is a parallelogram; P and Q are points in a straight line parallel to AB; PA and QB meet at R, and PD and QC meet at S: shew that RS is parallel to AD.


10. In the sides AB, AC of a triangle ABC two points D, E are taken such that BD is equal to CE; if DE, BC produced meet at F, shew that AB : AC :: EF : DF.

11. Find a point the perpendiculars from which on the sides of a given triangle shall be in a given ratio.



## PROPOSITION 7. THEOREM.

If two triangles have one angle of the one equal to one angle of the other, and the sides about one other angle in each proportional, so that the sides opposite to the equal angles are homologous, then the third angles are either equal or supplementary; and in the former case the triangles are similar.



Let  $ABC$ ,  $DEF$  be two triangles having the  $\angle ABC$  equal to the  $\angle DEF$ , and the sides about the angles at  $A$  and  $D$  proportional, namely

$$BA : AC :: ED : DF.$$

Then shall the  $\angle ACB$ ,  $DFE$  be either equal (as in Figs. 1 and 2) or supplementary (as in Figs. 1 and 3), and in the former case the triangles shall be similar.

If the  $\angle BAC =$  the  $\angle EDF$ , [Figs. 1 and 2.]

then the  $\angle ACB =$  the  $\angle DFE$ ; I. 32.

and the  $\Delta^s$  are equiangular, and therefore similar. VI. 4.

But if the  $\angle BAC$  is not equal to the  $\angle EDF$ , [Figs. 1 and 3.] one of them must be the greater.

Let the  $\angle EDF$  be greater than the  $\angle BAC$ .

At  $D$  in  $ED$  make the  $\angle EDF'$  equal to the  $\angle BAC$ . [Fig. 3.]

Then the  $\Delta^s BAC$ ,  $EDF'$  are equiangular, I. 32.

$\therefore BA : AC :: ED : DF'$ ; VI. 4.

but, by hypothesis,  $BA : AC :: ED : DF$ ;

$\therefore ED : DF :: ED : DF'$ , V. 1.

$\therefore DF = DF'$ ,


$\therefore$  the  $\angle DFF' =$  the  $\angle DF'F$ . I. 5.

But the  $\angle^s DF'F$ ,  $DF'E$  are supplementary, I. 13.

$\therefore$  the  $\angle^s DFF'$ ,  $DF'E$  are supplementary:

that is, the  $\angle^s DFE$ ,  $ACB$  are supplementary. Q.E.D.

COROLLARIES TO PROPOSITION 7.



Three cases of this theorem deserve special attention. It has been proved that if the angles  $ACB$ ,  $DFE$  are not *supplementary*, they are *equal*.

Hence, in addition to the hypothesis of this theorem,


- (i) If the angles  $ACB$ ,  $DFE$ , opposite to the two homologous sides  $AB$ ,  $DE$  are both acute or both obtuse, they cannot be supplementary, and are therefore equal: or if one of them is a right angle, the other must also be a right angle (whether considered as supplementary or equal to it):  
in either case the triangles are similar.
- (ii) If the two given angles at  $B$  and  $E$  are right angles or obtuse angles, it follows that the angles  $ACB$ ,  $DFE$  must be both acute, and therefore equal, by (i):  
so that the triangles are similar.
- (iii) If in each triangle the side opposite the *given* angle is not less than the other given side; that is, if  $AC$  and  $DF$  are not less than  $AB$  and  $DE$  respectively, then the angles  $ACB$ ,  $DFE$  cannot be greater than the angles  $ABC$ ,  $DEF$ , respectively;  
therefore the angles  $ACB$ ,  $DFE$ , are both acute;  
hence, as above, they are equal;  
and the triangles  $ABC$ ,  $DEF$  are similar.

*Obs.* We have given Euclid's demonstrations of Propositions 4, 5, 6; but these propositions also admit of easy proof by the method of superposition.

As an illustration, we will apply this method to Proposition 4.

PROPOSITION 4. [ALTERNATIVE PROOF.]

*If two triangles be equiangular to one another, the sides about the equal angles shall be proportionals, those sides which are opposite to equal angles being homologous.*



Let the  $\triangle ABC$  be equiangular to the  $\triangle DEF$ ,  
 having the  $\angle ABC$  equal to the  $\angle DEF$ ,  
 the  $\angle BCA$  equal to the  $\angle EFD$ ,

and consequently the  $\angle CAB$  equal to the  $\angle FDE$ . I. 32.

*Then shall the sides about these equal angles be proportionals.*

Apply the  $\triangle ABC$  to the  $\triangle DEF$ , so that  $B$  falls on  $E$ , and  $BA$  along  $ED$ :

then  $BC$  will fall along  $EF$ , since the  $\angle ABC = \angle DEF$ . *Hyp.*  
 Let  $G$  and  $H$  be the points in  $ED$  and  $EF$ , on which  $A$  and  $C$  fall;  
 then  $GH$  represents  $AC$  in its new position.

Then because the  $\angle EGH$  (*i.e.* the  $\angle BAC$ ) = the  $\angle EDF$ , *Hyp.*

$\therefore GH$  is par<sup>l</sup> to  $DF$ : vi. 2.

$\therefore DG : GE :: FH : HE$ ;

$\therefore$ , *componendo*,  $DE : GE :: FE : HE$ , v. 13.

$\therefore$ , *alternately*,  $DE : FE :: GE : HE$ , v. 11.

that is,  $DE : EF :: AB : BC$ .

Similarly by applying the  $\triangle ABC$  to the  $\triangle DEF$ , so that the point  $C$  may fall on  $F$ , it may be proved that

$EF : FD :: BC : CA$ .

$\therefore$ , *ex æquali*,  $DE : DF :: AB : AC$ .

Q.E.D.

QUESTIONS FOR REVISION, AND NUMERICAL ILLUSTRATIONS.

1. Distinguish between the use of the word *equiangular* in the following cases :

- (i) the figure ABCD is *equiangular* ;
- (ii) the figure ABCD is *equiangular* to the figure EFGH.

2. Define the terms *ratio*, *antecedent*, *consequent*. Why must the terms of a ratio be of the *same kind*? When are ratios said to be reciprocal?

3. When are four quantities *in proportion*? Quote the algebraical and geometrical tests of proportion; and deduce the latter from the former.

4. What is meant by *homologous terms* in a proportion? In the enunciation of Prop. 4, why is it necessary to add—*those sides which are opposite to equal angles being homologous*?

5. Quote the enunciation of the theorem known as *alternando* or *alternately*; and explain why the terms of a proportion to which this theorem is applied must be all of the same kind.

6. In the Particular Enunciation of Proposition 5 it is given that  
 $AB : BC :: DE : EF,$   
 and  $BC : CA :: EF : FD;$   
 Why do we add "*and consequently,*"  
 $AB : CA :: DE : FD?$

7. Define *similar figures*. In what way do the conditions of similarity in *triangles* differ from those in figures of more than three sides?

8. Two parallelograms whose areas are in the ratio 2·1 : 3·5 lie between the same parallels. If the base of the first is 6·6 inches in length, shew that the base of the second is 11 inches.


9. ABC is a triangle, and XY is drawn parallel to BC, cutting the other sides at X and Y :

- (i) If  $AB = 1$  foot,  $AC = 8$  inches, and  $AX = 7$  inches; shew that  $AY = 4\frac{2}{3}$  inches.
- (ii) If  $AB = 20$  inches,  $AC = 15$  inches, and  $AY = 9$  inches, shew that  $BX = 8$  inches.
- (iii) If X divides AB in the ratio 8 : 3, and if  $AC = 2\cdot2$  inches, shew that AY, YC measure respectively 1·6 and ·6 inches.

10. The vertical angle A of a triangle ABC is bisected by a line which cuts BC at X; if  $BC = 25$  inches in length, and if the sides BA, AC are in the ratio 7 : 3, shew that the segments of the base are 17·5 and 7·5 inches respectively.

## PROPOSITION 8. THEOREM.

*In a right-angled triangle, if a perpendicular is drawn from the right angle to the hypotenuse, the triangles on each side of it are similar to the whole triangle and to one another.*



Let BAC be a triangle right-angled at A, and let AD be drawn perp. to BC.

*Then shall the  $\triangle$  BDA, ADC be similar to the  $\triangle$  BAC and to one another.*

In the  $\triangle$  BDA, BAC,  
 the  $\angle$  BDA = the  $\angle$  BAC, being rt. angles,  
 and the angle at B is common to both ;  
 $\therefore$  the remaining  $\angle$  BAD = the remaining  $\angle$  BCA, I. 32.  
 that is, the  $\triangle$  BDA is equiangular to the  $\triangle$  BAC ;  
 $\therefore$  their sides about the equal angles are proportionals ; VI. 4.  
 $\therefore$  the  $\triangle$  BDA, BAC are similar.

In the same way it may be proved that the  $\triangle$  ADC, BAC are similar.

Hence the  $\triangle$  BDA, ADC, having their angles severally equal to those of the  $\triangle$  BAC, are equiangular to one another ;  
 $\therefore$  they are similar. VI. 4.

Q.E.D.

COROLLARY. Because the  $\triangle$  BDA, ADC are similar,  
 $\therefore$  BD : DA :: DA : DC ;  
 and because the  $\triangle$  CBA, ABD are similar,  
 $\therefore$  CB : BA :: BA : BD ;  
 and because the  $\triangle$  BCA, ACD are similar,  
 $\therefore$  BC : CA :: CA : CD.

## EXERCISES.


1. In the figure of Prop. 8 prove that the hypotenuse is to one side as the second side is to the perpendicular.

2. Shew that the radius of a circle is a mean proportional between the segments of any tangent between its point of contact and a pair of parallel tangents.

DEFINITION. One magnitude is said to be a **submultiple** of another, when the first is contained an *exact* number of times in the second. [Book v. Def. 2.]

PROPOSITION 9. PROBLEM.

*From a given straight line to cut off any required submultiple.*



Let AB be the given straight line.

*It is required to cut off a certain submultiple from AB.*

From A draw a straight line AG of indefinite length, making any angle with AB.

In AG take *any* point D; and, by cutting off successive parts each equal to AD, make AE to contain AD as many times as AB contains the required submultiple.

Join EB.

Through D draw DF  $\text{par}^l$  to EB, meeting AB in F.

*Then shall AF be the required submultiple.*

Because DF is  $\text{par}^l$  to EB, a side of the  $\triangle$  AEB,

$$\therefore BF : FA :: ED : DA; \quad \text{VI. 2.}$$

$$\therefore, \text{ componendo, } BA : AF :: EA : AD. \quad \text{V. 13.}$$

But AE contains AD the required number of times; *Constr.*


$\therefore$  AB contains AF the required number of times;  
that is, AF is the required submultiple. Q.E.F.

EXERCISES.

1. Divide a straight line into five equal parts.
2. Give a geometrical construction for cutting off two-sevenths of a given straight line.

## PROPOSITION 10. PROBLEM.

To divide a straight line similarly to a given divided straight line.



Let AB be the given straight line to be divided, and AC the given straight line divided at the points D and E.

It is required to divide AB similarly to AC.

Let AB, AC be placed so as to form any angle.

Join CB.

Through D draw DF  $\text{par}^1$  to CB, i. 31.  
and through E draw EG  $\text{par}^1$  to CB.

Then AB shall be divided at F and G similarly to AC.

Through D draw DHK  $\text{par}^1$  to AB.

Now by construction each of the figs. FH, HB is a  $\text{par}^m$ ;  
 $\therefore$  DH = FG, and HK = GB. i. 34.

Now since HE is  $\text{par}^1$  to KC, a side of the  $\triangle$  DKC,  
 $\therefore$  KH : HD :: CE : ED. vi. 2.

But KH = BG, and HD = GF ;  
 $\therefore$  BG : GF :: CE : ED. v. 1.

Again, because FD is  $\text{par}^1$  to GE, a side of the  $\triangle$  AGE,  
 $\therefore$  GF : FA :: ED : DA ; vi. 2.


$\therefore$ , *ex æquali*, BG : FA :: CE : DA ; v. 14.  
 $\therefore$  AB is divided similarly to AC. Q.E.F.

## EXERCISE.

Divide a straight line internally and externally in a given ratio.  
Is this always possible?

PROPOSITION 11. PROBLEM.

To find a third proportional to two given straight lines.



Let A, B be two given straight lines.

It is required to find a third proportional to A and B.

Take two st. lines DL, DK of indefinite length, containing any angle.

From DL cut off DG equal to A, and GE equal to B;  
and from DK cut off DH also equal to B. I. 3.  
Join GH.

Through E draw EF par<sup>l</sup> to GH, meeting DK in F. I. 31.

Then shall HF be a third proportional to A and B.

Because GH is par<sup>l</sup> to EF, a side of the  $\triangle DEF$ ;

$\therefore DG : GE :: DH : HF.$  VI. 2.

But  $DG = A$ ; and  $GE, DH$  each = B; Constr.

$\therefore A : B :: B : HF;$

that is, HF is a third proportional to A and B.

Q.E.F.

EXERCISES.

1. AB is a diameter of a circle, and through A any straight line is drawn to cut the circumference in C and the tangent at B in D; shew that AC is a third proportional to AD and AB.


2. ABC is an isosceles triangle having each of the angles at the base double of the vertical angle BAC; the bisector of the angle BCA meets AB at D. Shew that AB, BC, BD are three proportionals.

3. Two circles intersect at A and B; and at A tangents are drawn, one to each circle, to meet the circumferences at C and D; shew that if CB, BD are joined, BD is a third proportional to CB, BA.



## PROPOSITION 12. PROBLEM.

To find a fourth proportional to three given straight lines.



Let A, B, C be the three given straight lines.

It is required to find a fourth proportional to A, B, C.

Take two straight lines DL, DK of indefinite length, containing any angle.

From DL cut off DG equal to A, and GE equal to B;  
and from DK cut off DH equal to C. I. 3.

Join GH.

Through E draw EF par<sup>l</sup> to GH. I. 31.

Then shall HF be a fourth proportional to A, B, C.

Because GH is par<sup>l</sup> to EF, a side of the  $\triangle DEF$ ;

$\therefore DG : GE :: DH : HF.$  VI. 2.

But  $DG = A$ ,  $GE = B$ , and  $DH = C$ ; Constr.

$\therefore A : B :: C : HF$ ;

that is, HF is a fourth proportional to A, B, C.

Q.E.F.

## EXERCISES.


1. If from D, one of the angular points of a parallelogram ABCD, a straight line is drawn meeting AB at E and CB at F; shew that CF is a fourth proportional to EA, AD, and AB.

2. In a triangle ABC the bisector of the vertical angle BAC meets the base at D and the circumference of the circumscribed circle at E: shew that BA, AD, EA, AC are four proportionals.

3. From a point P tangents PQ, PR are drawn to a circle whose centre is C, and QT is drawn perpendicular to RC produced: shew that QT is a fourth proportional to PR, RC, and RT.

PROPOSITION 13. PROBLEM.

*To find a mean proportional between two given straight lines.*



Let AB, BC be the two given straight lines.

*It is required to find a mean proportional between AB and BC.*

Place AB, BC in a straight line, and on AC describe the semicircle ADC.

From B draw BD at rt. angles to AC. I. 11.

*Then shall BD be a mean proportional between AB and BC.*

Join AD, DC.

Now the  $\angle ADC$ , being in a semicircle, is a rt. angle; III. 31. and because in the right-angled  $\triangle ADC$ , DB is drawn from the rt. angle perp. to the hypotenuse,

$\therefore$  the  $\triangle$ 's ABD, BDC are similar; VI. 8.

$\therefore AB : BD :: BD : BC$ ;

that is, BD is a mean proportional between AB and BC.

Q.E.F.

EXERCISES.

1. If from one angle A of a parallelogram a straight line is drawn cutting the diagonal in E and the sides in P, Q, shew that AE is a mean proportional between PE and EQ.

2. A, B, C are three points in order in a straight line: find a point P in the straight line so that PB may be a mean proportional between PA and PC.

3. The diameter AB of a semicircle is divided at any point C, and CD is drawn at right angles to AB meeting the circumference in D; DO is drawn to O the centre, and CE is perpendicular to OD: shew that DE is a third proportional to AO and DC.

4. AC is the diameter of a semicircle on which a point B is taken so that BC is equal to the radius: shew that AB is a mean proportional between BC and the sum of BC, CA.

5. A is any point in a semicircle on BC as diameter; from D any point in BC a perpendicular is drawn meeting AB, AC, and the circumference in E, G, F respectively; shew that DG is a third proportional to DE and DF.

6. Two circles have external contact, and a common tangent touches them at A and B: prove that AB is a mean proportional between the diameters of the circles. [See Ex. 21, p. 237.]

7. If a straight line is divided at two given points, determine a third point such that its distances from the extremities may be proportional to its distances from the given points.

8. AB is a straight line divided at C and D so that AB, AC, AD are in continued proportion; from A a line AE is drawn in any direction and equal to AC; shew that BC and CD subtend equal angles at E.

9. In a given triangle draw a straight line parallel to one of the sides, so that it may be a mean proportional between the segments of the base.

10. On the radius OA of a quadrant OAB, a semicircle ODA is described, and at A a tangent AE is drawn; from O any line ODFE is drawn meeting the circumferences in D and F and the tangent in E: if DG is drawn perpendicular to OA, shew that OE, OF, OD, and OG are in continued proportion.


11. From any point A, on the circumference of the circle ABE, as centre, and with any radius, a circle BDC is described cutting the former circle in B and C; from A any line AFE is drawn meeting the chord BC in F, and the circumferences BDC, ABE in D, E respectively: shew that AD is a mean proportional between AF and AE.

DEFINITION. Two figures are said to have their sides about one angle in each **reciprocally proportional**, when a side of the *first* figure is to a side of the *second* as the remaining side of the *second* figure is to the remaining side of the *first*. [Book VI. Def. 4.]

PROPOSITION 14. THEOREM.

*Parallelograms which are equal in area, and which have one angle of the one equal to one angle of the other, have their sides about the equal angles reciprocally proportional.*

*Conversely, parallelograms which have one angle of the one equal to one angle of the other, and the sides about these angles reciprocally proportional, are equal in area.*



Let the par<sup>m</sup> AB, BC be of equal area, and have the  $\angle$  DBF equal to the  $\angle$  GBE.

Then shall the sides about the  $\angle$  DBF, GBE be reciprocally proportional, namely,

$$DB : BE :: GB : BF.$$

Place the par<sup>m</sup> so that DB, BE may be in the same straight line ;

$\therefore$  FB, BG are also in one straight line. I. 14.

Complete the par<sup>m</sup> FE.

Then because the par<sup>m</sup> AB = the par<sup>m</sup> BC, Hyp.  
and FE is another par<sup>m</sup>,

$\therefore$  the par<sup>m</sup> AB : the par<sup>m</sup> FE :: the par<sup>m</sup> BC : the par<sup>m</sup> FE ;  
but the par<sup>m</sup> AB : the par<sup>m</sup> FE :: DB : BE, VI. 1. Cor.

and the par<sup>m</sup> BC : the par<sup>m</sup> FE :: GB : BF ;  
 $\therefore$  DB : BE :: GB : BF. V. 1.

*Conversely.* Let the  $\angle$  DBF be equal to the  $\angle$  GBE,  
and let DB : BE :: GB : BF.

Then shall the par<sup>m</sup> AB be equal in area to the par<sup>m</sup> BC.

For, with the same construction as before,  
by hypothesis, DB : BE :: GB : BF ;  
but DB : BE :: the par<sup>m</sup> AB : the par<sup>m</sup> FE, VI. 1.  
and GB : BF :: the par<sup>m</sup> BC : the par<sup>m</sup> FE,  
 $\therefore$  the par<sup>m</sup> AB : the par<sup>m</sup> FE :: the par<sup>m</sup> BC : the par<sup>m</sup> FE ; V. 1.  
 $\therefore$  the par<sup>m</sup> AB = the par<sup>m</sup> BC. Q.E.D.


H.S.E.

Z

## PROPOSITION 15. THEOREM.

*Triangles which are equal in area, and which have one angle of the one equal to one angle of the other, have their sides about the equal angles reciprocally proportional.*

*Conversely, triangles which have one angle of the one equal to one angle of the other, and the sides about these angles reciprocally proportional, are equal in area.*



Let the  $\triangle CAB$ ,  $EAD$  be of equal area, and have the  $\angle CAB$  equal to the  $\angle EAD$ .

Then shall the sides about the  $\angle CAB$ ,  $EAD$  be reciprocally proportional, namely,

$$CA : AD :: EA : AB.$$

Place the  $\triangle$  so that  $CA$  and  $AD$  may be in the same st. line;  
 $\therefore BA$ ,  $AE$  are also in one st. line. I. 14.  
 Join  $BD$ .

Then because the  $\triangle CAB =$  the  $\triangle EAD$ , *Hyp.*  
 and  $ABD$  is another triangle,

$\therefore$  the  $\triangle CAB : \text{the } \triangle ABD :: \text{the } \triangle EAD : \text{the } \triangle ABD$  ;  
 but the  $\triangle CAB : \text{the } \triangle ABD :: CA : AD$ , VI. 1.  
 and the  $\triangle EAD : \text{the } \triangle ABD :: EA : AB$  ;  
 $\therefore CA : AD :: EA : AB$ . v. 1.

*Conversely.* Let the  $\angle CAB$  be equal to the  $\angle EAD$ ,  
 and let  $CA : AD :: EA : AB$ .

Then shall the  $\triangle CAB =$  the  $\triangle EAD$ .

For, with the same construction as before,  
 by hypothesis,  $CA : AD :: EA : AB$  ;

but  $CA : AD :: \text{the } \triangle CAB : \text{the } \triangle ABD$ , VI. 1.  
 and  $EA : AB :: \text{the } \triangle EAD : \text{the } \triangle ABD$  ;  
 $\therefore$  the  $\triangle CAB : \text{the } \triangle ABD :: \text{the } \triangle EAD : \text{the } \triangle ABD$  ; v. 1.  
 $\therefore$  the  $\triangle CAB =$  the  $\triangle EAD$ . Q.E.D.

## EXERCISES.

## ON PROPOSITIONS 14 AND 15.

1. *Parallelograms which are equal in area and which have their sides reciprocally proportional, have their angles respectively equal.*

2. *Triangles which are equal in area, and which have the sides about a pair of angles reciprocally proportional, have those angles equal or supplementary.*

3. AC, BD are the diagonals of a trapezium which intersect in O; if the side AB is parallel to CD, use Prop. 15 to prove that the triangle AOD is equal to the triangle BOC.

4. From the extremities A, B of the hypotenuse of a right-angled triangle ABC lines AE, BD are drawn perpendicular to AB, and meeting BC and AC produced in E and D respectively: employ Prop. 15 to shew that the triangles ABC, ECD are equal in area.

5. On AB, AC, two sides of any triangle, squares are described externally to the triangle. If the squares are ABDE, ACFG, shew that the triangles DAG, FAE are equal in area.

6. ABCD is a parallelogram; from A and C any two parallel straight lines are drawn meeting DC and AB in E and F respectively; EG, which is parallel to the diagonal AC, meets AD in G: shew that the triangles DAF, GAB are equal in area.

7. Describe an isosceles triangle equal in area to a given triangle and having its vertical angle equal to one of the angles of the given triangle.


8. Prove that the equilateral triangle described on the hypotenuse of a right-angled triangle is equal to the sum of the equilateral triangles described on the sides containing the right angle.

[Let ABC be the triangle right-angled at C; and let BXC, CYA, AZB be the equilateral triangles. Draw CD perpendicular to AB; and join DZ. Then shew by Prop. 15 that the  $\triangle AYC =$  the  $\triangle DAZ$ ; and similarly that the  $\triangle BXC =$  the  $\triangle BDZ$ .]

## PROPOSITION 16. THEOREM.

If four straight lines are proportional, the rectangle contained by the extremes is equal to the rectangle contained by the means.

Conversely, if the rectangle contained by the extremes is equal to the rectangle contained by the means, the four straight lines are proportional.



Let the st. lines  $AB$ ,  $CD$ ,  $EF$ ,  $GH$  be proportional, so that  
 $AB : CD :: EF : GH$ .

Then shall the rect.  $AB$ ,  $GH =$  the rect.  $CD$ ,  $EF$ .

From  $A$  draw  $AK$  perp. to  $AB$ , and equal to  $GH$ . I. 11, 3.

From  $C$  draw  $CL$  perp. to  $CD$ , and equal to  $EF$ .

Complete the par<sup>ms</sup>  $KB$ ,  $LD$ .

Then because  $AB : CD :: EF : GH$ ;  
 and  $EF = CL$ , and  $GH = AK$ ;

$\therefore AB : CD :: CL : AK$ ;

that is, the par<sup>ms</sup>  $KB$ ,  $LD$  have their sides about the equal angles at  $A$  and  $C$  reciprocally proportional;

$\therefore KB = LD$ .

*Hyp.*  
*Constr.*

VI. 14.

But  $KB$  is the rect.  $AB$ ,  $GH$ , for  $AK = GH$ ,  
 and  $LD$  is the rect.  $CD$ ,  $EF$ , for  $CL = EF$ ;

*Constr.*

$\therefore$  the rect.  $AB$ ,  $GH =$  the rect.  $CD$ ,  $EF$ .

*Conversely.* Let the rect. AB, GH = the rect. CD, EF.

Then shall  $AB : CD :: EF : GH$ .

For, with the same construction as before,

because the rect. AB, GH = the rect. CD, EF; *Hyp.*

and the rect. AB, GH = KB, for GH = AK, *Constr.*

and the rect. CD, EF = LD, for EF = CL;

$\therefore KB = LD$ ;

that is, the par<sup>ms</sup> KB, LD, which have the angle at A equal to the angle at C, are equal in area;

$\therefore$  the sides about the equal angles are reciprocally proportional;

that is,  $AB : CD :: CL : AK$ ;

$\therefore AB : CD :: EF : GH$ .

Q.E.D.

#### QUESTIONS FOR REVISION.

1. State and prove the *algebraical* theorem corresponding to Proposition 16.

2. Define the terms: *multiple, submultiple, fourth proportional, third proportional, mean proportional.*

3. ABC is a triangle right-angled at A, and AD is drawn perpendicular to BC: if AB, AC measure respectively 12 and 5 inches, shew that the segments of the hypotenuse are  $11\frac{1}{3}$  and  $1\frac{2}{3}$  inches.

4. Find in inches the length of the mean proportional between 1 inch and 3 inches. Hence give a geometrical construction for drawing a line  $\sqrt{3}$  inches in length: and extend the method to finding a line  $\sqrt{n}$  inches long.

5. A straight line AB, 21 inches in length, is divided at F and G into parts of 5, 7, 9 inches respectively. If a second line AC, 35 inches long, is similarly divided by the method of Proposition 10, shew that the lengths of the parts are  $8\frac{1}{3}$ ,  $11\frac{2}{3}$  and 15 inches respectively.


6. When are figures said to have their sides about one angle in each *reciprocally proportional*? Two equal parallelograms ABCD, EFGH have their angles at B and F equal: if AB = 2 inches, BC = 10 inches, and EF = 5 inches; find the length of FG.



## PROPOSITION 17. THEOREM.

If three straight lines are proportional the rectangle contained by the extremes is equal to the square on the mean.

Conversely, if the rectangle contained by the extremes is equal to the square on the mean, the three straight lines are proportional.



Let the three st. lines A, B, C be proportional, so that

$$A : B :: B : C.$$

Then shall the rect. A, C be equal to the sq. on B.

Take D equal to B.

Then because  $A : B :: B : C$ , and  $D = B$ ;

$$\therefore A : B :: D : C;$$

$\therefore$  the rect. A, C = the rect. B, D; VI. 16.

but the rect. B, D = the sq. on B, for  $D = B$ ;

$\therefore$  the rect. A, C = the sq. on B.

*Conversely.* Let the rect. A, C = the sq. on B.

Then shall  $A : B :: B : C$ .

For, with the same construction as before,

because the rect. A, C = the sq. on B, *Hyp.*

and the sq. on B = the rect. B, D, for  $D = B$ ;

$\therefore$  the rect. A, C = the rect. B, D;

$$\therefore A : B :: D : C, \quad \text{VI. 16.}$$

that is,  $A : B :: B : C$ . Q.E.D.

## QUESTIONS FOR REVISION.

1. State and prove the *algebraical* theorem corresponding to Proposition 17.

2. Two adjacent sides of a rectangle measure 12.1 and .9 inches in length; shew that the side of an equal square is 3.3 inches.

3. ABC is an isosceles triangle, the equal sides each measuring 12 inches. DAE is a triangle of equal area, having the angle DAE equal to the angle CAB. If AD = 36 inches, find the length of AE.

## EXERCISES.

## ON PROPOSITIONS 16 AND 17.

1. Apply Proposition 16 to prove that if two chords of a circle intersect, the rectangle contained by the segments of the one is equal to the rectangle contained by the segments of the other.

2. Prove that the rectangle contained by the sides of a right-angled triangle is equal to the rectangle contained by the hypotenuse and the perpendicular drawn to it from the right angle.

3. On a given straight line construct a rectangle equal to a given rectangle.

4. ABCD is a parallelogram ; from B any straight line is drawn cutting the diagonal AC at F, the side DC at G, and the side AD produced at E : shew that the rectangle EF, FG is equal to the square on BF.

5. On a given straight line as base describe an isosceles triangle equal to a given triangle.

6. AB is a diameter of a circle, and any line ACD cuts the circle in C and the tangent at B in D ; shew by Prop. 17 that the rectangle AC, AD is constant.

7. The exterior angle at A of a triangle ABC is bisected by a straight line which meets the base in D and the circumscribed circle in E : shew that the rectangle BA, AC is equal to the rectangle EA, AD.

8. If two chords AB, AC drawn from any point A in the circumference of the circle ABC are produced to meet the tangent at the other extremity of the diameter through A in D and E, shew that the triangle AED is similar to the triangle ABC.

9. At the extremities of a diameter of a circle tangents are drawn ; these meet the tangent at a point P in Q and R : shew that the rectangle QP, PR is constant for all positions of P.

10. A is the vertex of an isosceles triangle ABC inscribed in a circle, and ADE is a straight line which cuts the base in D and the circle in E ; shew that the rectangle EA, AD is equal to the square on AB.


11. Two circles touch one another externally at A ; a straight line touches the circles at B and C, and is produced to meet the straight line joining the centres at S : shew that the rectangle SB, SC is equal to the square on SA.

12. Divide a triangle into two equal parts by a straight line drawn at right angles to one of the sides.

DEFINITION. Two similar rectilinear figures are said to be **similarly situated** with respect to two of their sides when these sides are *homologous*. [Book VI. Def. 3.]

PROPOSITION 18. PROBLEM.

*On a given straight line to describe a rectilinear figure similar and similarly situated to a given rectilinear figure.*



Let AB be the given st. line, and CDEF the given rectilinear figure.

*It is required to describe on the st. line AB a rectilinear figure similar and similarly situated to CDEF.*

First suppose CDEF to be a quadrilateral.

Join DF.

At A in BA make the  $\angle BAG$  equal to the  $\angle DCF$ , I. 23.  
and at B in AB make the  $\angle ABG$  equal to the  $\angle CDF$ ;

$\therefore$  the remaining  $\angle AGB =$  the remaining  $\angle CFD$ ; I. 32.  
and the  $\triangle AGB$  is equiangular to the  $\triangle CFD$ .

Again at B in GB make the  $\angle GBH$  equal to the  $\angle FDE$ ,  
and at G in BG make the  $\angle BGH$  equal to the  $\angle DFE$ ; I. 23.

$\therefore$  the remaining  $\angle BHG =$  the remaining  $\angle DEF$ ; I. 32.  
and the  $\triangle BHG$  is equiangular to the  $\triangle DEF$ .

*Then shall ABHG be the required figure.*

(i) To prove that the fig. ABHG is equiangular to the fig. CDEF.


Because the  $\angle AGB =$  the  $\angle CFD$ , *Proved.*  
and the  $\angle BGH =$  the  $\angle DFE$ ; *Constr.*

$\therefore$  the whole  $\angle AGH =$  the whole  $\angle CFE$ .

Similarly the  $\angle ABH =$  the  $\angle CDE$ ;

and the angles at A and H are respectively equal to the angles at C and E; *Constr. and proof.*

$\therefore$  the fig. ABHG is equiangular to the fig. CDEF.



(ii) To prove that the figs. ABHG, CDEF have the sides about their equal angles proportional.

Because the  $\triangle AGB$  is equiangular to the  $\triangle CFD$ ,  
 $\therefore AG : GB :: CF : FD.$  VI. 4.

And because the  $\triangle BGH$  is equiangular to the  $\triangle DFE$ ,

$\therefore BG : GH :: DF : FE ;$   
 $\therefore, \text{ex æquali, } AG : GH :: CF : FE.$  V. 14.

Similarly it may be shewn that

$AB : BH :: CD : DE.$

Also  $BA : AG :: DC : CF,$  VI. 4.  
 and  $GH : HB :: FE : ED.$

$\therefore$  the figs. ABHG, CDEF are equiangular and have their sides about the equal angles proportional ;  
 that is, ABHG is similar to CDEF. VI. Def 2.


In like manner the process of construction may be extended to a figure of five or more sides.

Q.E.F.

DEFINITION. When three magnitudes are proportionals the *first* is said to have to the *third* the **duplicate ratio** of that which it has to the *second*. [Book v. Def. 13.]

## PROPOSITION 19. THEOREM.

*Similar triangles are to one another in the duplicate ratio of their homologous sides.*



Let  $\triangle ABC$ ,  $\triangle DEF$  be similar triangles, having the  $\angle ABC$  equal to the  $\angle DEF$ , and let  $BC$  and  $EF$  be homologous sides. Then shall the  $\triangle ABC$  be to the  $\triangle DEF$  in the duplicate ratio of  $BC$  to  $EF$ .

To  $BC$  and  $EF$  take a *third* proportional  $BG$ ,  
so that  $BC : EF :: EF : BG$ . VI. 11.  
Join  $AG$ .

Then because the  $\triangle ABC$ ,  $\triangle DEF$  are similar, *Hyp.*

$\therefore AB : BC :: DE : EF$  ;

$\therefore$ , *alternately*,  $AB : DE :: BC : EF$  ; V. 11.

but  $BC : EF :: EF : BG$  ; *Constr.*

$\therefore AB : DE :: EF : BG$  ; V. 1.

that is, the sides of the  $\triangle ABG$ ,  $\triangle DEF$  about the equal angles at  $B$  and  $E$  are reciprocally proportional ;

$\therefore$  the  $\triangle ABG =$  the  $\triangle DEF$ . VI. 15.

Again, because  $BC : EF :: EF : BG$ , *Constr.*

$\therefore BC : BG$  in the duplicate ratio of  $BC$  to  $EF$ . V. *Def.* 13.

But the  $\triangle ABC : \text{the } \triangle ABG :: BC : BG$  ; VI. 1.

$\therefore$  the  $\triangle ABC : \text{the } \triangle ABG$  in the duplicate ratio of  $BC$  to  $EF$  ; V. 1.

and the  $\triangle ABG =$  the  $\triangle DEF$  ; *Proved.*

$\therefore$  the  $\triangle ABC : \text{the } \triangle DEF$  in the duplicate ratio of  $BC : EF$ . Q.E.D.

QUESTIONS FOR REVISION, AND NUMERICAL ILLUSTRATIONS.

1. Quote the Geometrical and Algebraical definitions of the *duplicate of the ratio*  $a : b$ ; and deduce the latter from the former. Estimate numerically the duplicate of the ratio 36 : 21.

2. The smaller of two similar triangles has an area of 20 square feet, and two corresponding sides are 3 ft. 6 in. and 2 ft. 4 in. respectively: shew that the area of the greater triangle is 45 square feet.

3. XY is drawn parallel to BC, the base of a triangle ABC, to meet the other sides at X and Y: if AX and XB measure respectively 3 inches and 7 inches, shew that the areas of the triangles  $AXY$ ,  $ABC$  are in the ratio 9 : 100.

4. Two similar triangles have areas in the ratio 529 : 361; shew that any pair of homologous sides are to one another as 23 : 19.

5. When are similar figures said to be *similarly situated*? Shew that similar and similarly situated triangles are to one another in the duplicate ratio of their altitudes.

6. Two similar and similarly situated triangles have areas in the ratio 1369 : 1681; if the altitude of the greater is 10 ft. 3 in., shew that the altitude of the other is 1 foot less.

7. The sides of a triangle are 11, 23, 29; find the sides of a similar triangle whose area is 289 times that of the former.

8. Shew how to draw a straight line XY parallel to BC the base of a triangle ABC, so that the area of the triangle  $AXY$  may be *nine-sixteenths* of that of the triangle ABC.

9. XY is drawn parallel to the base BC of a triangle ABC, so that the triangle  $AXY$  has to the figure  $XBCY$  the ratio 4 : 5; shew that AB and AC are cut by XY in the ratio 2 : 1.

10. A triangle ABC is bisected by a straight line XY drawn parallel to the base BC. In what ratio is AB divided at X?


Hence shew how to bisect a triangle by a straight line drawn parallel to the base.

11. ABC is a triangle whose area is 16 square feet; and XY is drawn parallel to BC, dividing AB in the ratio 3 : 5; shew that if BY is joined, the area of the triangle  $BXY$  is 3 sq. ft. 108 sq. in.

12. ABC is a triangle right-angled at A, and AD is the perpendicular drawn from A to the hypotenuse: if the area of the triangle ABC is 54 square inches and AB is 1 foot, shew that the area of the triangle ADC is  $19\frac{1}{4}$  square inches.

## PROPOSITION 20. THEOREM.

Similar polygons may be divided into the same number of similar triangles, having the same ratio each to each that the polygons have; and the polygons are to one another in the duplicate ratio of their homologous sides.



Let  $ABCDE$ ,  $FGHKL$  be similar polygons, and let  $AB$  and  $FG$  be homologous sides.

Then (i) the polygons may be divided into the same number of similar triangles;

(ii) these triangles shall have each to each the same ratio that the polygons have;

(iii) the polygon  $ABCDE$  shall be to the polygon  $FGHKL$  in the duplicate ratio of  $AB$  to  $FG$ .

Join  $EB$ ,  $EC$ ,  $LG$ ,  $LH$ .

(i) Then because the polygon  $ABCDE$  is similar to the polygon  $FGHKL$ ,

$\therefore$  the  $\angle EAB =$  the  $\angle LFG$ ,

and  $EA : AB :: LF : FG$ ; VI. Def. 2.

$\therefore$  the  $\triangle EAB$  is similar to the  $\triangle LFG$ ; VI. 6.

$\therefore$  the  $\angle ABE =$  the  $\angle FGL$ .

But because the polygons are similar, Hyp.

$\therefore$  the  $\angle ABC =$  the  $\angle FGH$ ; VI. Def. 2.

$\therefore$  the remaining  $\angle EBC =$  the remaining  $\angle LGH$ .

And because the  $\triangle^s EAB$ ,  $LFG$  are similar, Proved.

$\therefore EB : BA :: LG : GF$ ;

and because the polygons are similar, Hyp.

$\therefore AB : BC :: FG : GH$ ; VI. Def. 2.

$\therefore$ , *ex æquali*,  $EB : BC :: LG : GH$ ; V. 14.

that is, the sides about the equal  $\angle^s EBC$ ,  $LGH$  are proportionals;

$\therefore$  the  $\triangle EBC$  is similar to the  $\triangle LGH$ . VI. 6.

In the same way it may be proved that the  $\triangle ECD$  is similar to the  $\triangle LHK$ .

$\therefore$  the polygons have been divided into the same number of similar triangles.

(ii) Again, because the  $\triangle EAB$  is similar to the  $\triangle LFG$ ,  
 $\therefore$  the  $\triangle EAB$  is to the  $\triangle LFG$  in the duplicate ratio  
of  $EB : LG$ ; VI. 19.

and, in like manner,

the  $\triangle EBC$  is to the  $\triangle LGH$  in the duplicate ratio  
of  $EB$  to  $LG$ ;

$\therefore$  the  $\triangle EAB : \text{the } \triangle LFG :: \text{the } \triangle EBC : \text{the } \triangle LGH$ . V. 1.

In like manner it can be shewn that

the  $\triangle EBC : \text{the } \triangle LGH :: \text{the } \triangle ECD : \text{the } \triangle LHK$ ;

$\therefore$  the  $\triangle EAB : \text{the } \triangle LFG :: \text{the } \triangle EBC : \text{the } \triangle LGH$   
 $:: \text{the } \triangle ECD : \text{the } \triangle LHK$ .

But in a series of equal ratios, as each antecedent is to its consequent so is the sum of the antecedents to the sum of the consequents; [Addendo. v. 12.]

$\therefore$  the  $\triangle EAB : \text{the } \triangle LFG :: \text{the fig. } ABCDE : \text{the fig. } FGHKL$ .

(iii) Now the  $\triangle EAB : \text{the } \triangle LFG$  in the duplicate ratio  
of  $AB : FG$ , VI. 19.

and the  $\triangle EAB : \text{the } \triangle LFG :: \text{the fig. } ABCDE : \text{the fig. } FGHKL$ ;

$\therefore$  the fig.  $ABCDE : \text{the fig. } FGHKL$  in the duplicate ratio  
of  $AB : FG$ . Q.E.D.

**COROLLARY 1.** Let a third proportional  $X$  be taken to  $AB$  and  $FG$ ,

then  $AB$  is to  $X$  in the duplicate ratio of  $AB : FG$ ;

but the fig.  $ABCDE : \text{the fig. } FGHKL$  in the duplicate  
ratio of  $AB : FG$ ; *Proved.*

$\therefore AB : X :: \text{the fig. } ABCDE : \text{the fig. } FGHKL$ .

*Hence, if three straight lines are proportionals, as the first is to the third, so is any rectilineal figure described on the first to a similar and similarly described rectilineal figure on the second.*


**COROLLARY 2.** It follows that *similar rectilineal figures are to one another as the squares on their homologous sides.* For squares are similar figures and therefore are to one another in the duplicate ratio of their sides.



*Obs.* The following theorem, taken from Euclid's Twelfth Book, is given here as an important application of the preceding proposition.

BOOK XII. PROPOSITION 1.

*The areas of similar polygons inscribed in circles are to one another as the squares on the diameters.*



Let  $ABCDE$  and  $FGHKL$  be two similar polygons, inscribed in the circles  $ACE$ ,  $FHL$ , of which  $AM$ ,  $FN$  are diameters.

*Then shall*

*the fig.  $ABCDE$  : the fig.  $FGHKL$  :: the sq. on  $AM$  : the sq. on  $FN$ .*

Join  $BM$ ,  $AC$  and  $GN$ ,  $FH$ .

Then since the polygon  $ABCDE$  is similar to the polygon  $FGHKL$ ,

$\therefore$  the  $\angle ABC =$  the  $\angle FGH$ ,

and  $AB : BC :: FG : GH$ ; vi. Def. 2.

$\therefore$  the  $\triangle ABC$  is similar to the  $\triangle FGH$ ; vi. 6.

$\therefore$  the  $\angle ACB =$  the  $\angle FHG$ .

But the  $\angle ACB =$  the  $\angle AMB$ ; iii. 21.

and the  $\angle FHG =$  the  $\angle FNG$ ;

$\therefore$  the  $\angle AMB =$  the  $\angle FNG$ .

Also in the  $\triangle ABM$ ,  $FGN$ , the  $\angle ABM$ ,  $FGN$  are equal, being rt. angles; iii. 31.

hence the remaining  $\angle BAM$ ,  $GFN$  are equal; i. 32.

and the  $\triangle ABM$ ,  $FGN$  are similar: vi. 4.

$\therefore AB : FG :: AM : FN$ .

But the fig.  $ABCDE$  : the fig.  $FGHKL$  in the duplicate ratio of  $AB : FG$ , vi. 20.

that is, in the duplicate ratio of  $AM : FN$ . v. 16.

Hence

the fig.  $ABCDE$  : the fig.  $FGHKL$  :: the sq. on  $AM$  : the sq. on  $FN$ . vi. 20, Cor. 2.

*Obs.* The following theorem, which forms Proposition 3 of Euclid's Twelfth Book, may be derived as a corollary from the preceding proof.

**COROLLARY.** *The areas of circles are to one another as the squares on their diameters.*

It has been shewn that  
the fig. ABCDE : the fig. FGHKL :: the sq. on AM : the sq. on FN :  
and this is true however many sides the two polygons may have.

Suppose the polygons are *regular*; then by sufficiently increasing the number of their sides, we may make their areas differ from the areas of their circumscribed circles by quantities smaller than any that can be named; hence ultimately,

the  $\odot$  ACE : the  $\odot$  FHL :: the sq. on AM : the sq. on FN.

#### EXERCISES ON PROPOSITIONS 19, 20.

1. If ABC is a triangle right-angled at A, and AD is drawn perpendicular to BC, shew that

- (i) CB : BD in the duplicate ratio of CB to BA ;
- (ii) The square on CB : the square on BA :: CB : BD ;
- (iii) The  $\triangle$  ABD : the  $\triangle$  CAD in the duplicate ratio of BA to AC.

2. In any triangle ABC, the sides AB, AC are cut by a line XY drawn parallel to BC. If AX is one-third of AB, what part is the triangle AXY of the triangle ABC ?

3. A trapezium ABCD has its sides AB, CD parallel, and its diagonals intersect at O. If AB is double of CD, find the ratio of the triangle AOB to the triangle COD.

4. ABC and XYZ are two similar triangles whose areas are respectively 245 and 5 square inches. If AB is 21 inches in length, find XY.


5. Shew how to draw a straight line XY parallel to the base BC of a triangle ABC, so that the area of the triangle AXY may be four-ninths of the triangle ABC.

6. Two circles intersect at A and B, and at A tangents are drawn, one to each circle, meeting the circumferences at C and D. If AB, CB and BD are joined, shew that

the  $\triangle$  CBA : the  $\triangle$  ABD :: CB : BD.

## PROPOSITION 21. THEOREM.

*Rectilinear figures which are similar to the same rectilinear figure, are also similar to each other.*



Let each of the rectilinear figures A and B be similar to C.

*Then shall A be similar to B.*

For because A is similar to C, *Hyp.*

$\therefore$  A is equiangular to C,  
and the sides about their equal angles are proportionals.  
VI. Def. 2.

Again, because B is similar to C, *Hyp.*

$\therefore$  B is equiangular to C,  
and the sides about their equal angles are proportionals.  
VI. Def. 2.

$\therefore$  A and B are each of them equiangular to C, and have their sides about the equal angles proportional to the corresponding sides of C;

$\therefore$  A is equiangular to B, *Ax. 1.*  
and the sides of A and B about their equal angles are proportionals;  
V. 1.


$\therefore$  A is similar to B.

Q.E.D.

PROPOSITION 22. THEOREM.

If four straight lines be proportional and a pair of similar rectilinear figures be similarly described on the first and second, and also a pair on the third and fourth, these figures shall be proportional.

Conversely, if a rectilinear figure on the first of four straight lines be to the similar and similarly described figure on the second as a rectilinear figure on the third is to the similar and similarly described figure on the fourth, the four straight lines shall be proportional.



First. Let AB, CD, EF, GH be proportionals,  
 so that  $AB : CD :: EF : GH$  ;  
 and let similar figures KAB, LCD be similarly described on  
 AB, CD, and also let similar figures MF, NH be similarly  
 described on EF, GH.

Then shall

the fig. KAB : the fig. LCD :: the fig. MF : the fig. NH.

To AB and CD take a *third* proportional X ; VI. 11.  
 and to EF and GH take a *third* proportional O ;

then  $AB : CD :: CD : X$ , Constr.

and  $EF : GH :: GH : O$ .

But  $AB : CD :: EF : GH$  ; Hyp.

$\therefore CD : X :: GH : O$ , V. 1.


$\therefore$ , *ex æquali*,  $AB : X :: EF : O$ . V. 14.

But  $AB : X ::$  the fig. KAB : the fig. LCD ; VI. 20, Cor.

and  $EF : O ::$  the fig. MF : the fig. NH ;

$\therefore$  the fig. KAB : the fig. LCD :: the fig. MF : the fig. NH.

V. 1.



*Conversely.*

Let the fig. KAB : the fig. LCD :: the fig. MF : the fig. NH.

*Then shall*  $AB : CD :: EF : GH$ .

To AB, CD, and EF take a *fourth* proportional PR : VI. 12.  
and on PR describe the fig. SR similar and similarly situated  
to either of the figs. MF, NH. VI. 18.

Then because  $AB : CD :: EF : PR$ , *Constr.*

$\therefore$ , by the former part of the proposition,  
the fig. KAB : the fig. LCD :: the fig. MF : the fig. SR.

But, *by hypothesis*,

the fig. KAB : the fig. LCD :: the fig. MF : the fig. NH ;

$\therefore$  the fig. MF : the fig. SR :: the fig. MF : the fig. NH, v. 1.

$\therefore$  the fig. SR = the fig. NH.

And since the figs. SR and NH are similar and similarly  
situated, *Constr.*

$\therefore PR = GH^*$ .

Now  $AB : CD :: EF : PR$ ; *Constr.*

$\therefore AB : CD :: EF : GH$ .


Q.E.D.

\* Euclid here assumes that if two similar and similarly situated figures are equal, their homologous sides are equal. The proof is easy and may be left as an exercise for the student.

**DEFINITION.** When there are any number of magnitudes of the same kind, the *first* is said to have to the *last* the **ratio compounded** of the ratios of the *first* to the *second*, of the *second* to the *third*, and so on up to the ratio of the *last but one* to the *last* magnitude. [Book v. Def. 11.]

PROPOSITION 23. THEOREM.

*Parallelograms which are equiangular to one another have to one another the ratio which is compounded of the ratios of their sides.*



Let the par<sup>m</sup> AC be equiangular to the par<sup>m</sup> CF, having the  $\angle BCD$  equal to the  $\angle ECG$ .

*Then shall the par<sup>m</sup> AC have to the par<sup>m</sup> CF the ratio compounded of the ratios BC : CG and DC : CE.*

Let the par<sup>m</sup> be placed so that BC and CG are in a st. line; then DC and CE are also in a st. line. I. 14.

Complete the par<sup>m</sup> DG.

Take any st. line K,  
and to BC, CG, and K find a *fourth* proportional L; VI. 12.  
and to DC, CE, and L take a *fourth* proportional M;  
then  $BC : CG :: K : L$ ,  
and  $DC : CE :: L : M$ .

But  $K : M$  is the ratio compounded of the ratios  
 $K : L$  and  $L : M$ ; v. Def. 11.  
that is,  $K : M$  is the ratio compounded of the ratios  
 $BC : CG$  and  $DC : CE$ .


Now the par<sup>m</sup> AC : the par<sup>m</sup> DG :: BC : CG VI. 1.  
:: K : L, Constr.  
and the par<sup>m</sup> DG : the par<sup>m</sup> CF :: DC : CE VI. 1.  
:: L : M; Constr.

$\therefore$ , *ex æquali*, the par<sup>m</sup> AC : the par<sup>m</sup> CF :: K : M. v. 14.  
But  $K : M$  is the ratio compounded of the ratios of the sides;  
 $\therefore$  the par<sup>m</sup> AC has to the par<sup>m</sup> CF the ratio compounded  
of the ratios of the sides. Q.E.D.

**EXERCISE.** The areas of two triangles or parallelograms are to one another in the ratio compounded of the ratios of their bases and of their altitudes,

## PROPOSITION 24. THEOREM.

*Parallelograms about a diagonal of any parallelogram are similar to the whole parallelogram and to one another.*



Let  $ABCD$  be a  $\text{par}^m$  of which  $AC$  is a diagonal ;  
and let  $EG, HK$  be  $\text{par}^{ms}$  about  $AC$ .

*Then shall the  $\text{par}^{ms}$   $EG, HK$  be similar to the  $\text{par}^m$   $ABCD$ , and to one another.*

For because  $DC$  is  $\text{par}^1$  to  $GF$ ,  
 $\therefore$  the  $\angle ADC =$  the  $\angle AGF$  ; I. 29.

and because  $BC$  is  $\text{par}^1$  to  $EF$ ,  
 $\therefore$  the  $\angle ABC =$  the  $\angle AEF$  ; I. 29.

and each of the  $\angle^s$   $BCD, EFG$  is equal to the opp.  $\angle$   $BAD$ ,  
 $\therefore$  the  $\angle BCD =$  the  $\angle EFG$  ; I. 34.

$\therefore$  the  $\text{par}^m$   $ABCD$  is equiangular to the  $\text{par}^m$   $AEGF$ .

Again in the  $\triangle^s$   $BAC, EAF$ ,  
because the  $\angle ABC =$  the  $\angle AEF$ , I. 29.  
and the  $\angle BAC$  is common ;

$\therefore$  the remaining  $\angle BCA =$  the remaining  $\angle EFA$  ; I. 32.

$\therefore$  the  $\triangle^s$   $BAC, EAF$  are equiangular to one another ;  
 $\therefore$   $AB : BC :: AE : EF$ . VI. 4.

But  $BC = AD$ , and  $EF = AG$  ; I. 34.

$\therefore$   $AB : AD :: AE : AG$ .

Similarly  $DC : CB :: GF : FE$ ,  
and  $CD : DA :: FG : GA$  ;

$\therefore$  the sides of the  $\text{par}^{ms}$   $ABCD, AEGF$  about their equal angles are proportional ;

$\therefore$  the  $\text{par}^m$   $ABCD$  is similar to the  $\text{par}^m$   $AEGF$ . VI. Def. 2.


In the same way the  $\text{par}^m$   $ABCD$  may be proved similar to the  $\text{par}^m$   $FHCK$ ,

$\therefore$  each of the  $\text{par}^{ms}$   $EG, HK$  is similar to the whole  $\text{par}^m$  ;  
 $\therefore$  the  $\text{par}^m$   $EG$  is similar to the  $\text{par}^m$   $HK$ . VI. 21.

Q.E.D.

PROPOSITION 25. PROBLEM.

To describe a rectilinear figure which shall be equal to one and similar to another rectilinear figure.



Let E and S be the two given rectilinear figures.

It is required to describe a figure equal to the fig. E and similar to the fig. S.

On AB a side of the fig. S describe a par<sup>m</sup> ABCD equal to S ;  
 and on BC describe a par<sup>m</sup> CBGF equal to the fig. E, and  
 having the  $\angle$  CBG equal to the  $\angle$  DAB ; I. 45.  
 then AB and BG are in one st. line, and also DC and CF in  
 one st. line.

Between AB and BG find a mean proportional HK ; VI. 13.  
 and on HK describe the fig. P, similar and similarly situated  
 to the fig. S. VI. 18.

Then P shall be the figure required.

Because  $AB : HK :: HK : BG$ , Constr.

$\therefore AB : BG ::$  the fig. S : the fig. P. VI. 20, Cor.

But  $AB : BG ::$  the par<sup>m</sup> AC : the par<sup>m</sup> BF ; VI. 1.  
 $\therefore$  the fig. S : the fig. P :: the par<sup>m</sup> AC : the par<sup>m</sup> BF ; V. 1.  
 and the fig. S = the par<sup>m</sup> AC ; Constr.  
 $\therefore$  the fig. P = the par<sup>m</sup> BF  
 = the fig. E. Constr.


And since, by construction, the fig. P is similar to the fig. S,  
 $\therefore$  P is the figure required.

Q.E.F.



## PROPOSITION 26. THEOREM.

*If two similar parallelograms have a common angle, and are similarly situated, they are about the same diagonal.*



Let the par<sup>ms</sup> ABCD, AEFB be similar and similarly situated, and have the common angle BAD.

*Then shall the par<sup>ms</sup> ABCD, AEFB be about the same diagonal.*

Join AC.

Then if AC does not pass through F, if possible let it cut FG, or FG produced, at H.

Through H draw HK par<sup>l</sup> to AD or BC. I. 31.

Then the par<sup>ms</sup> BD and KG are similar, since they are about the same diagonal AHC; VI. 24.

$$\therefore DA : AB :: GA : AK.$$

But because the par<sup>ms</sup> BD and EG are similar; *Hyp.*

$$\therefore DA : AB :: GA : AE; \quad \text{VI. Def. 2.}$$

$$\therefore GA : AK :: GA : AE;$$

$$\therefore AK = AE, \text{ which is impossible;}$$

$$\therefore AC \text{ must pass through F;}$$

that is, the par<sup>ms</sup> BD, EG are about the same diagonal.

Q.E.D.


*Obs.* Propositions 27, 28, 29 being cumbrous in form and of little value as geometrical results are now very generally omitted.

**DEFINITION.** A straight line is said to be divided in **extreme and mean ratio**, when the whole is to the greater segment as the greater segment is to the less.

[Book VI. Def. 5.]

**PROPOSITION 30. PROBLEM.**

*To divide a given straight line in extreme and mean ratio.*



Let **AB** be the given st. line.

*It is required to divide AB in extreme and mean ratio.*

Divide **AB** in **C** so that the rect. **AB, BC** may be equal to the sq. on **AC**. II. 11.


Then because the rect. **AB, BC** = the sq. on **AC**,  
 $\therefore AB : AC :: AC : BC$ . VI. 17.  
Q.E.F.

**EXERCISES.**

1. **ABCDE** is a regular pentagon; if the lines **BE** and **AD** intersect in **O**, shew that each of them is divided in extreme and mean ratio.
2. If the radius of a circle is cut in extreme and mean ratio, the greater segment is equal to the side of a regular decagon inscribed in the circle.

## PROPOSITION 31. THEOREM.

*In a right-angled triangle, any rectilineal figure described on the hypotenuse is equal to the sum of the two similar and similarly described figures on the sides containing the right angle.*



Let  $ABC$  be a right-angled triangle of which  $BC$  is the hypotenuse; and let  $P$ ,  $Q$ ,  $R$  be similar and similarly described figures on  $BC$ ,  $CA$ ,  $AB$  respectively.

*Then shall the fig.  $P$  be equal to the sum of the figs.  $Q$  and  $R$ .*

Draw  $AD$  perp. to  $BC$ .

Then the  $\triangle^s CBA$ ,  $ABD$  are similar; VI. 8.

$\therefore CB : BA :: BA : BD$ ;

$\therefore CB : BD ::$  the fig.  $P$  : the fig.  $R$ ; VI. 20, *Cor.*

$\therefore$ , *inversely*,  $BD : BC ::$  the fig.  $R$  : the fig.  $P$ . V. 2.

In like manner,  $DC : BC ::$  the fig.  $Q$  : the fig.  $P$ ;

$\therefore$  the sum of  $BD$ ,  $DC : BC ::$  the sum of figs.  $R$ ,  $Q$  : fig.  $P$ ;  
V. 15.

but  $BC =$  the sum of  $BD$ ,  $DC$ ;

$\therefore$  the fig.  $P =$  the sum of the figs.  $R$  and  $Q$ .

Q.E.D.

NOTE. This proposition is a generalization of Book I., Prop. 47. It will be a useful exercise for the student to deduce the general theorem (VI. 31) from the particular case (I. 47) with the aid of VI. 20, Cor. 2.

## EXERCISES.

1. In a right-angled triangle if a perpendicular is drawn from the right angle to the opposite side, the segments of the hypotenuse are in the duplicate ratio of the sides containing the right angle.

2. If, in Proposition 31, the figure on the hypotenuse is equal to the given triangle, the figures on the other two sides are respectively equal to the parts into which the triangle is divided by the perpendicular from the right angle to the hypotenuse.

3. AX and BY are medians of the triangle ABC which meet in G : if XY is joined, compare the areas of the triangles AGB, XGY.

4. *Shew that similar triangles are to one another in the duplicate ratio of (i) corresponding medians, (ii) the radii of their inscribed circles, (iii) the radii of their circumscribed circles.*

5. DEF is the pedal triangle of the triangle ABC ; prove that the triangle ABC is to the triangle DBF in the duplicate ratio of AB to BD. Hence shew that

$$\text{the fig. AFDC} : \text{the } \triangle \text{ BFD} :: \text{AD}^2 : \text{BD}^2.$$

6. The base BC of a triangle ABC is produced to a point D such that BD : DC in the duplicate ratio of BA : AC. Shew that AD is a mean proportional between BD and DC.

7. Bisect a triangle by a line drawn parallel to one of its sides.

8. Shew how to draw a line parallel to the base of a triangle so as to form with the other two sides produced a triangle double of the given triangle.

9. If through any point within a triangle lines are drawn from the angles to cut the opposite sides, the segments of any one side will have to each other the ratio compounded of the ratios of the segments of the other sides.


10. Draw a straight line parallel to the base of an isosceles triangle so as to cut off a triangle which has to the whole triangle the ratio of the base to a side.

11. Through a given point, between two straight lines containing a given angle, draw a line which shall cut off a triangle equal to a given rectilineal figure.

*Obs.* The 32nd Proposition as given by Euclid is defective, and as it is never applied, we have omitted it.

## PROPOSITION 33. THEOREM.

*In equal circles, angles, whether at the centres or the circumferences, have the same ratio as the arcs on which they stand: so also have the sectors.*



Let  $ABC$  and  $DEF$  be equal circles, and let  $BGC$ ,  $EHF$  be angles at the centres, and  $BAC$  and  $EDF$  angles at the  $\odot^{\circ}$ .

Then shall

- (i) the  $\angle BGC$  : the  $\angle EHF$  :: the arc  $BC$  : the arc  $EF$  ;
- (ii) the  $\angle BAC$  : the  $\angle EDF$  :: the arc  $BC$  : the arc  $EF$  ;
- (iii) the sector  $BGC$  : the sector  $EHF$  :: the arc  $BC$  : the arc  $EF$ .

Along the  $\odot^{\circ}$  of the  $\odot ABC$  take *any* number of arcs  $CK$ ,  $KL$  each equal to  $BC$  ;  
and along the  $\odot^{\circ}$  of the  $\odot DEF$  take *any* number of arcs  $FM$ ,  $MN$ ,  $NR$  each equal to  $EF$ .

Join  $GK$ ,  $GL$ ,  $HM$ ,  $HN$ ,  $HR$ .

(i) Then the  $\angle BGC$ ,  $CGK$ ,  $KGL$  are all equal,  
for they stand on the equal arcs  $BC$ ,  $CK$ ,  $KL$  : III. 27.  
 $\therefore$  the  $\angle BGL$  is the same multiple of the  $\angle BGC$  that the arc  $BL$  is of the arc  $BC$ .

Similarly, the  $\angle EHR$  is the same multiple of the  $\angle EHF$  that the arc  $ER$  is of the arc  $EF$ .

And if the arc  $BL =$  the arc  $ER$ ,

the  $\angle BGL =$  the  $\angle EHR$  ;

III. 27.

and if the arc  $BL$  is greater than the arc  $ER$ ,

the  $\angle BGL$  is greater than the  $\angle EHR$  ;

and if the arc  $BL$  is less than the arc  $ER$ ,

the  $\angle BGL$  is less than the  $\angle EHR$ .


Now since there are four magnitudes, namely the  $\angle^s$  BGC, EHF and the arcs BC, EF; and of the antecedents any equimultiples have been taken, namely the  $\angle$  BGL and the arc BL; and of the consequents any equimultiples have been taken, namely the  $\angle$  EHR and the arc ER:

and since it has been proved that the  $\angle$  BGL is greater than, equal to, or less than the  $\angle$  EHR, according as BL is greater than, equal to, or less than ER;

$\therefore$  the four original magnitudes are proportionals; v. *Def.* 5.  
that is, the  $\angle$  BGC : the  $\angle$  EHF :: the arc BC : the arc EF.

(ii) And since the  $\angle$  BGC = twice the  $\angle$  BAC, III. 20.  
and the  $\angle$  EHF = twice the  $\angle$  EDF;

$\therefore$  the  $\angle$  BAC : the  $\angle$  EDF :: the arc BC : the arc EF. v. 8.



(iii) Join BC, CK; and in the arcs BC, CK take any points X, O.

Join BX, XC, CO, OK.

Then in the  $\triangle^s$  BGC, CGK,

Because  $\left\{ \begin{array}{l} BG = CG, \\ GC = GK, \\ \text{and the } \angle BGC = \text{the } \angle CGK; \end{array} \right. \quad \text{III. 27.}$

$\therefore BC = CK;$  I. 4.

and the  $\triangle BGC = \text{the } \triangle CGK.$

And because the arc BC = the arc CK, *Constr.*

$\therefore$  the remaining arc BAC = the remaining arc CAK:

$\therefore$  the  $\angle BXC = \text{the } \angle COK;$  III. 27.

$\therefore$  the segment BXC is similar to the segment COK;


III. *Def.* 10.

and these segments stand on equal chords BC, CK;

$\therefore$  the segment BXC = the segment COK. III. 24.

And the  $\triangle BGC = \text{the } \triangle CGK;$

$\therefore$  the sector BGC = the sector CGK.



Similarly it may be shewn that the sectors BGC, CGK, KGL are all equal ;  
 and likewise the sectors EHF, FHM, MHN, NHR are all equal.  
 $\therefore$  the sector BGL is the same multiple of the sector BGC  
 that the arc BL is of the arc BC ;  
 and the sector EHR is the same multiple of the sector EHF  
 that the arc ER is of the arc EF.

And if the arc BL = the arc ER,  
 the sector BGL = the sector EHR : *Proved.*  
 and if the arc BL is greater than the arc ER,  
 the sector BGL is greater than the sector EHR :  
 and if the arc BL is less than the arc ER,  
 the sector BGL is less than the sector EHR.

Now since there are four magnitudes, namely, the sectors BGC, EHF and the arcs BC, EF ; and of the antecedents any equimultiples have been taken, namely the sector BGL and the arc BL ; and of the consequents any equimultiples have been taken, namely the sector EHR and the arc ER :  
 and since it has been shewn that the sector BGL is greater than, equal to, or less than the sector EHR, according as the arc BL is greater than, equal to, or less than the arc ER ;

$\therefore$  the four original magnitudes are proportionals ;  
 v. *Def.* 5.

that is,  
 the sector BGC : the sector EHF :: the arc BC : the arc EF.  
 Q.E.D.

## QUESTIONS FOR REVISION.

1. Explain why the operation known as *Alternately* requires that the four terms of a proportion should be of the same kind. Shew that this is unnecessary in the case of *Inversely*.

2. State and prove algebraically the theorem known as *Componendo*. In what proposition is this principle applied?

3. Enunciate and prove algebraically the operation used in Book vi. under the name *Ex Aequali*.

Also prove the same theorem in the following more general form:

*If there are two sets of magnitudes, such that the first is to the second of the first set as the first to the second of the other set, and the second to the third of the first set as the second to the third of the other, and so on throughout: then the first shall be to the last of the first set as the first to the last of the other.*

4. Explain the operation *Addendo*, and give an algebraical proof of it. In what proposition of Book vi. is this operation employed?

5. Give the geometrical and algebraical definitions of the *ratio compounded of given ratios*, and shew that the two definitions agree.

By what artifice would Euclid represent the ratio compounded of the ratios  $A : B$  and  $C : D$ ?

6. Two parallelograms ABCD, EFGH are equiangular to one another: if AB, BC are respectively 21 and 18 inches in length, and if EF, FG are 27 and 35 inches; shew that the areas of the parallelograms are in the ratio 2 : 5.

7. If  $A : B = X : Y$ , and  $C : B = Z : Y$ ;  
shew that  $A + C : B = X + Z : Y$ .

In what proposition of Book vi. is this principle used?

Explain and illustrate the necessity of the step *invertendo* in this proposition.

8. When is a straight line said to be divided in *extreme and mean ratio*?


If a line 10 inches in length is so divided, shew that the lengths of the segments are approximately 6.2 inches and 3.8 inches.

Shew also that the segments of *any* line divided in *extreme and mean ratio* are incommensurable.



## PROPOSITION B. THEOREM.

If the vertical angle of a triangle be bisected by a straight line which cuts the base, the rectangle contained by the sides of the triangle shall be equal to the rectangle contained by the segments of the base, together with the square on the straight line which bisects the angle.



Let  $ABC$  be a triangle, having the  $\angle BAC$  bisected by  $AD$ .  
Then shall  
the rect.  $BA, AC =$  the rect.  $BD, DC$ , with the sq. on  $AD$ .

Describe a circle about the  $\triangle ABC$ , IV. 5.  
and produce  $AD$  to meet the  $\circ^{\circ}$  in  $E$ .  
Join  $EC$ .

Then in the  $\triangle BAD, EAC$ ,  
because the  $\angle BAD =$  the  $\angle EAC$ , Hyp.  
and the  $\angle ABD =$  the  $\angle AEC$  in the same segment; III. 21.  
 $\therefore$  the remaining  $\angle BDA =$  the remaining  $\angle ECA$ ; I. 32.  
that is, the  $\triangle BAD$  is equiangular to the  $\triangle EAC$ .  
 $\therefore BA : AD :: EA : AC$ ; VI. 4.  
 $\therefore$  the rect.  $BA, AC =$  the rect.  $EA, AD$ , VI. 16.  
 $=$  the rect.  $ED, DA$ , with the sq. on  $AD$ . II. 3.


But the rect.  $ED, DA =$  the rect.  $BD, DC$ ; III. 35.  
 $\therefore$  the rect.  $BA, AC =$  the rect.  $BD, DC$ , with the sq. on  $AD$ .  
Q. E. D.

## EXERCISE.

If the vertical angle  $BAC$  is *externally* bisected by a straight line which meets the base in  $D$ , shew that the rectangle contained by  $BA, AC$  together with the square on  $AD$  is equal to the rectangle contained by the segments of the base.

PROPOSITION C. THEOREM.

*If from the vertical angle of a triangle a straight line be drawn perpendicular to the base, the rectangle contained by the sides of the triangle shall be equal to the rectangle contained by the perpendicular and the diameter of the circle described about the triangle.*



Let  $ABC$  be a triangle, and let  $AD$  be the perp. from  $A$  to the base  $BC$ .

*Then the rect.  $BA, AC$  shall be equal to the rectangle contained by  $AD$  and the diameter of the circle circumscribed about the  $\triangle ABC$ .*

Describe a circle about the  $\triangle ABC$ ;                    IV. 5.  
draw the diameter  $AE$ , and join  $EC$ .

Then in the  $\triangle BAD, EAC$ ,  
the rt. angle  $BDA =$  the rt. angle  $ECA$ , in the semicircle  $ECA$ ,  
and the  $\angle ABD =$  the  $\angle AEC$ , in the same segment; III. 21.

$\therefore$  the remaining  $\angle BAD =$  the remaining  $\angle EAC$ ; I. 32.

that is, the  $\triangle BAD$  is equiangular to the  $\triangle EAC$ ;


$\therefore BA : AD :: EA : AC$ ;    VI. 4.

$\therefore$  the rect.  $BA, AC =$  the rect.  $EA, AD$ .                    VI. 16.

Q.E.D.

## PROPOSITION D. THEOREM.

*The rectangle contained by the diagonals of a quadrilateral inscribed in a circle is equal to the sum of the two rectangles contained by its opposite sides.*



Let ABCD be a quadrilateral inscribed in a circle, and let AC, BD be its diagonals.

*Then the rect. AC, BD shall be equal to the sum of the rectangles AB, CD and BC, AD.*

Make the  $\angle DAE$  equal to the  $\angle BAC$ ; I. 23.  
to each add the  $\angle EAC$ ,  
then the  $\angle DAC =$  the  $\angle EAB$ .

Then in the  $\triangle EAB, DAC$ ,  
the  $\angle EAB =$  the  $\angle DAC$ ,

and the  $\angle ABE =$  the  $\angle ACD$  in the same segment; III. 21.

$\therefore$  the  $\triangle EAB, DAC$  are equiangular to one another; I. 32.

$\therefore AB : BE :: AC : CD$ ; VI. 4.

$\therefore$  the rect. AB, CD = the rect. AC, EB. VI. 16.

Again in the  $\triangle DAE, CAB$ ,  
the  $\angle DAE =$  the  $\angle CAB$ ,

*Constr.*

and the  $\angle ADE =$  the  $\angle ACB$ , in the same segment, III. 21.

$\therefore$  the  $\triangle DAE, CAB$  are equiangular to one another; I. 32.

$\therefore AD : DE :: AC : CB$ ; VI. 4.

$\therefore$  the rect. BC, AD = the rect. AC, DE. VI. 16.

But the rect. AB, CD = the rect. AC, EB. *Proved.*

$\therefore$  the sum of the rects. BC, AD and AB, CD = the sum of the rects. AC, DE and AC, EB;

that is, the sum of the rects. BC, AD and AB, CD

= the rect. AC, BD. II. 1.

Q.E.D.

**NOTE.** Propositions B, C, and D do not occur in Euclid, but were added by Robert Simson, who edited Euclid's text in 1756.

Prop. D is usually known as Ptolemy's theorem, and it is the particular case of the following more general theorem :

*The rectangle contained by the diagonals of a quadrilateral is less than the sum of the rectangles contained by its opposite sides, unless a circle can be circumscribed about the quadrilateral, in which case it is equal to that sum.*

## EXERCISES.

1. ABC is an isosceles triangle, and on the base, or base produced, any point X is taken : shew that the circumscribed circles of the triangles ABX, ACX are equal.

2. From the extremities B, C of the base of an isosceles triangle ABC, straight lines are drawn perpendicular to AB, AC respectively, and intersecting at D : shew that the rectangle BC, AD is double of the rectangle AB, DB.

3. If the diagonals of a quadrilateral inscribed in a circle are at right angles, the sum of the rectangles contained by the opposite sides is double the area of the figure.

4. ABCD is a quadrilateral inscribed in a circle, and the diagonal BD bisects AC : shew that the rectangle AD, AB is equal to the rectangle DC, CB.

5. If the vertex A of a triangle ABC is joined to any point in the base, it will divide the triangle into two triangles such that their circumscribed circles have radii in the ratio of AB to AC.

6. Construct a triangle, having given the base, the vertical angle, and the rectangle contained by the sides.

7. Two triangles of equal area are inscribed in the same circle : shew that the rectangle contained by any two sides of the one is to the rectangle contained by any two sides of the other as the base of the second is to the base of the first.


8. A circle is described round an equilateral triangle, and from any point in the circumference straight lines are drawn to the angular points of the triangle : shew that one of these straight lines is equal to the sum of the other two.

9. ABCD is a quadrilateral inscribed in a circle, and BD bisects the angle ABC : if the points A and C are fixed on the circumference of the circle and B is variable in position, shew that the sum of AB and BC has a constant ratio to BD.

## THEOREMS AND EXAMPLES ON BOOK VI.

### I. ON HARMONIC SECTION.

1. *To divide a given straight line internally and externally so that its segments may be in a given ratio.*



Let  $AB$  be the given st. line, and  $L, M$  two other st. lines which determine the given ratio.

*It is required to divide  $AB$  internally and externally in the ratio  $L : M$ .*

Through  $A$  and  $B$  draw any two par<sup>l</sup> st. lines  $AH, BK$ .

From  $AH$  cut off  $Aa$  equal to  $L$ , and from  $BK$  cut off  $Bb$  and  $Bb'$  each equal to  $M$ ,  $Bb'$  being taken in the same direction as  $Aa$ , and  $Bb$  in the opposite direction.

Join  $ab$ , cutting  $AB$  in  $P$ ;

join  $ab'$ , and produce it to cut  $AB$  externally at  $Q$ .

Then  $AB$  shall be divided internally at  $P$  and externally at  $Q$ ,  
so that  
and


$$AP : PB = L : M.$$

$$AQ : QB = L : M.$$

The proof follows at once from Euclid vi. 4.

NOTE. The solution is *singular*; that is, only *one* internal and *one* external point can be found that will divide the given straight line into segments which have the given ratio.

**DEFINITION.** A finite straight line is said to be **cut harmonically** when it is divided internally and externally into segments which have the same ratio.



Thus AB is divided harmonically at P and Q, if

$$AP : PB = AQ : QB.$$

P and Q are said to be **harmonic conjugates** of A and B.

Now by taking the above proportion *alternately*, we have

$$PA : AQ = PB : BQ;$$

from which it is seen that if P and Q divide AB internally and externally in the same ratio, then A and B divide PQ internally and externally in the same ratio; hence A and B are harmonic conjugates of P and Q.

*Example.* The base of a triangle is divided harmonically by the internal and external bisectors of the vertical angle :

for in each case the segments of the base are in the ratio of the other sides of the triangle. [Euclid vi. 3 and A.]


*Obs.* We shall use the terms *Arithmetic, Geometric, and Harmonic Means* in their ordinary Algebraical sense.

1. If AB is divided internally at P and externally at Q in the same ratio, then AB is the harmonic mean between AQ and AP.

For, by hypothesis,  $AQ : QB = AP : PB$ ;  
 $\therefore$ , alternately,  $AQ : AP = QB : PB$ ;  
 that is,  $AQ : AP = AQ - AB : AB - AP$ ;  
 $\therefore$  AP, AB, AQ are in Harmonic Progression.

2. If AB is divided harmonically at P and Q, and O is the middle point of AB;

$$\text{then } OP \cdot OQ = OA^2.$$



For since AB is divided harmonically at P and Q,


$\therefore AP : PB = AQ : QB$ ;  
 $\therefore AP - PB : AP + PB = AQ - QB : AQ + QB$ ;  
 or,  $2OP : 2OA = 2OQ : 2OB$ ;  
 $\therefore OP \cdot OQ = OA^2$ .

*Conversely*, if  $OP \cdot OQ = OA^2$ ,  
 it may be shewn that

$AP : PB = AQ : QB$ ;  
 that is, that AB is divided harmonically at P and Q.

3. *The Arithmetic, Geometric and Harmonic means of two straight lines may be thus represented graphically.*

In the adjoining figure, two tangents AH, AK are drawn from any external point A to the circle PHQK; HK is the chord of contact, and the st. line joining A to the centre O cuts the  $\odot^{\text{ce}}$  at P and Q.



Then (i) AO is the Arithmetic mean between AP and AQ: for clearly  $AO = \frac{1}{2}(AP + AQ)$ .

(ii) AH is the Geometric mean between AP and AQ: for  $AH^2 = AP \cdot AQ$ . III. 36.

(iii) AB is the Harmonic mean between AP and AQ: for  $OA \cdot OB = OP^2$ ; Ex. 1, p. 251.  
 $\therefore$  AB is cut harmonically at P and Q. Ex. 2, p. 385.

That is, AB is the Harmonic mean between AP and AQ.

And from the similar triangles OAH, HAB,

$OA : AH = AH : AB$ ,  
 $\therefore AO \cdot AB = AH^2$ ; VI. 17.


$\therefore$  the Geometric mean between two straight lines is the mean proportional between their Arithmetic and Harmonic means.

4. *Given the base of a triangle and the ratio of the other sides, to find the locus of the vertex.*

Let BC be the given base, and let BAC be any triangle standing upon it, such that  $BA : AC =$  the given ratio.

It is required to find the locus of A.

Bisect the  $\angle$  BAC internally and externally by AP, AQ.



Then BC is divided internally at P, and externally at Q, so that  $BP : PC = BQ : QC =$  the given ratio;  $\therefore$  P and Q are fixed points.

And since AP, AQ are the internal and external bisectors of the  $\angle$  BAC,

$\therefore$  the  $\angle$  PAQ is a rt. angle;  
 $\therefore$  the locus of A is a circle described on PQ as diameter.

EXERCISE. Given three points B, P, C in a straight line: find the locus of points at which BP and PC subtend equal angles.

## DEFINITIONS.

1. A series of points in a straight line is called a **range**. If the range consists of four points, of which one pair are harmonic conjugates with respect to the other pair, it is said to be a **harmonic range**.

2. A series of straight lines drawn through a point is called a **pencil**.

The point of concurrence is called the **vertex** of the pencil, and each of the straight lines is called a **ray**.

A pencil of four rays drawn from any point to a harmonic range is said to be a **harmonic pencil**.

3. A straight line drawn to cut a system of lines is called a **transversal**.

4. A system of four straight lines, no three of which are concurrent, is called a **complete quadrilateral**.

These straight lines will intersect two and two in six points, called the **vertices** of the quadrilateral; the *three* straight lines which join the opposite vertices are **diagonals**.

## THEOREMS ON HARMONIC SECTION.

1. *If a transversal is drawn parallel to one ray of a harmonic pencil, the other three rays intercept equal parts upon it: and conversely.*

2. *Any transversal is cut harmonically by the rays of a harmonic pencil.*

3. *In a harmonic pencil, if one ray bisect the angle between the other pair of rays, it is perpendicular to its conjugate ray. Conversely, if one pair of rays form a right angle, then they bisect internally and externally the angle between the other pair.*

4. *If  $A, P, B, Q$  and  $a, p, b, q$  are harmonic ranges, one on each of two given straight lines, and if  $Aa, Pp, Bb$ , the straight lines which join three pairs of corresponding points, meet at  $S$ ; then will  $Qq$  also pass through  $S$ .*


5. *If two straight lines intersect at  $A$ , and if  $A, P, B, Q$  and  $A, p, b, q$  are two harmonic ranges one on each straight line (the points corresponding as indicated by the letters), then  $Pp, Bb, Qq$  will be concurrent: also  $Pq, Bb, Qp$  will be concurrent.*

6. *Use Theorem 5 to prove that in a complete quadrilateral in which the three diagonals are drawn, the straight line joining any pair of opposite vertices is cut harmonically by the other two diagonals.*



## II. ON CENTRES OF SIMILARITY AND SIMILITUDE.

1. *If any two unequal similar figures are placed so that their homologous sides are parallel, the lines joining corresponding points in the two figures meet in a point, whose distances from any two corresponding points are in the ratio of any pair of homologous sides.*



Let  $ABCD$ ,  $A'B'C'D'$  be two similar figures, and let them be placed so that their homologous sides are parallel; namely,  $AB$ ,  $BC$ ,  $CD$ ,  $DA$  parallel to  $A'B'$ ,  $B'C'$ ,  $C'D'$ ,  $D'A'$  respectively.

*Then shall  $AA'$ ,  $BB'$ ,  $CC'$ ,  $DD'$  meet in a point, whose distances from any two corresponding points shall be in the ratio of any pair of homologous sides.*

Let  $AA'$  meet  $BB'$ , produced if necessary, in  $S$ .

Then because  $AB$  is par<sup>l</sup> to  $A'B'$ ; *Hyp.*

$\therefore$  the  $\triangle SAB$ ,  $SA'B'$  are equiangular;

$\therefore SA : SA' = AB : A'B'$ ; VI. 4.

$\therefore AA'$  divides  $BB'$ , externally or internally, in the ratio of  $AB$  to  $A'B'$ .

Similarly it may be shewn that  $CC'$  divides  $BB'$  in the ratio of  $BC$  to  $B'C'$ .

But since the figures are similar,

$BC : B'C' = AB : A'B'$ ;

$\therefore AA'$  and  $CC'$  divide  $BB'$  in the same ratio;  
that is,  $AA'$ ,  $BB'$ ,  $CC'$  meet in the same point  $S$ .


In like manner it may be proved that  $DD'$  meets  $CC'$  in the point  $S$ .

$\therefore AA'$ ,  $BB'$ ,  $CC'$ ,  $DD'$  are concurrent, and each of these lines is divided at  $S$ , externally or internally, in the ratio of a pair of homologous sides of the two figures. Q.E.D.

*COR.* *If any line is drawn through  $S$  meeting any pair of homologous sides in  $K$  and  $K'$ , the ratio  $SK : SK'$  is constant, and equal to the ratio of any pair of homologous sides.*

**NOTE.** It will be seen that the lines joining corresponding points are divided externally or internally at  $S$  according as the corresponding sides are drawn in the same or in opposite directions. In either case the point of concurrence  $S$  is called a **centre of similarity** of the two figures.

2. A common tangent  $STT'$  to two circles whose centres are  $C, C'$ , meets the line of centres in  $S$ . If through  $S$  any straight line is drawn meeting these two circles in  $P, Q$ , and  $P', Q'$ , respectively, then the radii  $CP, CQ$  shall be respectively parallel to  $C'P', C'Q'$ . Also the rectangles  $SQ \cdot SP', SP \cdot SQ'$  shall each be equal to the rectangle  $ST \cdot ST'$ .



Join  $CT, CP, CQ$  and  $C'T', C'P', C'Q'$ .

Then since each of the  $\angle^s CTS, C'T'S$  is a right angle, III. 18.

$\therefore CT$  is par<sup>l</sup> to  $C'T'$ ;

$\therefore$  the  $\Delta^s SCT, SC'T'$  are equiangular;

$\therefore SC : SC' = CT : C'T'$   
 $= CP : C'P'$ ;

$\therefore$  the  $\Delta^s SCP, SC'P'$  are similar; VI. 7.

$\therefore$  the  $\angle SCP =$  the  $\angle SC'P'$ ;

$\therefore CP$  is par<sup>l</sup> to  $C'P'$ .

Similarly  $CQ$  is par<sup>l</sup> to  $C'Q'$ .

Again, it easily follows that  $TP, TQ$  are par<sup>l</sup> to  $T'P', T'Q'$  respectively;

$\therefore$  the  $\Delta^s STP, ST'P'$  are similar.

Now the rect.  $SP \cdot SQ =$  the sq. on  $ST$ ; III. 36.

$\therefore SP : ST = ST : SQ$ , VI. 16.

and  $SP : ST = SP' : ST'$ ;

$\therefore ST : SQ = SP' : ST'$ ;

$\therefore$  the rect.  $ST \cdot ST' = SQ \cdot SP'$ .

In the same way it may be proved that

the rect.  $SP \cdot SQ' =$  the rect.  $ST \cdot ST'$ .

Q.E.D.

COR. 1. It has been proved that

$SC : SC' = CP : C'P'$ ;

thus the external common tangents to the two circles meet at a point  $S$  which divides the line of centres externally in the ratio of the radii.

Similarly it may be shewn that the transverse common tangents meet at a point  $S'$  which divides the line of centres internally in the ratio of the radii.

COR. 2.  $CC'$  is divided harmonically at  $S$  and  $S'$ .

DEFINITION. The points  $S$  and  $S'$  which divide externally and internally the line of centres of two circles in the ratio of their radii are called the **external and internal centres of similitude** respectively.

## EXAMPLES ON CENTRES OF SIMILITUDE.

1. Inscribe a square in a given triangle.
2. In a given triangle inscribe a triangle similar and similarly situated to a given triangle.
3. Inscribe a square in a given sector of circle, so that two angular points shall be on the arc of the sector and the other two on the bounding radii.
4. *In the figure on page 298, if  $D_1$  meets the inscribed circle in  $X$ , shew that  $A, X, D_1$  are collinear. Also if  $A_1$  meets the base in  $Y$  shew that  $AA_1$  is divided harmonically at  $Y$  and  $A$ .*
5. *With the notation on page 302 shew that  $O$  and  $G$  are respectively the external and internal centres of similitude of the circum-scribed and nine-points circle.*
6. *If a variable circle touches two fixed circles, the line joining their points of contact passes through a centre of similitude. Distinguish between the different cases.*
7. *Describe a circle which shall touch two given circles and pass through a given point.*
8. *Describe a circle which shall touch three given circles.*
9.  $C_1, C_2, C_3$  are the centres of three given circles;  $S'_1, S_1$  are the internal and external centres of similitude of the pair of circles whose centres are  $C_1, C_2$ , and  $S'_2, S_2, S'_3, S_3$  have similar meanings with regard to the other two pairs of circles: shew that
  - (i)  $S'_1C_1, S'_2C_2, S'_3C_3$  are concurrent;
  - (ii) the six points  $S_1, S_2, S_3, S'_1, S'_2, S'_3$  lie three and three on four straight lines. [See Ex. 1 and 2, pp. 400, 401.]

## III. ON POLE AND POLAR.

## DEFINITIONS.


1. If in any straight line drawn from the centre of a circle two points are taken such that the rectangle contained by their distances from the centre is equal to the square on the radius, each point is said to be the **inverse** of the other.

Thus in the figure given on the following page, if  $O$  is the centre of the circle, and if  $OP \cdot OQ = (\text{radius})^2$ , then each of the points  $P$  and  $Q$  is the inverse of the other.

It is clear that if one of these points is within the circle the other must be without it.

2. The **polar** of a given point with respect to a given circle is the straight line drawn through the inverse of the given point at right angles to the line which joins the given point to the centre: and with reference to the polar the given point is called the **pole**.

Thus in the adjoining figure, if  $OP \cdot OQ = (\text{radius})^2$ , and if through



$P$  and  $Q$ ,  $LM$  and  $HK$  are drawn perp. to  $OP$ ; then  $HK$  is the polar of the point  $P$ , and  $P$  is the pole of the st. line  $HK$  with respect to the given circle; also  $LM$  is the polar of the point  $Q$ , and  $Q$  the pole of  $LM$ .

It is clear that the polar of an *external* point must intersect the circle, and that the polar of an *internal* point must fall without it: also that the polar of a point *on the circumference* is the tangent at that point.


1. Now it has been proved [see Ex. 1, page 251] that if from an external point  $P$  two tangents  $PH, PK$  are drawn to a circle, of which  $O$  is the centre, then  $OP$  cuts the chord of contact  $HK$  at right angles at  $Q$ , so that

$$OP \cdot OQ = (\text{radius})^2;$$

$\therefore HK$  is the polar of  $P$  with respect to the circle. Def. 2.

Hence we conclude that

*The polar of an external point with reference to a circle is the chord of contact of tangents drawn from the given point to the circle.*



2. If  $A$  and  $P$  are any two points, and if the polar of  $A$  with respect to any circle passes through  $P$ , then the polar of  $P$  must pass through  $A$ .

Let  $BC$  be the polar of the point  $A$  with respect to a circle whose centre is  $O$ , and let  $BC$  pass through  $P$ .

Then shall the polar of  $P$  pass through  $A$ .

Join  $OP$ ; and from  $A$  draw  $AQ$  perp. to  $OP$ . We shall shew that  $AQ$  is the polar of  $P$ .

Now since  $BC$  is the polar of  $A$ ,

$\therefore$  the  $\angle ABP$  is a rt. angle;

*Def. 2, page 391.*

and the  $\angle AQP$  is a rt. angle: *Constr.*

$\therefore$  the four points  $A, B, P, Q$  are concyclic;

$\therefore OQ \cdot OP = OA \cdot OB$  III. 36.

$= (\text{radius})^2$ , for  $CB$  is the polar of  $A$ :


$\therefore P$  and  $Q$  are inverse points with respect to the given circle.

And since  $AQ$  is perp. to  $OP$ ,

$\therefore AQ$  is the polar of  $P$ .

That is, the polar of  $P$  passes through  $A$ .

Q. E. D.



NOTE. A similar proof applies to the case when the given point  $A$  is without the circle, and the polar  $BC$  cuts it.

The above Theorem is known as the **Reciprocal Property of Pole and Polar**.

3. To prove that the locus of the intersection of tangents drawn to a circle at the extremities of all chords which pass through a given point within the circle is the polar of that point.


Let  $A$  be the given point within the circle. Let  $HK$  be any chord passing through  $A$ ; and let the tangents at  $H$  and  $K$  intersect at  $P$ .

It is required to prove that the locus of  $P$  is the polar of the point  $A$ .

I. To shew that  $P$  lies on the polar of  $A$ .

Since  $HK$  is the chord of contact of tangents drawn from  $P$ ,

$\therefore HK$  is the polar of  $P$ . Ex. 1, p. 391.



But HK, the polar of P, passes through A ;

$\therefore$  the polar of A passes through P: Ex. 2, p. 392.

that is, the point P lies on the polar of A.

II. To shew that *any* point on the polar of A satisfies the given conditions.

Let BC be the polar of A, and let P be any point on it.

Draw tangents PH, PK, and let HK be the chord of contact.

Now from Ex. 1, p. 391, we know that the chord of contact HK is the polar of P,

and we also know that the polar of P must pass through A ; for P is on BC, the polar of A: Ex. 2, p. 392.

that is, HK passes through A.

$\therefore$  P is the point of intersection of tangents drawn at the extremities of a chord passing through A.

From I. and II. we conclude that the required locus is the polar of A.

NOTE. If A is *without* the circle, the theorem demonstrated in Part I. of the above proof still holds good ; but the converse theorem in Part II. is not true for *all* points in BC. For if A is without the circle, the polar BC will intersect it ; and no point on that part of the polar which is within the circle can be the point of intersection of tangents.

We now see that

(i) *The Polar of an external point with respect to a circle is the chord of contact of tangents drawn from it.*

(ii) *The Polar of an internal point is the locus of the intersections of tangents drawn at the extremities of all chords which pass through it.*

(iii) *The Polar of a point on the circumference is the tangent at that point.*

The following theorem is known as the Harmonic Property of Pole and Polar.


4. *Any straight line drawn through a point is cut harmonically by the point, its polar, and the circumference of the circle.*

Let  $AHB$  be a circle,  $P$  the given point and  $HK$  its polar; let  $Paqb$  be any straight line drawn through  $P$  meeting the polar at  $q$  and the  $\odot^{ce}$  of the circle at  $a$  and  $b$ .

Then shall  $P, a, q, b$  be a harmonic range.

In the case here considered,  $P$  is an external point.

Join  $P$  to the centre  $O$ , and let  $PO$  cut the  $\odot^{ce}$  at  $A$  and  $B$ : let the polar of  $P$  cut the  $\odot^{ce}$  at  $H$  and  $K$ , and  $PO$  at  $Q$ .



Join  $Qa, Qb, Oa, Ob, PH$ .

Then  $PH$  is a tangent to the  $\odot AHB$ . Ex. 1, p. 391.

From the similar triangles  $OPH, HPQ,$   
 $OP : PH = PH : PQ.$

$$\therefore PQ \cdot PO = PH^2 \\ = Pa \cdot Pb.$$

$\therefore$  the points  $O, Q, a, b$  are concyclic:

$$\therefore \begin{aligned} \angle aQA &= \angle abO && \text{Ex. 5, p. 241.} \\ &= \angle Oab && \text{I. 5.} \\ &= \angle OQb, \text{ in the same segment.} \end{aligned}$$

And since  $QH$  is perp. to  $AB$ ,

$$\therefore \angle aQH = \angle bQH.$$

$\therefore Qq$  and  $QP$  are the internal and external bisectors of the  $\angle aQb$ :

$$\therefore P, a, q, b \text{ is a harmonic range. Ex. 1, p. 385.}$$

The student should investigate for himself the case when  $P$  is an internal point.

*Conversely, it may be shewn that if through a fixed point  $P$  any secant is drawn cutting the circumference of a given circle at  $a$  and  $b$ , and if  $q$  is the harmonic conjugate of  $P$  with respect to  $a, b$ ; then the locus of  $q$  is the polar of  $P$  with respect to the given circle.*

#### DEFINITION.

A triangle so related to a circle that each side is the polar of the opposite vertex is said to be **self-conjugate** with respect to the circle.



## EXAMPLES ON POLE AND POLAR.

1. *The straight line which joins any two points is the polar with respect to a given circle of the point of intersection of their polars.*
2. *The point of intersection of any two straight lines is the pole of the straight line which joins their poles.*
3. *Find the locus of the poles of all straight lines which pass through a given point.*
4. *Find the locus of the poles, with respect to a given circle, of tangents drawn to a concentric circle.*
5. *If two circles cut one another orthogonally and  $PQ$  be any diameter of one of them; shew that the polar of  $P$  with regard to the other circle passes through  $Q$ .*
6. *If two circles cut one another orthogonally, the centre of each circle is the pole of their common chord with respect to the other circle.*
7. *Any two points subtend at the centre of a circle an angle equal to one of the angles formed by the polars of the given points.*
8.  *$O$  is the centre of a given circle, and  $AB$  a fixed straight line.  $P$  is any point in  $AB$ ; find the locus of the point inverse to  $P$  with respect to the circle.*
9. *Given a circle, and a fixed point  $O$  on its circumference:  $P$  is any point on the circle. find the locus of the point inverse to  $P$  with respect to any circle whose centre is  $O$ .*
10. *Given two points  $A$  and  $B$ , and a circle whose centre is  $O$ ; shew that the rectangle contained by  $OA$  and the perpendicular from  $B$  on the polar of  $A$  is equal to the rectangle contained by  $OB$  and the perpendicular from  $A$  on the polar of  $B$ .*
11. *Four points  $A, B, C, D$  are taken in order on the circumference of a circle;  $DA, CB$  intersect at  $P$ ,  $AC, BD$  at  $Q$ , and  $BA, CD$  in  $R$ : shew that the triangle  $PQR$  is self-conjugate with respect to the circle.*
12. *Give a linear construction for finding the polar of a given point with respect to a given circle. Hence find a linear construction for drawing a tangent to a circle from an external point.*
13. *If a triangle is self-conjugate with respect to a circle, the centre of the circle is at the orthocentre of the triangle.*
14. *The polars, with respect to a given circle, of the four points of a harmonic range form a harmonic pencil: and conversely.*



## IV. ON THE RADICAL AXIS.

1. To find the locus of points from which the tangents drawn to two given circles are equal.



Let  $A$  and  $B$  be the centres of the given circles, whose radii are  $a$  and  $b$ ; and let  $P$  be any point such that the tangent  $PQ$  drawn to the circle ( $A$ ) is equal to the tangent  $PR$  drawn to the circle ( $B$ ).

*It is required to find the locus of  $P$ .*

Join  $PA$ ,  $PB$ ,  $AQ$ ,  $BR$ ,  $AB$ ; and from  $P$  draw  $PS$  perp. to  $AB$ .

Then because  $PQ = PR$ ,  $\therefore PQ^2 = PR^2$ .

But  $PQ^2 = PA^2 - AQ^2$ ; and  $PR^2 = PB^2 - BR^2$ ; I. 47.

$\therefore PA^2 - AQ^2 = PB^2 - BR^2$ ;

that is,  $PS^2 + AS^2 - a^2 = PS^2 + SB^2 - b^2$ ; I. 47.

or,  $AS^2 - a^2 = SB^2 - b^2$ .

Hence  $AB$  is divided at  $S$ , so that  $AS^2 - SB^2 = a^2 - b^2$ :

$\therefore S$  is a *fixed point*.

Hence all points from which equal tangents can be drawn to the two circles lie on the straight line which cuts  $AB$  at rt. angles, so that the difference of the squares on the segments of  $AB$  is equal to the difference of the squares on the radii.


Again, by simply retracing these steps, it may be shewn that in Fig. 1 every point in  $SP$ , and in Fig. 2 every point in  $SP$  exterior to the circles, is such that tangents drawn from it to the two circles are equal.

Hence we conclude that in Fig. 1 the whole line  $SP$  is the required locus, and in Fig. 2 that part of  $SP$  which is without the circles.

In either case  $SP$  is said to be the **Radical Axis** of the two circles.

**COROLLARY.** *If the circles cut one another as in Fig. 2, it is clear that the Radical Axis is identical with the straight line which passes through the points of intersection of the circles; for it follows readily from III. 36 that tangents drawn to two intersecting circles from any point in the common chord produced are equal.*

2. *The Radical Axes of three circles taken in pairs are concurrent.*



Let there be three circles whose centres are A, B, C.

Let OZ be the radical axis of the  $\odot^s$  (A) and (B);  
and OY the Radical Axis of the  $\odot^s$  (A) and (C), O being the point  
of their intersection.

*Then shall the radical axis of the  $\odot^s$  (B) and (C) pass through O.*

It will be found that the point O is either *without* or *within* all  
the circles.

I. When O is without the circles.

From O draw OP, OQ, OR tangents to the  $\odot^s$  (A), (B), (C).

Then because O is a point on the radical axis of (A) and (B); *Hyp.*

$$\therefore OP = OQ.$$

And because O is a point on the radical axis of (A) and (C), *Hyp.*

$$\therefore OP = OR;$$

$$\therefore OQ = OR;$$

$\therefore$  O is a point on the radical axis of (B) and (C);


that is, the radical axis of (B) and (C) passes through O.

II. If the circles intersect in such a way that O is within them  
all;

the radical axes are then the common chords of the three circles  
taken two and two; and it is required to prove that these common  
chords are concurrent. This may be shewn indirectly by III. 35.

**DEFINITION.** The point of intersection of the radical axes  
of three circles taken in pairs is called the **radical centre**.

3. To draw the radical axis of two given circles.



Let *A* and *B* be the centres of the given circles.

*It is required to draw their radical axis.*

If the given circles intersect, then the st. line drawn through their points of intersection will be the radical axis. [Ex. 1, Cor. p. 397.]

But if the given circles do not intersect, describe any circle so as to cut them in *E*, *F* and *G*, *H*.

Join *EF* and *HG*, and produce them to meet in *P*.

Join *AB*; and from *P* draw *PS* perp. to *AB*.

*Then PS shall be the radical axis of the  $\odot^s$  (*A*), (*B*).*

[The proof follows from III. 36 and Ex. 1, p. 396.]

**DEFINITION.** If each pair of circles in a given system have the same radical axis, the circles are said to be **co-axal**.

#### EXAMPLES ON THE RADICAL AXIS.

1. Shew that the radical axis of two circles bisects any one of their common tangents.

2. If tangents are drawn to two circles from any point on their radical axis; shew that a circle described with this point as centre and any one of the tangents as radius, cuts both the given circles orthogonally.

3. *O* is the radical centre of three circles, and from *O* a tangent *OT* is drawn to any one of them: shew that a circle whose centre is *O* and radius *OT* cuts all the given circles orthogonally.

4. If three circles touch one another, taken two and two, shew that their common tangents at the points of contact are concurrent.

5. *If circles are described on the three sides of a triangle as diameter, their radical centre is the orthocentre of the triangle.*

6. *All circles which pass through a fixed point and cut a given circle orthogonally, pass through a second fixed point.*

7. *Find the locus of the centres of all circles which pass through a given point and cut a given circle orthogonally.*

8. *Describe a circle to pass through two given points and cut a given circle orthogonally.*

9. *Find the locus of the centres of all circles which cut two given circles orthogonally.*

10. *Describe a circle to pass through a given point and cut two given circles orthogonally.*

11. *The difference of the squares on the tangents drawn from any point to two circles is equal to twice the rectangle contained by the straight line joining their centres and the perpendicular from the given point on their radical axis.*

12. *In a system of co-axial circles which do not intersect, any point is taken on the radical axis; shew that a circle described from this point as centre, with radius equal to the tangent drawn from it to any one of the circles, will meet the line of centres in two fixed points.*

[These fixed points are called the **Limiting Points** of the system.]

13. *In a system of co-axial circles the two limiting points and the points in which any one circle of the system cuts the line of centres form a harmonic range.*

14. *In a system of co-axial circles a limiting point has the same polar with regard to all the circles of the system.*


15. *If two circles are orthogonal any diameter of one is cut harmonically by the other.*

## V. ON TRANSVERSALS.

In the two following theorems we are to suppose that the segments of straight lines are expressed numerically in terms of some common unit; and the ratio of one such segment to another will be denoted by the fraction of which the first is the numerator and the second the denominator.

**DEFINITION.** A straight line drawn to cut a given system of lines is called a **transversal**.

1. *If three concurrent straight lines are drawn from the angular points of a triangle to meet the opposite sides, then the product of three alternate segments taken in order is equal to the product of the other three segments.*



Let  $AD, BE, CF$  be drawn from the vertices of the  $\triangle ABC$  to intersect at  $O$ , and cut the opposite sides at  $D, E, F$ .

Then shall  $BD \cdot CE \cdot AF = DC \cdot EA \cdot FB$ .

Now the  $\triangle AOB, AOC$  have a common base  $AO$ ; and it may be shewn that

$BD : DC =$  the alt. of  $\triangle AOB$  : the alt. of  $\triangle AOC$  ;

$$\therefore \frac{BD}{DC} = \frac{\triangle AOB}{\triangle AOC} ;$$

similarly,

$$\frac{CE}{EA} = \frac{\triangle BOC}{\triangle BOA} ;$$

and

$$\frac{AF}{FB} = \frac{\triangle COA}{\triangle COB} .$$

Multiplying these ratios, we have

$$\frac{BD}{DC} \cdot \frac{CE}{EA} \cdot \frac{AF}{FB} = 1 ;$$

or,

$$BD \cdot CE \cdot AF = DC \cdot EA \cdot FB .$$


Q. E. D.

**NOTE.** The converse of this theorem, which may be proved indirectly, is very important : it may be enunciated thus :

*If three straight lines drawn from the vertices of a triangle cut the opposite sides so that the product of three alternate segments taken in order is equal to the product of the other three, then the three straight lines are concurrent.*

*That is, if  $BD \cdot CE \cdot AF = DC \cdot EA \cdot FB$ ,  
then  $AD, BE, CF$  are concurrent.*

2. If a transversal is drawn to cut the sides, or the sides produced, of a triangle, the product of three alternate segments taken in order is equal to the product of the other three segments.



Let ABC be a triangle, and let a transversal meet the sides BC, CA, AB, or these sides produced, at D, E, F.

Then shall  $BD \cdot CE \cdot AF = DC \cdot EA \cdot FB$ .

Draw AH par<sup>l</sup> to BC, meeting the transversal at H.

Then from the similar  $\triangle^s$  DFB, HAF,

$$\frac{BD}{FB} = \frac{HA}{AF};$$

and from the similar  $\triangle^s$  DCE, HAE,

$$\frac{CE}{DC} = \frac{EA}{HA};$$

$\therefore$ , by multiplication,  $\frac{BD}{FB} \cdot \frac{CE}{DC} = \frac{EA}{AF};$

that is,  $\frac{BD \cdot CE \cdot AF}{DC \cdot EA \cdot FB} = 1,$

or,  $BD \cdot CE \cdot AF = DC \cdot EA \cdot FB.$

Q. E. D.

NOTE. In this theorem the transversal must either meet two sides and the third side produced, as in Fig. 1; or all three sides produced, as in Fig. 2.

The converse of this theorem may be proved indirectly:

If three points are taken in two sides of a triangle and the third side produced, or in all three sides produced, so that the product of three alternate segments taken in order is equal to the product of the other three segments, the three points are collinear.

## DEFINITIONS.

1. If two triangles are such that three straight lines joining corresponding vertices are concurrent, they are said to be **co-polar**.

2. If two triangles are such that the points of intersection of corresponding sides are collinear, they are said to be **co-axial**.

The propositions given on pages 111-114 relating to the concurrence of straight lines in a triangle, may be proved by the method of transversals, and in addition to these the following important theorems may be established.

## THEOREMS TO BE PROVED BY TRANSVERSALS.

1. *The straight lines which join the vertices of a triangle to the points of contact of the inscribed circle (or any of the three escribed circles) are concurrent.*

2. *The middle points of the diagonals of a complete quadrilateral are collinear.* [See Def. 4, p. 387.]

3. *Co-polar triangles are also co-axial; and conversely co-axial triangles are also co-polar.*

4. *The six centres of similitude of three circles lie three by three on four straight lines.*

MISCELLANEOUS EXAMPLES ON BOOK VI.

1. Through  $D$ , any point in the base of a triangle  $ABC$ , straight lines  $DE$ ,  $DF$  are drawn parallel to the sides  $AB$ ,  $AC$ , and meeting the sides at  $E$ ,  $F$  : shew that the triangle  $AEF$  is a mean proportional between the triangles  $FBD$ ,  $EDC$ .

2. If two triangles have one angle of the one equal to one angle of the other, and a second angle of the one supplementary to a second angle of the other, then the sides about the third angles are proportional.

3.  $AE$  bisects the vertical angle of the triangle  $ABC$  and meets the base in  $E$  ; shew that if circles are described about the triangles  $ABE$ ,  $ACE$ , the diameters of these circles are to each other in the same ratio as the segments of the base.

4. Through a fixed point  $O$  draw a straight line so that the parts intercepted between  $O$  and the perpendiculars drawn to the straight line from two other fixed points may have a given ratio.

5. The angle  $A$  of a triangle  $ABC$  is bisected by  $AD$  meeting  $BC$  in  $D$ , and  $AX$  is the median bisecting  $BC$  : shew that  $XD$  has the same ratio to  $XB$  as the difference of the sides has to their sum.

6.  $AD$  and  $AE$  bisect the vertical angle of a triangle internally and externally, meeting the base in  $D$  and  $E$  ; shew that if  $O$  is the middle point of  $BC$ , then  $OB$  is a mean proportional between  $OD$  and  $OE$ .

7.  $P$  and  $Q$  are fixed points ;  $AB$  and  $CD$  are fixed parallel straight lines ; any straight line is drawn from  $P$  to meet  $AB$  at  $M$ , and a straight line is drawn from  $Q$  parallel to  $PM$  meeting  $CD$  at  $N$  : shew that the ratio of  $PM$  to  $QN$  is constant, and thence shew that the straight line through  $M$  and  $N$  passes through a fixed point.

8. If  $C$  is the middle point of an arc of a circle whose chord is  $AB$ , and  $D$  is any point in the conjugate arc ; shew that  

$$AD + DB : DC :: AB : AC.$$

9. In the triangle  $ABC$  the side  $AC$  is double of  $BC$ . If  $CD$ ,  $CE$  bisect the angle  $ACB$  internally and externally meeting  $AB$  in  $D$  and  $E$ , shew that the areas of the triangles  $CBD$ ,  $ACD$ ,  $ABC$ ,  $CDE$  are as 1, 2, 3, 4.

10.  $AB$ ,  $AC$  are two chords of a circle ; a line parallel to the tangent at  $A$  cuts  $AB$ ,  $AC$  in  $D$  and  $E$  respectively : shew that the rectangle  $AB$ ,  $AD$  is equal to the rectangle  $AC$ ,  $AE$ .



11. If from any point on the hypotenuse of a right-angled triangle perpendiculars are drawn to the two sides, the rectangle contained by the segments of the hypotenuse will be equal to the sum of the rectangles contained by the segments of the sides.

12. D is a point in the side AC of the triangle ABC, and E is a point in AB. If BD, CE divide each other into parts in the ratio 4 : 1, then D, E divide CA, BA in the ratio 3 : 1.

13. If the perpendiculars from two fixed points on a straight line passing between them be in a given ratio, the straight line must pass through a third fixed point.

14. PA, PB are two tangents to a circle; PCD any chord through P: shew that the rectangle contained by one pair of opposite sides of the quadrilateral ACBD is equal to the rectangle contained by the other pair.

15. A, B, C are any three points on a circle, and the tangent at A meets BC produced in D: shew that the diameters of the circles circumscribed about ABD, ACD are as AD to CD.

16. AB, CD are two diameters of the circle AD BC at right angles to each other, and EF is any chord; CE, CF are drawn meeting AB produced in G and H; prove that  
the rect. CE, HG = the rect. EF, CH.

17. From the vertex A of any triangle ABC draw a line meeting BC produced in D so that AD may be a mean proportional between the segments of the base.

18. Two circles touch internally at O; AB a chord of the larger circle touches the smaller in C which is cut by the lines OA, OB in the points P, Q: shew that  $OP : OQ :: AC : CB$ .

19. AB is any chord of a circle; AC, BC are drawn to any point C in the circumference and meet the diameter perpendicular to AB at D, E: if O is the centre, shew that the rect. OD, OE is equal to the square on the radius.

20. YD is a tangent to a circle drawn from a point Y in the diameter AB produced; from D a perpendicular DX is drawn to the diameter; shew that the points X, Y divide AB internally and externally in the same ratio.

21. Determine a point in the circumference of a circle, from which lines drawn to two other given points shall have a given ratio.

22.  $O$  is the centre and  $OA$  a radius of a given circle, and  $V$  is a fixed point in  $OA$ ;  $P$  and  $Q$  are two points on the circumference on opposite sides of  $A$  and equidistant from it;  $QV$  is produced to meet the circle in  $L$ ; shew that, whatever be the length of the arc  $PQ$ , the chord  $LP$  will always meet  $OA$  produced in a fixed point.

23.  $EA, EA'$  are diameters of two circles touching each other externally at  $E$ ; a chord  $AB$  of the former circle, when produced, touches the latter at  $C'$ , while a chord  $A'B'$  of the latter touches the former at  $C$ : prove that the rectangle, contained by  $AB$  and  $A'B'$ , is four times as great as that contained by  $BC'$  and  $B'C$ .

24. If a circle be described touching externally two given circles, the straight line passing through the points of contact will intersect the line of centres of the given circles at a fixed point.

25. Two circles touch externally in  $C$ ; if any point  $D$  be taken without them so that the radii  $AC, BC$  subtend equal angles at  $D$ , and  $DE, DF$  be tangents to the circles, shew that  $DC$  is a mean proportional between  $DE$  and  $DF$ .

26. If through the middle point of the base of a triangle any line be drawn intersecting one side of the triangle, the other produced, and the line drawn parallel to the base from the vertex, it will be divided harmonically.

27. If from either base angle of a triangle a line be drawn intersecting the median from the vertex, the opposite side, and the line drawn parallel to the base from the vertex, it will be divided harmonically.

28. Any straight line drawn to cut the arms of an angle and its internal and external bisectors is cut harmonically.

29.  $P, Q$  are harmonic conjugates of  $A$  and  $B$ , and  $C$  is an external point; if the angle  $PCQ$  is a right angle, shew that  $CP, CQ$  are the internal and external bisectors of the angle  $ACB$ .

30. From  $C$ , one of the base angles of a triangle, draw a straight line meeting  $AB$  in  $G$ , and a straight line through  $A$  parallel to the base in  $E$ , so that  $CE$  may be to  $EG$  in a given ratio.

31.  $P$  is a given point outside the angle formed by two given lines  $AB, AC$ ; shew how to draw a straight line from  $P$  such that the parts of it intercepted between  $P$  and the lines  $AB, AC$  may have a given ratio.

32. Through a given point within a given circle, draw a straight line such that the parts of it intercepted between that point and the circumference may have a given ratio. How many solutions does the problem admit of?

33. If a common tangent be drawn to any number of circles which touch each other internally, and from any point of this tangent as a centre a circle be described, cutting the other circles; and if from this centre lines be drawn through the intersections of the circles, the segments of the lines within each circle shall be equal.

34. APB is a quadrant of a circle, SPT a line touching it at P; C is the centre, and PM is perpendicular to CA; prove that

$$\text{the } \triangle SCT : \text{the } \triangle ACB :: \text{the } \triangle ACB : \text{the } \triangle CMP,$$

35. ABC is a triangle inscribed in a circle, AD, AE are lines drawn to the base BC parallel to the tangents at B, C respectively; shew that AD=AE, and  $BD : CE :: AB^2 : AC^2$ .

36. AB is the diameter of a circle, E the middle point of the radius OB; on AE, EB as diameters circles are described; PQL is a common tangent touching the circles at P and Q, and AB produced at L: shew that BL is equal to the radius of the smaller circle.

37. The vertical angle C of a triangle is bisected by a straight line which meets the base at D, and is produced to a point E, such that the rectangle contained by CD and CE is equal to the rectangle contained by AC and CB: shew that if the base and vertical angle be given, the position of E is invariable.

38. ABC is an isosceles triangle having the base angles at B and C each double of the vertical angle: if BE and CD bisect the base angles and meet the opposite sides in E and D, shew that DE divides the triangle into figures whose ratio is equal to that of AB to BC.

39. If AB, the diameter of a semicircle, be bisected in C, and on AC and CB circles be described, and in the space between the three circumferences a circle be inscribed, shew that its diameter will be to that of the equal circles in the ratio of 2 to 3.

40. O is the centre of a circle inscribed in a quadrilateral ABCD; a line EOF is drawn and making equal angles with AD and BC, and meeting them in E and F respectively: shew that the triangles AEO, BOF are similar, and that

$$AE : ED = CF : FB.$$

41. From the last exercise deduce the following : The inscribed circle of a triangle  $ABC$  touches  $AB$  in  $F$  ;  $XOY$  is drawn through the centre making equal angles with  $AB$  and  $AC$ , and meeting them in  $X$  and  $Y$  respectively : shew that  $BX : XF = AY : YC$ .

42. Inscribe a square in a given semicircle.

43. Inscribe a square in a given segment of a circle.

44. Describe an equilateral triangle equal to a given isosceles triangle.

45. Describe a square having given the difference between a diagonal and a side.

46. Given the vertical angle, the ratio of the sides containing it, and the diameter of the circumscribing circle, construct the triangle.

47. Given the vertical angle, the line bisecting the base, and the angle the bisector makes with the base, construct the triangle.

48. In a given circle inscribe a triangle so that two sides may pass through two given points and the third side be parallel to a given straight line.

49. In a given circle inscribe a triangle so that the sides may pass through the three given points.

50.  $A, B, X, Y$  are four points in a straight line, and  $O$  is such a point in it that the rectangle  $OA, OY$  is equal to the rectangle  $OB, OX$  ; if a circle is described with centre  $O$  and radius equal to a mean proportional between  $OA$  and  $OY$ , shew that at every point on this circle  $AB$  and  $XY$  will subtend equal angles.

51.  $O$  is a fixed point, and  $OP$  is any line drawn to meet a fixed straight line in  $P$  ; if on  $OP$  a point  $Q$  is taken so that  $OQ$  to  $OP$  is a constant ratio, find the locus of  $Q$ .

52.  $O$  is a fixed point, and  $OP$  is any line drawn to meet the circumference of a fixed circle in  $P$  ; if on  $OP$  a point  $Q$  is taken so that  $OQ$  to  $OP$  is a constant ratio, find the locus of  $Q$ .

53. If from a given point two straight lines are drawn including a given angle, and having a fixed ratio, find the locus of the extremity of one of them when the extremity of the other lies on a fixed straight line.

54. On a straight line  $PAB$ , two points  $A$  and  $B$  are marked and the line  $PAB$  is made to revolve round the fixed extremity  $P$ .  $C$  is a fixed point in the plane in which  $PAB$  revolves ; prove that if  $CA$  and  $CB$  be joined and the parallelogram  $CADB$  be completed, the locus of  $D$  will be a circle.

55. Find the locus of a point whose distances from two fixed points are in a given ratio.

56. Find the locus of a point from which two given circles subtend the same angle.

57. Find the locus of a point such that its distances from two intersecting straight lines are in a given ratio.

58. In the figure on page 389, shew that  $QT$ ,  $P'T'$  meet on the radical axis of the two circles.

59.  $ABC$  is any triangle, and on its sides equilateral triangles are described externally: if  $X, Y, Z$  are the centres of their inscribed circles, shew that the triangle  $XYZ$  is equilateral.

60. If  $S, I$  are the centres, and  $R, r$  the radii of the circumscribed and inscribed circles of a triangle, and if  $N$  is the centre of its nine-points circle,

$$\text{prove that} \quad \begin{aligned} \text{(i)} \quad SI^2 &= R^2 - 2Rr, \\ \text{(ii)} \quad NI &= \frac{1}{2}R - r. \end{aligned}$$

Establish corresponding properties for the escribed circles, and hence prove that the nine-points circle touches the inscribed and escribed circles of a triangle.

# SOLID GEOMETRY.

## EUCLID. BOOK XI.

### DEFINITIONS.

FROM the Definitions of Book I. it will be remembered that

(i) A **line** is that which has *length*, without breadth or thickness.

(ii) A **surface** is that which has *length* and *breadth*, without thickness.

To these definitions we have now to add :

(iii) **Space** is that which has *length*, *breadth*, and *thickness*.


Thus a line is said to be of **one dimension** ;  
a surface is said to be of **two dimensions** ;  
and space is said to be of **three dimensions**.

The Propositions of Euclid's Eleventh Book here given establish the first principles of the *geometry of space*, or *solid geometry*. They deal with the properties of straight lines which are not all in the same plane, the relations which straight lines bear to planes which do not contain those lines, and the relations which two or more planes bear to one another. Unless the contrary is stated the straight lines are supposed to be of indefinite length, and the planes of infinite extent.

Solid geometry then proceeds to discuss the properties of solid figures, of surfaces which are not planes, and of lines which cannot be drawn on a plane surface.

## LINES AND PLANES.

1. A straight line is **perpendicular to a plane** when it is perpendicular to *every* straight line which meets it in that plane.




**NOTE.** It will be proved in Proposition 4 that if a straight line is perpendicular to *two* straight lines which meet it in a plane, it is also perpendicular to *every* straight line which meets it in that plane.

A straight line drawn perpendicular to a plane is said to be a **normal** to that plane.

2. The foot of the perpendicular let fall from a given point on a plane is called the **projection of that point** on the plane.


3. The **projection of a line** on a plane is the locus of the feet of perpendiculars drawn from all points in the given line to the plane.



Thus in the above figure the line  $ab$  is the projection of the line  $AB$  on the plane  $PQ$ .

**NOTE.** It will be proved hereafter (see page 446) that the projection of a straight line on a plane is also a straight line.

4. The **inclination of a straight line to a plane** is the acute angle contained by that line and another drawn from the point at which the first line meets the plane to the point at which a perpendicular to the plane let fall from any point of the first line meets the plane.




Thus in the above figure, if from any point  $X$  in the given straight line  $AB$ , which intersects the plane  $PQ$  at  $A$ , a perpendicular  $Xx$  is let fall on the plane, and the straight line  $Ax$  is drawn from  $A$  through  $x$ , then the inclination of the straight line  $AB$  to the plane  $PQ$  is measured by the acute angle  $BAx$ . In other words:—

The **inclination of a straight line to a plane** is the acute angle contained by the given straight line and its *projection on the plane*.

**AXIOM.** If two surfaces intersect one another, they meet in a *line or lines*.

5. The **common section** of two intersecting surfaces is the line (or lines) in which they meet.




**NOTE.** It is proved in Proposition 3 that the common section of two planes is a straight line.

Thus  $AB$ , the common section of the two planes  $PQ$ ,  $XY$  is proved to be a straight line.




6. One plane is perpendicular to another plane when any straight line drawn in one of the planes perpendicular to the common section is also perpendicular to the other plane.



Thus in the above figure, the plane EB is perpendicular to the plane CD, if any straight line PQ, drawn in the plane EB at right angles to the common section AB, is also at right angles to the plane CD.

7. The inclination of a plane to a plane is the acute angle contained by two straight lines drawn from any point in the common section at right angles to it, one in one plane and one in the other.

Thus in the adjoining figure, the straight line AB is the common section of the two intersecting planes BC, AD; and from Q, any point in AB, two straight lines QP, QR are drawn perpendicular to AB, one in each plane: then the inclination of the two planes is measured by the acute angle PQR.



NOTE. This definition assumes that the angle PQR is of constant magnitude whatever point Q is taken in AB: the truth of which assumption is proved in Proposition 10.

The angle formed by the intersection of two planes is called a **dihedral angle**.


It may be proved that two planes are perpendicular to one another when the dihedral angle formed by them is a right angle.

8. **Parallel planes** are such as do not meet when produced.

9. A straight line is **parallel to a plane** if it does not meet the plane when produced.

10. The angle between two straight lines which do not meet is the angle contained by two *intersecting* straight lines respectively parallel to the two non-intersecting lines.


Thus if  $AB$  and  $CD$  are two straight lines which do not meet, and  $ab$ ,  $bc$  are two intersecting lines parallel respectively to  $AB$  and  $CD$ ; then the angle between  $AB$  and  $CD$  is measured by the angle  $abc$ .



11. A **solid angle** is that which is made by three or more plane angles which have a common vertex, but are not in the same plane.

A solid angle made by *three* plane angles is said to be **trihedral**; if made by more than three, it is said to be **polyhedral**.

A solid angle is sometimes called a **corner**.



12. A **solid figure** is any portion of space bounded by one or more surfaces, plane or curved.


These surfaces are called the **faces** of the solid, and the intersections of adjacent faces are called **edges**.

## POLYHEDRA.

13. A **polyhedron** is a solid figure bounded by plane faces.

**NOTE.** A plane rectilineal figure must at least have *three* sides; or *four*, if two of the sides are parallel. A polyhedron must at least have *four* faces; or, if two faces are parallel, it must at least have *five* faces.

14. A **prism** is a solid figure bounded by plane faces, of which two that are opposite are similar and equal polygons in parallel planes, and the other faces are parallelograms.



The polygons are called the **ends** of the prism. A prism is said to be **right** if the edges formed by each pair of adjacent parallelograms are perpendicular to the two ends; if otherwise the prism is **oblique**.

15. A **parallelepiped** is a solid figure bounded by three pairs of parallel plane faces.

Fig. 1.






Fig. 2.



A parallelepiped may be *rectangular* as in fig. 1, or *oblique* as in fig. 2. The name **cuboid** is sometimes given to a rectangular parallelepiped whose length, breadth, and thickness are not all equal.


16. A **pyramid** is a solid figure bounded by plane faces, of which one is a polygon, and the rest are triangles having as bases the sides of the polygon, and as a common vertex some point not in the plane of the polygon.



The polygon is called the **base** of the pyramid.

A pyramid having for its base a *regular* polygon is said to be **right** when the vertex lies in the straight line drawn perpendicular to the base from its central point (the centre of its inscribed or circumscribed circle).

17. A **tetrahedron** is a pyramid on a triangular base: it is thus contained by *four* triangular faces.



18. Polyhedra are classified according to the number of their *faces* :

thus a **hexahedron** has *six* faces ;  
 an **octahedron** has *eight* faces ;  
 a **dodecahedron** has *twelve* faces.


19. **Similar polyhedra** are such as have all their solid angles equal, each to each, and are bounded by the same number of similar faces.

20. A polyhedron is **regular** when its faces are similar and equal regular polygons.


21. It will be proved (see page 451) that there can only be *five* regular polyhedra.

They are defined as follows:—


(i) A **regular tetrahedron** is a solid figure bounded by *four* plane faces, which are equal and equilateral *triangles*.




(ii) A **cube** is a solid figure bounded by *six* plane faces, which are equal *squares*.




(iii) A **regular octahedron** is a solid figure bounded by *eight* plane faces, which are equal and equilateral *triangles*.



(iv) A **regular dodecahedron** is a solid figure bounded by *twelve* plane faces, which are equal and regular *pentagons*.



(v) A **regular icosahedron** is a solid figure bounded by *twenty* plane faces, which are equal and equilateral *triangles*.



### SOLIDS OF REVOLUTION.

22a. A **sphere** is a solid figure described by the revolution of a semicircle about its diameter, which remains fixed.

The **axis** of the sphere is the fixed straight line about which the semicircle revolves.

The **centre** of the sphere is the same as the centre of the semicircle.

A **diameter** of a sphere is any straight line which passes through the centre, and is terminated both ways by the surface of the sphere.


The following definition of a sphere, analogous to that given for a circle (I. Def. 15), may also be noted :

22b. A **sphere** is a solid figure contained by one surface, which is such that all straight lines drawn from a certain point within it to the surface are equal : this point is called the centre of the sphere.

A **radius** of a sphere is a straight line drawn from the centre to the surface.

It will be seen that the surface of a sphere is the locus of a point which moves *in space* so that its distance from a certain fixed point (the centre) is constant.

23. A **right cylinder** is a solid figure described by the revolution of a rectangle about one of its sides which remains fixed.



The **axis** of the cylinder is the fixed straight line about which the rectangle revolves.

The **bases, or ends,** of the cylinder are the circular faces described by the two revolving opposite sides of the rectangle.

24. A **right cone** is a solid figure described by the revolution of a right-angled triangle about one of the sides containing the right angle which remains fixed.



The **axis** of the cone is the fixed straight line about which the triangle revolves.


The **base** of the cone is the circular face described by that side which revolves.

The hypotenuse of the right-angled triangle in any one of its positions is called a **generating line** of the cone.

25. **Similar cones and cylinders** are those which have their axes and the diameters of their bases proportionals.

PROPOSITION 1. THEOREM.

*One part of a straight line cannot be in a plane and another part outside it.*



If possible, let AB, part of the st. line ABC, be in the plane PQ, and the part BC outside it.

Then since the st. line AB is in the plane PQ,  
 $\therefore$  it can be produced in that plane. I. Post. 2.

Produce AB to D;  
 and let any other plane which passes through AD be turned about AD until it passes also through C.

Then because the points B and C are in this plane,  
 $\therefore$  the st. line BC is in it: I. Def. 7.  
 $\therefore$  ABC and ABD are in the same plane and are both st. lines; which is impossible. I. Def. 4.  
 $\therefore$  the st. line ABC has not one part AB in the plane PQ, and another part BC outside it. Q.E.D.

**NOTE.** This proposition scarcely needs proof, for the truth of it follows immediately from the definitions of a straight line and a plane.


It should be observed that the method of proof used in this and the next proposition rests upon the following axiom:

*If a plane of unlimited extent turns about a fixed straight line as an axis, it can be made to pass through any point in space.*



## PROPOSITION 2. THEOREM.

*Any two intersecting straight lines are in one plane: and any three straight lines, of which each pair intersect one another, are in one plane.*



Let the two st. lines AB and CD intersect at E ;  
and let the st. line BC be drawn cutting AB and CD at B  
and C.

Then (i) AB and CD shall lie in one plane.  
(ii) AB, BC, CD shall lie in one plane.

(i) Let any plane pass through AB ;  
and let this plane be turned about AB until it passes  
through C.

Then, since C and E are points in this plane,  
∴ the whole st. line CED is in it. I. Def. 7 and XI. 1.  
That is, AB and CD lie in one plane.

(ii) And since B and C are points in the plane which  
contains AB and CD,  
∴ also the st. line BC lies in this plane. Q.E.D.


**COROLLARY.** *One, and only one, plane can be made to pass  
through two given intersecting straight lines.*

Hence the position of a plane is fixed,

- (i) if it passes through a given straight line and a given point  
outside it ; Ax. p. 419.
- (ii) if it passes through two intersecting straight lines ; XI. 2.
- (iii) if it passes through three points not collinear ; XI. 2.
- (iv) if it passes through two parallel straight lines. I. Def. 35.

PROPOSITION 3. THEOREM.

If two planes cut one another, their common section is a straight line.



Let the two planes  $XA$ ,  $CY$  cut one another, and let  $BD$  be their common section.

Then shall  $BD$  be a straight line.

For if not, from  $B$  to  $D$  in the plane  $XA$  draw the st. line  $BED$ ;

and in the plane  $CY$  draw the st. line  $BFD$ .

Then the st. lines  $BED$ ,  $BFD$  have the same extremities;

$\therefore$  they include a space;

but this is impossible.

*Ax.* 10.

$\therefore$  the common section  $BD$  cannot be otherwise than a st. line.

ALTERNATIVE PROOF.

Let the planes  $XA$ ,  $CY$  cut one another, and let  $B$  and  $D$  be two points in their common section.

Then because  $B$  and  $D$  are two points in the plane  $XA$ ,

$\therefore$  the st. line joining  $B$ ,  $D$  lies in that plane. *I. Def.* 7.

And because  $B$  and  $D$  are two points in the plane  $CY$ ,

$\therefore$  the st. line joining  $B$ ,  $D$  lies in that plane.


Hence the st. line  $BD$  lies in *both planes*,

and is therefore their common section.

That is, the common section of the two planes is a *straight line*. Q.E.D.

## PROPOSITION 4. THEOREM. [Alternative Proof.]

If a straight line is perpendicular to each of two straight lines at their point of intersection, it shall also be perpendicular to the plane in which they lie.



Let the straight line AD be perp. to each of the st. lines AB, AC at A their point of intersection.

Then shall AD be perp. to the plane in which AB and AC lie.

Produce DA to F, making AF equal to DA.

Draw any st. line BC in the plane of AB, AC, to cut AB, AC at B and C;  
and in the same plane draw through A any st. line AE to cut BC at E.

It is required to prove that AD is perp. to AE. XI. Def. 1.

Join DB, DE, DC; and FB, FE, FC.

Then in the  $\triangle$  BAD, BAF,

because DA = FA,

*Constr.*

and the common side AB is perp. to DA, FA;

$\therefore$  BD = BF.

I. 4.

Similarly CD = CF.

Now if the  $\triangle$  BFC be turned about its base BC until the vertex F comes into the plane of the  $\triangle$  BDC,

then F will coincide with D,

since the conterminous sides of the triangles are equal. I. 7

$\therefore$  EF will coincide with ED,

that is, EF = ED.

Hence in the  $\triangle^s$  DAE, FAE,  
 since DA, AE, ED = FA, AE, EF respectively,  
 $\therefore$  the  $\angle$  DAE = the  $\angle$  FAE.


That is, DA is perp. to AE.

Similarly it may be shewn that DA is perp. to every  
 st. line which meets it in the plane of AB, AC ;

$\therefore$  DA is perp. to this plane. Q.E.D.

PROPOSITION 4. THEOREM. [Euclid's Proof.]

*If a straight line is perpendicular to each of two straight lines at their point of intersection, it shall also be perpendicular to the plane in which they lie.*



Let the st. line EF be perp. to each of the st. lines  
 AB, DC at E their point of intersection.

*Then shall EF be also perp. to the plane XY, in which AB and  
 DC lie.*

Make EA, EC, EB, ED all equal, and join AD, BC.

Through E in the plane XY draw any st. line cutting  
 AD and BC in G and H.

Take any pt. F in EF; and join FA, FG, FD, FB, FH, FC.

Then in the  $\triangle^s$  AED, BEC,

because AE, ED = BE, EC respectively,


and the  $\angle$  AED = the  $\angle$  BEC ;

$\therefore$  AD = BC, and the  $\angle$  DAE = the  $\angle$  CBE.

*Constr.*

I. 15.

I. 4.



In the  $\triangle^s$  AEG, BEH,  
 because the  $\angle$  GAE = the  $\angle$  HBE,  
 and the  $\angle$  AEG = the  $\angle$  BEH,  
 and EA = EB ; *Proved.*  
I. 15.  
 $\therefore$  EG = EH, and AG = BH. *Constr.*  
I. 26.

Again in the  $\triangle^s$  FEA, FEB,  
 because EA = EB,  
 and the common side FE is perp. to EA, EB ; *Hyp.*  
I. 4.  
 $\therefore$  FA = FB.

Similarly FC = FD,

Again in the  $\triangle^s$  DAF, CBF,  
 because DA, AF, FD = CB, BF, FC, respectively,  
 $\therefore$  the  $\angle$  DAF = the  $\angle$  CBF. I. 8.

And in the  $\triangle^s$  FAG, FBH,  
 because FA, AG = FB, BH, respectively,  
 and the  $\angle$  FAG = the  $\angle$  FBH, *Proved.*  
I. 4.  
 $\therefore$  FG = FH.


Lastly in the  $\triangle^s$  FEG, FEH,  
 because FE, EG, GF = FE, EH, HF, respectively,  
 $\therefore$  the  $\angle$  FEG = the  $\angle$  FEH ; I. 8.  
 that is, FE is perp. to GH.

Similarly it may be shewn that FE is perp. to every  
 st. line which meets it in the plane XY,

$\therefore$  FE is perp. to this plane. XI. *Def.* 1.  
Q.E.D.

PROPOSITION 5. THEOREM.

*If a straight line is perpendicular to each of three concurrent straight lines at their point of intersection, these three straight lines shall be in one plane.*



Let the straight line AB be perpendicular to each of the straight lines BC, BD, BE, at B their point of intersection.

*Then shall BC, BD, BE be in one plane.*

Let XY be the plane which passes through BE, BD; XI. 2.  
and, if possible, suppose that BC is not in this plane.

Let AF be the plane which passes through AB, BC;  
and let the common section of the two planes XY, AF be the st. line BF. XI. 3.

Then since AB is perp. to BE and BD,  
 $\therefore$  AB is perp. to the plane containing BE, BD, namely the plane XY; XI. 4.

and since BF is in this plane,  
 $\therefore$  AB is also perp. to BF. XI. Def. 1.


But AB is perp. to BC; *Hyp.*  
 $\therefore$  the  $\angle^s$  ABF, ABC, which are in the same plane AF, are both rt. angles; which is impossible.

$\therefore$  BC is not outside the plane of BD, BE :  
that is, BC, BD, BE are in one plane.

Q.E.D.

## PROPOSITION 6. THEOREM.

*If two straight lines are perpendicular to the same plane, they shall be parallel to one another.*



Let the st. lines  $AB, CD$  be perp. to the plane  $XY$ .

*Then shall  $AB$  and  $CD$  be par<sup>l</sup>.\**

Let  $AB$  and  $CD$  meet the plane  $XY$  at  $B$  and  $D$ .

Join  $BD$ ;

and in the plane  $XY$  draw  $DE$  perp. to  $BD$ , making  $DE$  equal to  $AB$ .

Join  $BE, AE, AD$ .

Then since  $AB$  is perp. to the plane  $XY$ , *Hyp.*  
 $\therefore AB$  is also perp. to  $BD$  and  $BE$ , which meet it in that plane; xi. *Def.* 1.

that is, the  $\angle^s ABD, ABE$  are rt. angles.  
 Similarly the  $\angle^s CDB, CDE$  are rt. angles.

Now in the  $\triangle^s ABD, EDB$ ,  
 because  $AB, BD = ED, DB$ , respectively, *Constr.*  
 and the  $\angle ABD =$  the  $\angle EDB$ , being rt. angles ;  
 $\therefore AD = EB$ . I. 4.

Again in the  $\triangle^s ABE, EDA$ ,  
 because  $AB, BE = ED, DA$ , respectively,  
 and  $AE$  is common ;  
 $\therefore$  the  $\angle ABE =$  the  $\angle EDA$ . I. 8.

\*NOTE. In order to shew that  $AB$  and  $CD$  are parallel, it is necessary to prove that (i) they are *in the same plane*, (ii) the angles  $ABD, CDB$ , are supplementary.

But the  $\angle ABE$  is a rt. angle ; *Proved.*  
 $\therefore$  the  $\angle EDA$  is a rt. angle.

But the  $\angle EDB$  is a rt. angle by construction,  
 and the  $\angle EDC$  is a rt. angle, since  $CD$  is perp. to the  
 plane  $XY$ . *Hyp.*

Hence  $ED$  is perp. to the three lines  $DA$ ,  $DB$ , and  $DC$  ;  
 $\therefore DA$ ,  $DB$ ,  $DC$  are in one plane. *XI. 5.*


But  $AB$  is in the plane which contains  $DA$ ,  $DB$  ; *XI. 2.*  
 $\therefore AB$ ,  $BD$ ,  $DC$  are in one plane.

And each of the  $\angle^s ABD$ ,  $CDB$  is a rt. angle ; *Hyp.*  
 $\therefore AB$  and  $CD$  are par<sup>l</sup>. *I. 28.*

*Q.E.D.*

PROPOSITION 7. THEOREM.

*If two straight lines are parallel, the straight line which joins any point in one to any point in the other is in the same plane as the parallels.*



Let  $AB$  and  $CD$  be two par<sup>l</sup> st. lines,  
 and let  $E$ ,  $F$  be any two points, one in each st. line.

*Then shall the st. line which joins  $E$ ,  $F$  be in the same plane as  $AB$ ,  $CD$ .*

For since  $AB$  and  $CD$  are par<sup>l</sup>,  
 $\therefore$  they are in one plane. *I. Def. 35.*


And since the points  $E$  and  $F$  are in this plane,  
 $\therefore$  the st. line which joins them lies wholly in this plane.  
*I. Def. 7.*

That is,  $EF$  is in the plane of the par<sup>l</sup>  $AB$ ,  $CD$ .  
*Q.E.D.*



## PROPOSITION 8. THEOREM.

*If two straight lines are parallel, and if one of them is perpendicular to a plane, then the other shall also be perpendicular to the same plane.*



Let  $AB$ ,  $CD$  be two par<sup>l</sup> st. lines, of which  $AB$  is perp. to the plane  $XY$ .

*Then  $CD$  shall also be perp. to the same plane.*

Let  $AB$  and  $CD$  meet the plane  $XY$  at the points  $B$ ,  $D$ .

Join  $BD$ ;

and in the plane  $XY$  draw  $DE$  perp. to  $BD$ , making  $DE$  equal to  $AB$ .

Join  $BE$ ,  $AE$ ,  $AD$ .

Then because  $AB$  is perp. to the plane  $XY$ , *Hyp.*  
 $\therefore AB$  is also perp. to  $BD$  and  $BE$ , which meet it in that plane; *XI. Def. 1.*

that is, the  $\angle^s$   $ABD$ ,  $ABE$  are rt. angles.

Now in the  $\triangle^s$   $ABD$ ,  $EDB$ ,

because  $AB$ ,  $BD = ED$ ,  $DB$ , respectively, *Constr.*  
 and the  $\angle$   $ABD =$  the  $\angle$   $EDB$ , being rt. angles;

$\therefore AD = EB$ .

*I. 4.*

Again in the  $\triangle^s$   $ABE$ ,  $EDA$ ,

because  $AB$ ,  $BE = ED$ ,  $DA$ , respectively,  
 and  $AE$  is common;

$\therefore$  the  $\angle$   $ABE =$  the  $\angle$   $EDA$ .

*I. 8.*

But the  $\angle ABE$  is a rt. angle ; *Proved.*

$\therefore$  the  $\angle EDA$  is a rt. angle :

that is,  $ED$  is perp. to  $DA$ .

But  $ED$  is also perp. to  $DB$  : *Constr.*

$\therefore$   $ED$  is perp. to the plane containing  $DB, DA$ . XI. 4.

And  $DC$  is in this plane ;

for both  $DB$  and  $DA$  are in the plane of the par<sup>ls</sup>  $AB, CD$ .

XI. 7.

$\therefore$   $ED$  is also perp. to  $DC$  ;

XI. *Def.* 1.

that is, the  $\angle CDE$  is a rt. angle.

Again since  $AB$  and  $CD$  are par<sup>l</sup>,  
and since the  $\angle ABD$  is a rt. angle,

*Hyp.*

$\therefore$  the  $\angle CDB$  is also a rt. angle.

I. 29.

$\therefore$   $CD$  is perp. both to  $DB$  and  $DE$  ;

$\therefore$   $CD$  is also perp. to the plane  $XY$ , which contains

$DB, DE$ .

XI. 4.

Q.E.D.

#### EXERCISES.

1. The perpendicular is the least straight line that can be drawn from an external point to a plane.

2. Equal straight lines drawn from an external point to a plane are equally inclined to the perpendicular drawn from that point to the plane.

3. Shew that two observations with a spirit-level are sufficient to determine if a plane is horizontal : and prove that for this purpose the two positions of the level must not be parallel.


4. What is the locus of points in space which are equidistant from two fixed points ?

5. Shew how to determine in a given straight line the point which is equidistant from two fixed points. When is this impossible ?

6. If a straight line is parallel to a plane, shew that any plane passing through the given straight line will have with the given plane a common section which is parallel to the given straight line.

## PROPOSITION 9. THEOREM.

*Two straight lines which are parallel to a third straight line are parallel to one another.*



Let the st. lines  $AB$ ,  $CD$  be each par<sup>l</sup> to the st. line  $PQ$ .

*Then shall  $AB$  be par<sup>l</sup> to  $CD$ .*

CASE I. If  $AB$ ,  $CD$  and  $PQ$  are *in one plane*, the proposition has already been proved. I. 30.

CASE II. But if  $AB$ ,  $CD$  and  $PQ$  are not in one plane,  
in  $PQ$  take any point  $G$  ;  
and from  $G$ , in the plane of the par<sup>l</sup>  $AB$ ,  $PQ$ , draw  $GH$   
perp. to  $PQ$  ; I. 11.  
also from  $G$ , in the plane of the par<sup>l</sup>  $CD$ ,  $PQ$ , draw  $GK$   
perp. to  $PQ$ . I. 11.

Then because  $PQ$  is perp. to  $GH$  and  $GK$ , *Constr.*  
 $\therefore PQ$  is perp. to the plane  $HGK$ , which contains them.

XI. 4.

But  $AB$  is par<sup>l</sup> to  $PQ$  ;

*Hyp.*

$\therefore AB$  is also perp. to the plane  $HGK$ .

XI. 8.


Similarly,  $CD$  is perp. to the plane  $HGK$ .

Hence  $AB$  and  $CD$ , being perp. to the same plane, are par<sup>l</sup>  
to one another. XI. 6.

Q.E.D.

PROPOSITION 10. THEOREM.

*If two intersecting straight lines are respectively parallel to two other intersecting straight lines not in the same plane with them, then the first pair and the second pair shall contain equal angles.*



Let the st. lines AB, BC be respectively par<sup>l</sup> to the st. lines DE, EF, which are not in the same plane with them.

*Then shall the  $\angle ABC =$  the  $\angle DEF$ .*

In BA and ED, make BA equal to ED ;  
and in BC and EF, make BC equal to EF.  
Join AD, BE, CF, AC, DF.

Then because BA is equal and par<sup>l</sup> to ED,

*Hyp. and Constr.*  
 $\therefore$  AD is equal and par<sup>l</sup> to BE. I. 33.

And because BC is equal and par<sup>l</sup> to EF,  
 $\therefore$  CF is equal and par<sup>l</sup> to BE. I. 33.

Hence AD and CF, being each equal and par<sup>l</sup> to BE, are equal and par<sup>l</sup> to one another ; *Ax. 1 and XI. 9.*


hence it follows that AC is equal and par<sup>l</sup> to DF. I. 33.

Then in the  $\triangle ABC, DEF$ ,  
because AB, BC, AC = DE, EF, DF, respectively,  
 $\therefore$  the  $\angle ABC =$  the  $\angle DEF$ . I. 8.

**Q.E.D.**

## PROPOSITION 11. PROBLEM.

To draw a straight line perpendicular to a given plane from a given point outside it.



Let A be the given point outside the plane XY.

It is required to draw from A a st. line perp. to the plane XY.

Draw any st. line BC in the plane XY ;

and from A draw AD perp. to BC. I. 12.

Then if AD is also perp. to the plane XY, what was required is done.

But if not, from D draw DE in the plane XY perp. to BC ;

and from A draw AF perp. to DE. I. 12.

Then AF shall be perp. to the plane XY.

Through F draw FH par<sup>l</sup> to BC. I. 31.

Now because CD is perp. to DA and DE, Constr.

$\therefore$  CD is perp. to the plane containing DA, DE. XI. 4.

And HF is par<sup>l</sup> to CD ;

$\therefore$  HF is also perp. to the plane containing DA, DE. XI. 8.

And since FA meets HF in this plane,

$\therefore$  the  $\angle$  HFA is a rt. angle ; XI. Def. 1.

that is, AF is perp. to FH.


And AF is also perp. to DE ; Constr.

$\therefore$  AF is perp. to the plane containing FH, DE ;

that is, AF is perp. to the plane XY. Q.E.F.

PROPOSITION 12. PROBLEM.

To draw a straight line perpendicular to a given plane from a given point in the plane.



Let A be the given point in the plane XY.

It is required to draw from A a st. line perp. to the plane XY.

From any point B outside the plane XY draw BC perp. to the plane. XI. 11.

Then if BC passes through A, what was required is done.

But if not, from A draw AD par<sup>l</sup> to BC. I. 31.

Then AD shall be the perpendicular required.

For since BC is perp. to the plane XY, Constr.

and since AD is par<sup>l</sup> to BC, Constr.

$\therefore$  AD is also perp. to the plane XY. XI. 8.

Q.E.F.

EXERCISES.


1. Equal straight lines drawn to meet a plane from a point without it are equally inclined to the plane.

2. Find the locus of the foot of the perpendicular drawn from a given point upon any plane which passes through a given straight line.

3. From a given point A a perpendicular AF is drawn to a plane XY; and from F, FD is drawn perpendicular to BC, any line in that plane: shew that AD is also perpendicular to BC.

## PROPOSITION 13. THEOREM.

*Only one perpendicular can be drawn to a given plane from a given point either in the plane or outside it.*



CASE I. Let the given point  $A$  be *in* the given plane  $XY$ ; and, if possible, let two perps.  $AB$ ,  $AC$  be drawn from  $A$  to the plane  $XY$ .

Let  $DF$  be the plane which contains  $AB$  and  $AC$ ; and let the st. line  $DE$  be the common section of the planes  $DF$  and  $XY$ . XI. 3

Then the st. lines  $AB$ ,  $AC$ ,  $AE$  are in one plane.

And because  $BA$  is perp. to the plane  $XY$ , *Hyp.*

$\therefore BA$  is also perp. to  $AE$ , which meets it in this plane; XI. Def. 1.

that is, the  $\angle BAE$  is a rt. angle.

Similarly, the  $\angle CAE$  is a rt. angle.

$\therefore$  the  $\angle^s BAE$ ,  $CAE$ , which are in the same plane, are equal to one another; which is impossible.

$\therefore$  two perpendiculars cannot be drawn to the plane  $XY$  from the point  $A$  in that plane.

CASE II. Let the given point  $A$  be *outside* the plane  $XY$ .

Then two perp<sup>s</sup> cannot be drawn from  $A$  to the plane;


for if there could be two, they would be par<sup>l</sup>, XI. 6.

which is absurd.

Q.E.D.

PROPOSITION 14. THEOREM.

*Planes to which the same straight line is perpendicular are parallel to one another.*



Let the st. line AB be perp. to each of the planes CD, EF.

*Then shall the planes CD, EF be par<sup>l</sup>.*

For if not, they will meet when produced.

If possible, let the two planes meet, and let the st. line GH be their common section. XI. 3.

In GH take any point K;  
and join AK, BK.

Then because AB is perp. to the plane EF,

$\therefore$  AB is also perp. to BK, which meets it in this plane; XI. Def. 1.

that is, the  $\angle$  ABK is a rt. angle.

Similarly, the  $\angle$  BAK is a rt. angle.

$\therefore$  in the  $\triangle$  KAB, the two  $\angle$ 's ABK, BAK are together equal to two rt. angles;


which is impossible. I. 17.

$\therefore$  the planes CD, EF, though produced, do not meet:  
that is, they are par<sup>l</sup>. Q.E.D.



## PROPOSITION 15. THEOREM.

*If two intersecting straight lines are parallel respectively to two other intersecting straight lines which are not in the same plane with them, then the plane containing the first pair shall be parallel to the plane containing the second pair.*



Let the st. lines  $AB$ ,  $BC$  be respectively  $\text{par}^1$  to the st. lines  $DE$ ,  $EF$ , which are not in the same plane as  $AB$ ,  $BC$ .

*Then shall the plane containing  $AB$ ,  $BC$  be  $\text{par}^2$  to the plane containing  $DE$ ,  $EF$ .*

From  $B$  draw  $BG$  perp. to the plane of  $DE$ ,  $EF$ ; XI. 11.  
and let it meet that plane at  $G$ .

Through  $G$  draw  $GH$ ,  $GK$   $\text{par}^1$  respectively to  $DE$ ,  $EF$ . I. 31.

Then because  $BG$  is perp. to the plane of  $DE$ ,  $EF$ ,  
 $\therefore$   $BG$  is also perp. to  $GH$  and  $GK$ , which meet it in that plane:

that is, each of the  $\angle^s$   $BGH$ ,  $BGK$  is a rt. angle.

Now *by hypothesis*  $BA$  is  $\text{par}^1$  to  $ED$ ,  
and *by construction*  $GH$  is  $\text{par}^1$  to  $ED$ ;

$\therefore$   $BA$  is  $\text{par}^1$  to  $GH$ .

XI. 9.

And since the  $\angle$   $BGH$  is a rt. angle;

*Proved.*

$\therefore$  the  $\angle$   $ABG$  is a rt. angle.


I. 29.

Similarly the  $\angle$   $CBG$  is a rt. angle.

Then since BG is perp. to each of the st. lines BA, BC,  
 $\therefore$  BG is perp. to the plane containing them. XI. 4.  
 But BG is also perp. to the plane of ED, EF; *Constr.*  
 that is, BG is perp. to the two planes AC, DF;  
 $\therefore$  these planes are par<sup>l</sup>. XI. 14.  
 Q.E.D.

PROPOSITION 16. THEOREM.

*If two parallel planes are cut by a third plane, their common sections with it shall be parallel.*



Let the par<sup>l</sup> planes AB, CD be cut by the plane EFHG,  
 and let the st. lines EF, GH be their common sections  
 with it.

*Then shall EF, GH be par<sup>l</sup>.*

For if not, EF and GH will meet if produced.

If possible, let them meet at K.

Then since the whole st. line EFH is in the plane AB, XI. 1.

and K is a point in that line,

$\therefore$  the point K is in the plane AB.

Similarly the point K is in the plane CD.

Hence the planes AB, CD when produced meet at K;  
 which is impossible, since they are par<sup>l</sup>. *Hyp.*


$\therefore$  the st. lines EF and GH do not meet;  
 and they are in the same plane EFHG;

$\therefore$  they are par<sup>l</sup>. I. Def. 35.

Q.E.D.

## PROPOSITION 17. THEOREM.

*Straight lines which are cut by parallel planes are cut proportionally.*



Let the st. lines  $AB, CD$  be cut by the three par<sup>l</sup> planes  $GH, KL, MN$  at the points  $A, E, B,$  and  $C, F, D.$

Then shall  $AE : EB :: CF : FD.$

Join  $AC, BD, AD;$

and let  $AD$  meet the plane  $KL$  at the point  $X:$

join  $EX, XF.$

Then because the two par<sup>l</sup> planes  $KL, MN$  are cut by the plane  $ABD,$

$\therefore$  the common sections  $EX, BD$  are par<sup>l</sup>. XI. 16.

And because the two par<sup>l</sup> planes  $GH, KL$  are cut by the plane  $DAC,$

$\therefore$  the common sections  $XF, AC$  are par<sup>l</sup>. XI. 16.

Now since  $EX$  is par<sup>l</sup> to  $BD,$  a side of the  $\triangle ABD,$

$\therefore AE : EB :: AX : XD.$  VI. 2.

Again because  $XF$  is par<sup>l</sup> to  $AC,$  a side of the  $\triangle DAC,$

$\therefore AX : XD :: CF : FD.$  VI. 2.


Hence  $AE : EB :: CF : FD.$  V. 1.

Q.E.D.

DEFINITION. One plane is perpendicular to another plane, when *any* straight line drawn in one of the planes perpendicular to their common section is also perpendicular to the other plane. [Book XI. Def. 6.]

PROPOSITION 18. THEOREM.

*If a straight line is perpendicular to a plane, then every plane which passes through the straight line is also perpendicular to the given plane.*



Let the st. line AB be perp. to the plane XY ;  
and let DE be any plane passing through AB.

*Then shall the plane DE be perp. to the plane XY.*

Let the st. line CE be the common section of the planes XY, DE. XI. 3.

From F, any point in CE, draw FG in the plane DE perp. to CE. I. 11.

Then because AB is perp. to the plane XY, Hyp.  
 $\therefore$  AB is also perp. to CE, which meets it in that plane, XI. Def. 1.

that is, the  $\angle$  ABF is a rt. angle.

But the  $\angle$  GFB is also a rt. angle ;

$\therefore$  GF is par<sup>l</sup> to AB.

*Constr.*

I. 28.

And AB is perp. to the plane XY,

*Hyp.*

$\therefore$  GF is also perp. to the plane XY.

XI. 8.

Hence it has been shewn that any st. line GF drawn in the plane DE perp. to the common section CE is also perp. to the plane XY.

$\therefore$  the plane DE is perp. to the plane XY. XI. Def. 6.


Q.E.D.

EXERCISE.

*Shew that two planes are perpendicular to one another when the dihedral angle [see XI. Def. 7] formed by them is a right angle.*

## PROPOSITION 19. THEOREM.

*If two intersecting planes are each perpendicular to a third plane, their common section shall also be perpendicular to that plane.*



Let each of the planes AB, BC be perp. to the plane ADC, and let BD be their common section.

*Then shall BD be perp. to the plane ADC.*

For if not, from D draw in the plane AB the st. line DE perp. to AD, the common section of the planes ADB, ADC :

I. 11.

and from D draw in the plane BC the st. line DF perp. to DC, the common section of the planes BDC, ADC.

Then because the plane BA is perp. to the plane ADC,

*Hyp.*

and DE is drawn in the plane BA perp. to AD the common section of these planes,

*Constr.*

$\therefore$  DE is perp. to the plane ADC. XI. Def. 6.

Similarly DF is perp. to the plane ADC.

$\therefore$  from the point D two st. lines are drawn perp. to the plane ADC ; which is impossible.


XI. 13.

Hence DB cannot be otherwise than perp. to the plane ADC.

Q.E.D.

PROPOSITION 20. THEOREM.

*Of the three plane angles which form a trihedral angle, any two are together greater than the third.*



Let the trihedral angle at A be formed by the three plane  $\angle^s$  BAD, DAC, BAC.

*Then shall any two of them, such as the  $\angle^s$  BAD, DAC, be together greater than the third, the  $\angle$  BAC.*

CASE I. If the  $\angle$  BAC is less than, or equal to, either of the  $\angle^s$  BAD, DAC ; it is evident that the  $\angle^s$  BAD, DAC are together greater than the  $\angle$  BAC.

CASE II. But if the  $\angle$  BAC is greater than either of the  $\angle^s$  BAD, DAC ; then at the point A in the plane BAC make the  $\angle$  BAE equal to the  $\angle$  BAD ;

and cut off AE equal to AD.

Through E, and in the plane BAC, draw the st. line BEC cutting AB, AC at B and C :

join DB, DC.

Then in the  $\triangle^s$  BAD, BAE,  
since BA, AD = BA, AE, respectively,  
and the  $\angle$  BAD = the  $\angle$  BAE ;

*Constr.*

*Constr.*

$\therefore$  BD = BE.

I. 4.


Again in the  $\triangle$  BDC, since BD, DC are together greater than BC,

I. 20.

and BD = BE,

*Proved.*


$\therefore$  DC is greater than EC.



And in the  $\triangle^s$  DAC, EAC,  
 because DA, AC = EA, AC respectively,  
 but DC is greater than EC ;  
 $\therefore$  the  $\angle$  DAC is greater than the  $\angle$  EAC. *Constr.*  
*Proved.*  
I. 25.  
 But the  $\angle$  BAD = the  $\angle$  BAE ; *Constr.*  
 $\therefore$  the two  $\angle^s$  BAD, DAC are together greater than the  
 $\angle$  BAC. Q.E.D.

PROPOSITION 21. THEOREM.

*Every (convex) solid angle is formed by plane angles which are together less than four right angles.*



Let the solid angle at S be formed by the plane  $\angle^s$  ASB, BSC, CSD, DSE, ESA.  
 Then shall the sum of these plane angles be less than four  
 rt. angles.

For let a plane  $XY$  intersect all the arms of the plane angles on the same side of the vertex at the points  $A, B, C, D, E$ : and let  $AB, BC, CD, DE, EA$  be the common sections of the plane  $XY$  with the planes of the several angles.

Within the polygon  $ABCDE$  take any point  $O$ ;  
and join  $O$  to each of the vertices of the polygon.

Then since the  $\angle^s$   $SAE, SAB, EAB$  form the trihedral angle  $A$ ,

$\therefore$  the  $\angle^s$   $SAE, SAB$  are together greater than the  $\angle$   $EAB$ ;

XI. 20.

that is,

the  $\angle^s$   $SAE, SAB$  are together greater than the  $\angle^s$   $OAE, OAB$ .

Similarly,

the  $\angle^s$   $SBA, SBC$  are together greater than the  $\angle^s$   $OBA, OBC$ :  
and so on, for each of the angular points of the polygon.

Thus by addition,  
the sum of the base angles of the triangles whose vertices are at  $S$ , is greater than the sum of the base angles of the triangles whose vertices are at  $O$ .

But these two systems of triangles are equal in number;  
 $\therefore$  the sum of all the angles of the one system is equal to the sum of all the angles of the other.

It follows that the sum of the vertical angles at  $S$  is less than the sum of the vertical angles at  $O$ .

But the sum of the angles at  $O$  is four rt. angles;  
 $\therefore$  the sum of the angles at  $S$  is less than four rt. angles.

Q.E.D.

NOTE. This proposition was not given in this form by Euclid, who established its truth only in the case of *trihedral* angles. The above demonstration, however, applies to all cases in which the polygon  $ABCDE$  is *convex*, but it must be observed that without this condition the proposition is not necessarily true.

A solid angle is *convex* when it lies entirely on one side of each of the infinite planes which pass through its plane angles. If this is the case, the polygon  $ABCDE$  will have no *re-entrant* angle. And it is clear that it would not be possible to apply XI. 20 to a vertex at which a re-entrant angle existed.



## EXERCISES ON BOOK XI.

1. Equal straight lines drawn to a plane from a point without it have equal projections on that plane.
2. If  $S$  is the centre of the circle circumscribed about the triangle  $ABC$ , and if  $SP$  is drawn perpendicular to the plane of the triangle, shew that any point in  $SP$  is equidistant from the vertices of the triangle.
3. Find the locus of points in space equidistant from three given points.
4. From Example 2 deduce a practical method of drawing a perpendicular from a given point to a plane, having given ruler, compasses, and a straight rod longer than the required perpendicular.
5. Give a geometrical construction for drawing a straight line equally inclined to three straight lines which meet in a point, but are not in the same plane.
6. In a *gauche* quadrilateral (that is, a quadrilateral whose sides are not in the same plane) if the middle points of adjacent sides are joined, the figure thus formed is a parallelogram.
7.  $AB$  and  $AC$  are two straight lines intersecting at right angles, and from  $B$  a perpendicular  $BD$  is drawn to the plane in which they are: shew that  $AD$  is perpendicular to  $AC$ .
8. If two intersecting planes are cut by two parallel planes, the lines of section of the first pair with each of the second pair contain equal angles.
9. If a straight line is parallel to a plane, shew that any plane passing through the given straight line will intersect the given plane in a line of section which is parallel to the given line.
10. Two intersecting planes pass one through each of two parallel straight lines; shew that the common section of the planes is parallel to the given lines.
11. If a straight line is parallel to each of two intersecting planes, it is also parallel to the common section of the planes.
12. Through a given point in space draw a straight line to intersect each of two given straight lines which are not in the same plane.
13. If  $AB$ ,  $BC$ ,  $CD$  are straight lines not all in one plane, shew that a plane which passes through the middle point of each one of them is parallel both to  $AC$  and  $BD$ .
14. From a given point  $A$  a perpendicular  $AB$  is drawn to a plane  $XY$ ; and a second perpendicular  $AE$  is drawn to a straight line  $CD$  in the plane  $XY$ : shew that  $EB$  is perpendicular to  $CD$ .

15. From a point  $A$  two perpendiculars  $AP$ ,  $AQ$  are drawn one to each of two intersecting planes: shew that the common section of these planes is perpendicular to the plane of  $AP$ ,  $AQ$ .

16. From  $A$ , a point in one of two given intersecting planes,  $AP$  is drawn perpendicular to the first plane, and  $AQ$  perpendicular to the second: if these perpendiculars meet the second plane at  $P$  and  $Q$ , shew that  $PQ$  is perpendicular to the common section of the two planes.

17.  $A$ ,  $B$ ,  $C$ ,  $D$  are four points not in one plane, shew that the four angles of the *gauche* quadrilateral  $ABCD$  [see Ex. 6, p. 444] are together less than four right angles.

18.  $OA$ ,  $OB$ ,  $OC$  are three straight lines drawn from a given point  $O$  not in the same plane, and  $OX$  is another straight line within the solid angle formed by  $OA$ ,  $OB$ ,  $OC$ : shew that

(i) the sum of the angles  $AOX$ ,  $BOX$ ,  $COX$  is greater than half the sum of the angles  $AOB$ ,  $BOC$ ,  $COA$ .

(ii) the sum of the angles  $AOX$ ,  $COX$  is less than the sum of the angles  $AOB$ ,  $COB$ .

(iii) the sum of the angles  $AOX$ ,  $BOX$ ,  $COX$  is less than the sum of the angles  $AOB$ ,  $BOC$ ,  $COA$ .

19.  $OA$ ,  $OB$ ,  $OC$  are three straight lines forming a solid angle at  $O$ , and  $OX$  bisects the plane angle  $AOB$ : shew that the angle  $XOC$  is less than half the sum of the angles  $AOC$ ,  $BOC$ .

20. If a point is equidistant from the angles of a right-angled triangle and not in the plane of the triangle, the line joining it with the middle point of the hypotenuse is perpendicular to the plane of the triangle.

21. The angle which a straight line makes with its projection on a plane is less than that which it makes with any other straight line which meets it in that plane.

22. Find a point in a given plane such that the sum of its distances from two given points (not in the plane but on the same side of it) may be a minimum.

23. If two straight lines in one plane are equally inclined to another plane, they will be equally inclined to the common section of these planes.

24.  $PA$ ,  $PB$ ,  $PC$  are three concurrent straight lines, each of which is at right angles to the other two:  $PX$ ,  $PY$ ,  $PZ$  are perpendiculars drawn from  $P$  to  $BC$ ,  $CA$ ,  $AB$  respectively. Shew that  $XYZ$  is the pedal triangle of the triangle  $ABC$ .

25.  $PA$ ,  $PB$ ,  $PC$  are three concurrent straight lines, each of which is at right angles to the other two, and from  $P$  a perpendicular  $PO$  is drawn to the plane of  $ABC$ : shew that  $O$  is the orthocentre of the triangle  $ABC$ .


## THEOREMS AND EXAMPLES ON BOOK XI.

## DEFINITIONS.

(i) Lines which are drawn on a plane, or through which a plane may be made to pass, are said to be **co-planar**.

(ii) The **projection of a line on a plane** is the locus of the feet of perpendiculars drawn from all points in the given line to the plane.

**THEOREM 1.** *The projection of a straight line on a plane is itself a straight line.*



Let  $AB$  be the given st. line, and  $XY$  the given plane.

From  $P$ , any point in  $AB$ , draw  $Pp$  perp. to the plane  $XY$ .

*It is required to shew that the locus of  $p$  is a st. line.*

From  $A$  and  $B$  draw  $Aa$ ,  $Bb$  perp. to the plane  $XY$ .

Now since  $Aa$ ,  $Pp$ ,  $Bb$  are all perp. to the plane  $XY$ ,

$\therefore$  they are par<sup>l</sup>.

xi. 6.

And since these par<sup>l</sup>s all intersect  $AB$ ,

$\therefore$  they are co-planar.

xi. 7.


$\therefore$  the point  $p$  is in the common section of the planes  $Ab$ ,  $XY$ ;  
that is,  $p$  is in the st. line  $ab$ .

But  $p$  is any point in the projection of  $AB$ ,

$\therefore$  the projection of  $AB$  is the st. line  $ab$ .

Q. E. D.

**THEOREM 2.** Draw a perpendicular to each of two straight lines which are not in the same plane. Prove that this perpendicular is the shortest distance between the two lines.



Let  $AB$  and  $CD$  be the two straight lines, not in the same plane.

(i) *It is required to draw a st. line perp. to each of them.*

Through  $E$ , any point in  $AB$ , draw  $EF$  par<sup>l</sup> to  $CD$ .

Let  $XY$  be the plane which passes through  $AB$ ,  $EF$ .

From  $H$ , any point in  $CD$ , draw  $HK$  perp. to the plane  $XY$ . xi. 11.

And through  $K$ , draw  $KQ$  par<sup>l</sup> to  $EF$ , cutting  $AB$  at  $Q$ .

Then  $KQ$  is also par<sup>l</sup> to  $CD$ ;

xi. 9.

and  $CD$ ,  $HK$ ,  $KQ$  are in one plane.

xi. 7.

From  $Q$ , draw  $QP$  par<sup>l</sup> to  $HK$  to meet  $CD$  at  $P$ .

Then shall  $PQ$  be perp. to both  $AB$  and  $CD$ .

For, since  $HK$  is perp. to the plane  $XY$ , and  $PQ$  is par<sup>l</sup> to  $HK$ ,

$\therefore PQ$  is perp. to the plane  $XY$ ;

Constr.

xi. 8.

$\therefore PQ$  is perp. to  $AB$ , which meets it in that plane. xi. Def. 1.

For a similar reason  $PQ$  is perp. to  $CD$ ,

$\therefore PQ$  is also perp. to  $CD$ , which is par<sup>l</sup> to  $KQ$ .

(ii) *It is required to shew that  $PQ$  is the least of all st. lines drawn from  $AB$  to  $CD$ .*

Take  $HE$ , any other st. line drawn from  $AB$  to  $CD$ .

Then  $HE$ , being oblique to the plane  $XY$ , is greater than the perp.  $HK$ .

Ex. 1, p. 429.


$\therefore HE$  is also greater than  $PQ$ .

Q. E. D.

**DEFINITION.** A parallelepiped is a solid figure bounded by three pairs of parallel plane faces.

**THEOREM 3.** (i) *The faces of a parallelepiped are parallelograms, of which those which are opposite are identically equal.*

(ii) *The four diagonals of a parallelepiped are concurrent and bisect one another.*



Let  $ABA'B'$  be a  $\text{par}^{\text{ped}}$ , of which  $ABCD$ ,  $C'D'A'B'$  are opposite faces.

(i) *Then all the faces shall be  $\text{par}^{\text{m}}$ , and the opposite faces shall be identically equal.*

For since the planes  $DA'$ ,  $AD'$  are  $\text{par}^{\text{l}}$ , XI. Def. 15.  
and the plane  $DB$  meets them,

$\therefore$  the common sections  $AB$  and  $DC$  are  $\text{par}^{\text{l}}$ . XI. 16.

Similarly  $AD$  and  $BC$  are  $\text{par}^{\text{l}}$ .

$\therefore$  the fig.  $ABCD$  is a  $\text{par}^{\text{m}}$ ,

and  $AB = DC$ ; also  $AD = BC$ .

I. 34.

Similarly each of the faces of the  $\text{par}^{\text{ped}}$  is a  $\text{par}^{\text{m}}$ ;

so that the edges  $AB$ ,  $C'D'$ ,  $B'A'$ ,  $DC$  are equal and  $\text{par}^{\text{l}}$ ;

so also are the edges  $AD$ ,  $C'B'$ ,  $D'A'$ ,  $BC$ ; and likewise  $AC'$ ,  $BD'$ ,  $CA'$ ,  $DB'$ .

Then in the opp. faces  $ABCD$ ,  $C'D'A'B'$ ,

we have  $AB = C'D'$  and  $BC = D'A'$ ;

*Proved.*

and since  $AB$ ,  $BC$  are respectively  $\text{par}^{\text{l}}$  to  $C'D'$ ,  $D'A'$ ,

$\therefore$  the  $\angle ABC = \text{the } \angle C'D'A'$ ;

XI. 10.


$\therefore$  the  $\text{par}^{\text{m}} ABCD = \text{the } \text{par}^{\text{m}} C'D'A'B'$  identically. Ex. 11, p. 70.

(ii) *The diagonals  $AA'$ ,  $BB'$ ,  $CC'$ ,  $DD'$  shall be concurrent and bisect one another.*

Join  $AC$  and  $A'C'$ .

Then since  $AC'$  is equal and  $\text{par}^1$  to  $A'C$ ,  
 $\therefore$  the fig.  $ACA'C'$  is a  $\text{par}^m$ ;  
 $\therefore$  its diagonals  $AA'$ ,  $CC'$  bisect one another. Ex. 5, p. 70.  
 That is,  $AA'$  passes through  $O$ , the middle point of  $CC'$ .  
 Similarly if  $BC'$  and  $B'C$  were joined, the fig.  $BCB'C'$  would be a  
 $\text{par}^m$ ;  
 $\therefore$  the diagonals  $BB'$ ,  $CC'$  bisect one another.  
 That is,  $BB'$  also passes through  $O$  the middle point of  $CC'$ .  
 Similarly it may be shewn that  $DD'$  passes through, and is  
 bisected at,  $O$ . Q. E. D.

**THEOREM 4.** *The straight lines which join the vertices of a tetrahedron to the centroids of the opposite faces are concurrent.*



Let  $ABCD$  be a tetrahedron, and let  $g_1, g_2, g_3, g_4$  be the centroids of the faces opposite respectively to  $A, B, C, D$ .

*Then shall  $Ag_1, Bg_2, Cg_3, Dg_4$  be concurrent.*

Take  $X$  the middle point of the edge  $CD$ ;  
 then  $g_1$  and  $g_2$  must lie respectively in  $BX$  and  $AX$ ,  
 so that  $BX = 3 \cdot Xg_1$ , Ex. 4, p. 113.  
 and  $AX = 3 \cdot Xg_2$ ;  
 $\therefore g_1g_2$  is  $\text{par}^1$  to  $AB$ .

And  $Ag_1, Bg_2$  must intersect one another, since they are both in the plane of the  $\triangle AXB$ ;

let them intersect at the point  $G$ .

Then by similar  $\triangle^s$ ,  $AG : Gg_1 = AB : g_1g_2$   
 $= AX : Xg_2$   
 $= 3 : 1$ .


$\therefore Bg_2$  cuts  $Ag_1$  at a point  $G$  whose distance from  $g_1 = \frac{1}{4} \cdot Ag_1$ .

Similarly it may be shewn that  $Cg_3$  and  $Dg_4$  cut  $Ag_1$  at the same point;

$\therefore$  these lines are concurrent. Q. E. D.

**THEOREM 5.** (i) *If a pyramid is cut by planes drawn parallel to its base, the sections are similar to the base.*

(ii) *The areas of such sections are in the duplicate ratio of their perpendicular distances from the vertex.*



Let  $SABCD$  be a pyramid, and  $abcd$  the section formed by a plane drawn  $\text{par}^1$  to the base  $ABCD$ .

(i) *Then the figs.  $ABCD$ ,  $abcd$  shall be similar.*

Because the planes  $abcd$ ,  $ABCD$  are  $\text{par}^1$ ,  
and the plane  $ABba$  meets them,  
 $\therefore$  the common sections  $ab$ ,  $AB$  are  $\text{par}^1$ .

Similarly  $bc$  is  $\text{par}^1$  to  $BC$ ;  $cd$  to  $CD$ ; and  $da$  to  $DA$ .

And since  $ab$ ,  $bc$  are respectively  $\text{par}^1$  to  $AB$ ,  $BC$ ,

$\therefore$  the  $\angle abc = \text{the } \angle ABC$ .

XI. 10.

Similarly the remaining angles of the fig.  $abcd$  are equal to the corresponding angles of the fig.  $ABCD$ .

And since the  $\triangle^s Sab$ ,  $SAB$  are similar,

$\therefore ab : AB = Sb : SB$

$= bc : BC$ , for the  $\triangle^s Sbc$ ,  $SBC$  are similar.

Or,  $ab : bc = AB : BC$ .

In like manner,  $bc : cd = BC : CD$ ; and so on.

$\therefore$  the figs.  $abcd$ ,  $ABCD$  are equiangular to one another, and have their sides about the equal angles proportional;

$\therefore$  they are similar.

(ii) From  $S$  draw  $SxX$   $\text{perp.}$  to the  $\text{par}^1$  planes  $abcd$ ,  $ABCD$  and meeting them at  $x$  and  $X$ .

Then shall fig.  $abcd : \text{fig. } ABCD = Sx^2 : SX^2$ .

Join  $ax$ ,  $AX$ .

Then it is clear that the  $\triangle^s Sax$ ,  $SAX$  are similar.

And the fig.  $abcd : \text{fig. } ABCD = ab^2 : AB^2$

VI. 20.

$= aS^2 : AS^2$ ,

$= Sx^2 : SX^2$ .

Q. E. D.

DEFINITION. A polyhedron is *regular* when its faces are similar and equal regular polygons.

THEOREM 6. *There cannot be more than five regular polyhedra.*

This is proved by examining the number of ways in which it is possible to form a solid angle out of the plane angles of various regular polygons; bearing in mind that *three* plane angles at least are required to form a solid angle, and the sum the plane angles forming a solid angle *is less than four right angles.* XI. 21.

Suppose the faces of the regular polyhedron to be *equilateral triangles.*

Then since each angle of an equilateral triangle is  $\frac{2}{3}$  of a right angle, it follows that a solid angle may be formed (i) by *three*, (ii) by *four*, or (iii) by *five* such faces; for the sums of the plane angles would be respectively (i) two right angles, (ii)  $\frac{8}{3}$  of a right angle, (iii)  $\frac{10}{3}$  of a right angle; that is, in all three cases the sum of the plane angles would be less than four right angles.

But it is impossible to form a solid angle of *six* or more equilateral triangles, for then the sum of the plane angles would be equal to, or greater than four right angles.

Again, suppose that the faces of the polyhedron are *squares.*

(iv) Then it is clear that a solid angle could be formed of *three*, but not more than three, of such faces.

Lastly, suppose the faces are *regular pentagons.*

(v) Then, since each angle of a regular pentagon is  $\frac{6}{5}$  of a right angle, it follows that a solid angle may be formed of *three* such faces; but the sum of more than three angles of a regular pentagon is greater than four right angles.

Further, since each angle of a *regular hexagon* is equal to  $\frac{4}{3}$  of a right angle, it follows that no solid angle could be formed of such faces; for the sum of three angles of a hexagon is equal to four right angles.

Similarly, no solid angle can be formed of the angles of a polygon of more sides than six.


Thus there can be no more than *five* regular polyhedra.





## NOTE ON THE REGULAR POLYHEDRA.

(i) The polyhedron of which each solid angle is formed by *three equilateral triangles* is called a regular **tetrahedron**.

It has *four faces*,  
*four vertices*,  
*six edges*.





(ii) The polyhedron of which each solid angle is formed by *four equilateral triangles* is called a regular **octahedron**.



It has *eight faces*, *six vertices*, *twelve edges*.


(iii) The polyhedron of which each solid angle is formed by *five equilateral triangles* is called a regular **icosahedron**.





It has *twenty faces*, *twelve vertices*, *thirty edges*.

(iv) The regular polyhedron of which each solid angle is formed by *three squares* is called a **cube**.

It has *six faces*,  
*eight vertices*,  
*twelve edges*.



(v) The polyhedron of which each solid angle is formed by *three regular pentagons* is called a regular **dodecahedron**.



It has *twelve faces*, *twenty vertices*, *thirty edges*.

**THEOREM 7.** *If F denote the number of faces, E of edges, and V of vertices in any polyhedron, then will*

$$E + 2 = F + V.$$

Suppose the polyhedron to be formed by fitting together the faces in succession: suppose also that  $E_r$  denotes the number of edges, and  $V_r$  of vertices, when  $r$  faces have been placed in position, and that the polyhedron has  $n$  faces when complete.

Now when *one* face is taken there are as many vertices as edges, that is,

$$E_1 = V_1.$$

The *second* face on being adjusted has *two* vertices and *one* edge in common with the first; therefore by adding the second face we increase the number of edges by one more than the number of vertices;

$$\therefore E_2 - V_2 = 1.$$

Again, the *third* face on adjustment has *three* vertices and *two* edges in common with the former two faces; therefore on adding the third face we once more increase the number of edges by one more than the number of vertices;

$$\therefore E_3 - V_3 = 2.$$

Similarly, when all the faces but one have been placed in position,

$$E_{n-1} - V_{n-1} = n - 2.$$

But in fitting on the last face we add no new edges nor vertices;

$$\therefore E = E_{n-1}, \quad V = V_{n-1}, \quad \text{and } F = n.$$

$$\text{So that } E - V = F - 2,$$

$$\text{or, } E + 2 = F + V.$$

This is known as *Euler's Theorem*.

### MISCELLANEOUS EXAMPLES ON SOLID GEOMETRY.

1. The projections of parallel straight lines on any plane are parallel.

2. If  $ab$  and  $cd$  are the projections of two parallel straight lines  $AB$ ,  $CD$  on any plane, shew that  $AB : CD = ab : cd$ .

3. Draw two parallel planes one through each of two straight lines which do not intersect and are not parallel.

4. If two straight lines do not intersect and are not parallel, on what planes will their projections be parallel?

5. Find the locus of the middle point of a straight line of constant length whose extremities lie one on each of two non-intersecting straight lines.

6. Three points A, B, C are taken one on each of the conterminous edges of a cube: prove that the angles of the triangle ABC are all acute.

7. If a parallelepiped is cut by a plane which intersects two pairs of opposite faces, the common sections form a parallelogram.

8. The square on the diagonal of a rectangular parallelepiped is equal to the sum of the squares on the three edges conterminous with the diagonal.

9. The square on the diagonal of a cube is three times the square on one of its edges.

10. The sum of the squares on the four diagonals of a parallelepiped is equal to the sum of the squares on the twelve edges.

11. If a perpendicular is drawn from a vertex of a regular tetrahedron on its base, shew that the foot of the perpendicular will divide each median of the base in the ratio 2 : 1.

12. Prove that the perpendicular from the vertex of a regular tetrahedron upon the opposite face is three times that dropped from its foot upon any of the other faces.

13. If AP is the perpendicular drawn from the vertex of a regular tetrahedron upon the opposite face, shew that

$$3AP^2 = 2a^2,$$

where  $a$  is the length of an edge of the tetrahedron.

14. The straight lines which join the middle points of opposite edges of a tetrahedron are concurrent.

15. If a tetrahedron is cut by any plane parallel to two opposite edges, the section will be a parallelogram.

16. Prove that the shortest distance between two opposite edges of a regular tetrahedron is one half of the diagonal of the square on an edge.

17. In a tetrahedron if two pairs of opposite edges are at right angles, then the third pair will also be at right angles.

18. In a tetrahedron whose opposite edges are at right angles in pairs, the four perpendiculars drawn from the vertices to the opposite faces and the three shortest distances between opposite edges are concurrent.

19. In a tetrahedron whose opposite edges are at right angles, the sum of the squares on each pair of opposite edges is the same.

20. The sum of the squares on the edges of any tetrahedron is four times the sum of the squares on the straight lines which join the middle points of opposite edges.

21. In any tetrahedron the plane which bisects a dihedral angle divides the opposite edge into segments which are proportional to the areas of the faces meeting at that edge.

22. If the angles at one vertex of a tetrahedron are all right angles, and the opposite face is equilateral, shew that the sum of the perpendiculars dropped from any point in this face upon the other three faces is constant.

23. Shew that the polygons formed by cutting a prism by parallel planes are equal.

24. Three straight lines in space  $OA$ ,  $OB$ ,  $OC$ , are mutually at right angles, and their lengths are  $a$ ,  $b$ ,  $c$ : express the area of the triangle  $ABC$  in its simplest form.

25. Find the diagonal of a regular octahedron in terms of one of its edges.

26. Shew how to cut a cube by a plane so that the lines of section may form a regular hexagon.

27. Shew that every section of a sphere by a plane is a circle.

28. Find in terms of the length of an edge the radius of a sphere inscribed in a regular tetrahedron.

29. Find the locus of points in a given plane at which a straight line of fixed length and position subtends a right angle.

30. A fixed point  $O$  is joined to any point  $P$  in a given plane which does not contain  $O$ ; on  $OP$  a point  $Q$  is taken such that the rectangle  $OP$ ,  $OQ$  is constant: shew that  $Q$  lies on a fixed sphere.

**WORKS BY H. S. HALL, M.A., and F. H. STEVENS, M.A.**

**A SCHOOL GEOMETRY**, based on the recommendations of the Mathematical Association, and on the recent report of the Cambridge Syndicate on Geometry.

Parts I. and II.—*Part I.* Lines and Angles. Rectilinear Figures.  
*Part II.* Areas of Rectilinear Figures. Containing the substance of Euclid Book I. Crown 8vo. 1s. 6d.

Part III.—Circles. Containing the substance of Euclid Book III. 1-34, and part of Book IV. 1s.

Parts I., II., III. in one volume. 2s. 6d.

Part IV.—Squares and Rectangles. Geometrical equivalents of Certain Algebraical Formulæ. Containing the substance of Euclid Book II., and Book III. 35-37. 6d.

Parts III. and IV. in one volume. 1s. 6d.

Parts I.-IV. in one volume. 3s.

Part V.—Containing the substance of Euclid Book VI. 1s. 6d.

Parts IV. and V. in one volume. 2s.

Parts I.-V. in one volume. 4s. 6d.

Part VI.—Containing the substance of Euclid Book XI. 1-21, together with Theorems relating to the Surfaces and Volumes of the simpler Solid Figures. 2s.

**A TEXT-BOOK OF EUCLID'S ELEMENTS**, including Alternative Proofs, together with Additional Theorems and Exercises, classified and arranged. Third Edition enlarged. Books I.—VI. XI. and XII. Props. 1 and 3. Globe 8vo. 4s. 6d.

Also in parts, separately, as follows:—

Book I. - - - 1s.	Books III. and IV. - 2s.
Books I. and II. 1s. 6d.	Books III.—VI. - 3s.
Books II. and III. 2s.	Books IV.—VI. - 2s. 6d.
Books I.—III. - 2s. 6d.	Books V., VI., XI. and XII.
Books I.—IV. - 3s.	1 and 3. - - 2s. 6d.
Sewed, 2s. 6d.	Book XI. - - 1s.

*JOURNAL OF EDUCATION*—"The most complete introduction to plane geometry based on Euclid's Elements that we have yet seen."

**A KEY TO THE EXERCISES AND EXAMPLES CONTAINED IN A TEXT-BOOK OF EUCLID'S ELEMENTS.** Books I.—VI. and XI. By H. S. HALL, M.A., and F. H. STEVENS, M.A., Masters of the Military Side, Clifton College. Crown 8vo. 8s. 6d. Books I.—IV. 6s. 6d. Books VI. and XI. 3s. 6d.

**AN ELEMENTARY COURSE OF MATHEMATICS.** Comprising Arithmetic, Algebra, and Euclid. By H. S. HALL, M.A., and F. H. STEVENS, M.A. Globe 8vo. 2s. 6d.

MACMILLAN AND CO., LTD., LONDON.

**WORKS BY H. S. HALL, M.A., and S. R. KNIGHT, B.A.**

*EIGHTH EDITION. Revised and Enlarged.*

**ELEMENTARY ALGEBRA FOR SCHOOLS** (containing a chapter on Graphs). Globe 8vo. (bound in maroon-coloured cloth), 3s. 6d. With Answers (bound in green-coloured cloth, 4s. 6d.

*NATURE*.—"This is, in our opinion, the best Elementary Algebra for school use. It is the combined work of two teachers who have had considerable experience of actual school teaching . . . and so successfully grapples with difficulties which our present text-books in use, from their authors lacking such experience, ignore or slightly touch upon. . . . We confidently recommend it to mathematical teachers, who, we feel sure, will find it the best book of its kind for teaching purposes."

**ANSWERS TO EXAMPLES IN ELEMENTARY ALGEBRA.**  
Fcap. 8vo. Sewed 1s.

**SOLUTIONS OF THE EXAMPLES IN ELEMENTARY ALGEBRA FOR SCHOOLS.** Crown 8vo. 8s. 6d.

**HIGHER ALGEBRA.** A Sequel to Elementary Algebra for Schools. Fifth Edition, revised and enlarged. Crown 8vo. 7s. 6d.

*ATHENÆUM*.—"The *Elementary Algebra*, by the same authors, is a work of such exceptional merit that those acquainted with it will form high expectations of the sequel to it now issued. Nor will they be disappointed. Of the authors' *Higher Algebra* as of their *Elementary Algebra*, we unhesitatingly assert that it is by far the best work of its kind with which we are acquainted. It supplies a want much felt by teachers."

**SOLUTIONS OF THE EXAMPLES IN HIGHER ALGEBRA.** Crown 8vo. 10s. 6d.

**ALGEBRA FOR BEGINNERS.** Globe 8vo, 2s. With Answers. 2s. 6d.

*SCHOOLMASTER*.—"To teachers who have had experience of either the Elementary or the Higher Algebra it will only be necessary to say that this book is marked by the same qualities which have brought these works into such deserved repute. To those who are still in ignorance of these books, we can say that for clear, simple, and concise explanation, convenient order of subject-matter and copious and well-graduated exercises, these books have, to say the least, no superiors."

*ABERDEEN FREE PRESS*.—"To give the learner a sufficient knowledge of the ordinary algebraical processes without presenting him with too many of their difficulties requires a thorough knowledge of what is necessary for a beginner, and of what may be omitted until a more advanced stage. A very slight examination of the book shows that the authors possess this qualification in an eminent degree. . . . The definitions and explanations of symbols are full and simple. . . . An important feature of the work is the great number of problems given to show the practical applications of the science."

**MACMILLAN AND CO., LTD., LONDON.**

**WORKS BY H. S. HALL, M.A., and S. R. KNIGHT, B.A.**

**ALGEBRAICAL EXERCISES AND EXAMINATION PAPERS.** With or without Answers. Third Edition, revised and enlarged. Globe 8vo. 2s. 6d.

*IRISH TEACHERS JOURNAL*.—"We know of no better work to place in the hands of junior teachers, monitors, and senior pupils. Any person who works carefully and steadily through this book could not possibly fail in an examination of Elementary Algebra. . . . We congratulate the authors on the skill displayed in the selections of examples."

*SCHOOLMASTER*.—"We can strongly recommend the volume to teachers seeking a well-arranged series of tests in Algebra."

**ARITHMETICAL EXERCISES AND EXAMINATION PAPERS.** With an Appendix containing Questions in LOGARITHMS AND MENSURATION. With or without Answers. Third Edition, revised and enlarged. Globe 8vo. 2s. 6d.

*CAMBRIDGE REVIEW*.—"All the mathematical work these gentlemen have given to the public is of genuine worth, and these exercises are no exception to the rule. The addition of the logarithm and mensuration questions add greatly to the value."

**ELEMENTARY TRIGONOMETRY.** Third Edition. Globe 8vo. 4s. 6d.

*ACADEMY*.—"It is drawn up with that skill which shows the writers to be past masters in the art of teaching, and is quite on as high a platform for Trigonometry as their previous books on Algebra are for that subject. We have read the text with care and have thoroughly enjoyed it."

*EDUCATIONAL REVIEW*.—"On the whole it is the best elementary treatise on Trigonometry we have seen."

**SOLUTIONS OF THE EXAMPLES IN ELEMENTARY TRIGONOMETRY.** Crown 8vo. 8s. 6d.

**ALGEBRAICAL EXAMPLES** Supplementary to Hall and Knight's ALGEBRA FOR BEGINNERS and ELEMENTARY ALGEBRA (Chaps. I.-XXVII.). By H. S. HALL, M.A. With or without Answers. Globe 8vo. 2s.

**A SHORT INTRODUCTION TO GRAPHICAL ALGEBRA.**  
By H. S. HALL, M.A. Third Edition revised and enlarged.  
Globe 8vo. 1s. [Key, in the Press.]

MACMILLAN AND CO., LTD., LONDON.



**WORK BY H. S. HALL, M.A., and R. J. WOOD, B.A.**

**ALGEBRA FOR ELEMENTARY SCHOOLS.** Globe 8vo.  
Parts I., II., and III., 6*d.* each. Cloth, 8*d.* each. Answers,  
4*d.* each.

*HEAD TEACHER.*—"Any boy working conscientiously through this book will have laid a solid foundation for more advanced effort."

*TEACHERS' AID.*—"The aim of the authors is to provide an easy and well-graduated course of Algebra suited to the requirements and capacity of those who take it as a specific subject in elementary schools. The general plan and arrangement of Hall & Knight's well-known elementary text-book have been closely followed with such changes as the special needs of elementary pupils demand. It is needless to say that the book in every way is a creditable production."

*MADRAS EDUCATIONAL NEWS.*—"The examples are many and well chosen. It is well calculated to serve the purpose for which it is intended."

**WORKS BY F. H. STEVENS, M.A.**

**MENSURATION FOR BEGINNERS,** with the Rudiments  
of Geometrical Drawing. Globe 8vo. 1*s.* 6*d.*

*THE EDUCATIONAL TIMES* (March, 1897).—"A considerable amount of ground is covered, and the whole is written with rare judgment and clearness. We should judge that it would be practically impossible to produce a book better adapted to its purpose."

*THE GUARDIAN* (January 13th, 1897).—"The little book is really excellent. Mr. Stevens seems to us to have chosen just the elements that are of use in every-day life, and, in the notes and examples he has worked out, to have given sufficient illustration of right methods. Exercises are numerous and discreetly tabulated; and—a point worth noticing—a good many are set for graphic solution."

*UNIVERSITY CORRESPONDENT.*—"The main book seems to us to be admirably written and got up in attractive form, with just such differences of type as are a real help."

*EDUCATIONAL NEWS.*—"This little treatise will be found thoroughly practical . . . the explanations are clear and to the point, and the examples numerous and graded. It is worth the attention of teachers and pupil-teachers, who have this subject to take. The book is in all respects worthy of commendation."

**ELEMENTARY MENSURATION.** Globe 8vo. 3*s.* 6*d.*

*THE OXFORD MAGAZINE* (May 27th, 1896).—"Mr. Stevens seems bent on making students understand the meaning of the processes employed; and his examples are excellent. This is by far the best book extant on the subject."

*SCHOOL GUARDIAN.*—"The volume is systematically arranged, clearly and concisely written, and fully illustrated by numerous diagrams and a variety of examples and exercises. It is in every way a creditable piece of work, and worthy of commendation."

**MACMILLAN AND CO., LTD., LONDON.**



